

# Intermolecular 1,3-dipolar cycloadditions of azomethine imines

Raymond C. F. Jones,<sup>\*,a</sup> Stephen J. Hollis,<sup>b</sup> and James N. Iley<sup>\*,b</sup>

<sup>a</sup>Department of Chemistry, Loughborough University, Leics. LE11 3TU, UK

<sup>b</sup>Department of Chemistry, The Open University, Walton Hall, Milton Keynes MK7 6AA, UK

E-mail: [r.c.f.jones@lboro.ac.uk](mailto:r.c.f.jones@lboro.ac.uk)

Dedicated to Professor Lutz Tietze to mark his 65<sup>th</sup> birthday


## Abstract

Dipolar cycloadditions of azomethine imines, formed *in situ* from aldehydes and *N*<sup>1</sup>-alkyl-*N*<sup>2</sup>-acylhydrazines, with electron-deficient dipolarophiles produce pyrazolidines: mono-substituted dipolarophiles afford principally 4-substituted pyrazolidines

**Keywords:** 1,3-dipole, azomethine imine, pyrazolidine


## Introduction

As part of our research effort into the preparation of peptidomimetics containing 5-membered heterocyclic rings,<sup>1</sup> we wished to investigate the synthesis of pyrazolidines (tetrahydropyrazoles) **1**, Scheme 1.




**Scheme 1.** Retrosynthetic analysis of pyrazolidines.

Our ultimate targets were constrained mimics of the types **3** and **4** as replacements for the dipeptide unit **5**.<sup>2</sup> The incorporation of a cyclic moiety into a peptide backbone can restrict the conformational freedom of the peptide and these so-called peptidomimetics<sup>3</sup> can possess enhanced biological activity or increased selectivity towards specific receptor sites.



The pyrazolidine ring is accessible *via* 1,3-dipolar cycloaddition reaction of an azomethine imine **2** with an appropriately substituted alkene (Scheme 1),<sup>4</sup> that has the potential to generate three new chiral centers at C-3, C-4 and C-5. It is perhaps not surprising that many of the reported examples of cycloadditions involving azomethine imines have been intramolecular, allowing some control to be exercised over the regio- and stereochemistry of the product.<sup>5</sup> Azomethine imines are available from the addition to an aldehyde, **6**, of a 1,2-disubstituted hydrazine **7**,<sup>6</sup> resulting in the formation of an aminol, **8** (Scheme 2) that loses the elements of water to form the dipole.<sup>7</sup> Certain restrictions are placed on the hydrazine substituents; if  $R^2 = H$ , then the aminol eliminates water to form a stable hydrazone, and it is also usual for  $R^3$  to be an electron-withdrawing group in order to stabilize the formal negative charge on the dipole **2**.




**Scheme 2.** Formation of azomethine imines.

In order to progress towards our long-term objective, and since *intermolecular* cycloadditions with azomethine imines had been relatively little studied, we have examined the scope and limitations of the reaction, investigating the effects of varying the nature of the different components: the aldehyde, hydrazine and dipolarophile. The results of this exploratory work are described herein: our report is prompted by recent related studies using rigid cyclic azomethine ylides.<sup>8,9</sup>


## Results and Discussion

As an orienting cycloaddition, benzaldehyde (lacking  $\alpha$ -hydrogen atoms),  $N^1$ -acetyl- $N^2$ -methylhydrazine **9** (selected as meeting the criteria outlined earlier, and prepared simply from ethyl acetate and methylhydrazine<sup>10</sup>) and methyl propenoate (a simple mono-substituted dipolarophile) were reacted together (Scheme 3) in toluene at reflux for 72 h under Dean-Stark water removal.<sup>11</sup>




**Scheme 3.** Reagent: i, CH<sub>2</sub>=CHCO<sub>2</sub>Me, PhMe reflux, Dean-Stark trap.

Chromatographic purification of the crude product afforded both the *syn*- (**10**) and the *anti*- (**11**) 4-methoxycarbonylpyrazolidine diastereoisomers (17 and 22%, respectively). The regiochemistry of the products was elucidated by 2D- NMR spectroscopy, the <sup>1</sup>H–<sup>1</sup>H COSY spectrum revealing coupling between the methine protons, showing them to be adjacent (C-3 and C-4). The relative stereochemistry of the *anti* isomer **11** was deduced by n.O.e. experiments and confirmed by an X-ray crystal structure determination (Fig. 1).



**Figure 1.** X-Ray crystal structure of *anti*-tetrahydropyrazolidine **11**.

Whereas the *syn*-isomer **10** was obtained pure following column chromatography, preparative HPLC was necessary to separate the *anti*-isomer **11** from a trace impurity. Although it was not possible to isolate this latter material in pure form, it was tentatively assigned as the 5-methoxycarbonylpyrazolidine **12** based on its signals in the <sup>1</sup>H NMR spectrum of the mixture. Also isolated from the reaction mixture were the trisubstituted hydrazine **13** (33%; incompletely characterized), formed by conjugate addition of the hydrazine to methyl propenoate,<sup>8</sup> and the hexahydrotetrazine **14** (27%) presumably arising from dimerization of the dipole, a reaction that has been observed by others.<sup>6,7,12</sup> In an attempt to improve the efficiency of the reaction and minimize the formation of these unwanted by-products, the effect of altering the reaction conditions was investigated.




Although the use of a Dean-Stark trap would appear necessary to promote formation of the dipole, cycloadditions of azomethine imines have been carried out in the presence of a molar equivalent of water.<sup>8</sup> When the reaction was attempted either without water removal or with one molar equivalent of water added, a reduction in yield of the cycloadducts **10** and **11** and the hexahydrotetrazine **14** was observed, suggesting that dipole formation had been inhibited. A significant proportion of the azomethine imine dimerizes to form **14** in preference to cycloaddition. Triethylamine has been reported as an effective suppressant of dipole dimerization during the cycloaddition of azomethine imines;<sup>13</sup> however, a reaction performed in the presence of 3 molar equivalents of base did not result in a significant drop in the quantity of dimer **14** formed. Lowering the reaction temperature to that of benzene at reflux, or to ambient temperature (using 4Å molecular sieves to remove water), gave reduced yields of **10** and **11**. To minimize the conjugate addition (to form **13**) that competes with dipole formation, an extra molar equivalent of aldehyde was used but this resulted in only a slight increase in the yields of **10** and **11**. Increasing the amount of hydrazine present so that complete conversion of the aldehyde to the dipole could take place despite any conjugate addition likewise produced a similar, marginal improvement. We were thus unable to improve significantly upon the initial yield, and so turned our attention towards the reaction of the azomethine imine with other dipolarophiles (Scheme 4).

Cycloaddition of the azomethine imine with acrylonitrile gave the *syn*- and *anti*- 4-cyanotetrahydropyrazoles **15** (15%) and **16** (10%) respectively. The regiochemistry of the products was again determined by 2D NMR spectroscopy. The relative stereochemistry was assigned by comparison with the <sup>1</sup>H NMR spectra of the 4-methoxycarbonyltetrahydropyrazoles **10** and **11**, for which C-4(H) resonates at δ 3.94 for the *syn*-isomer **10** and δ 3.45 for **11**. The corresponding values for cycloadducts **15** and **16** are δ 3.78 and δ 3.44, respectively. In addition, the coupling constant <sup>3</sup>J<sub>3,4</sub> is larger in isomer **15** than in **16** (6.6 Hz compared to 6.0 Hz) which compares favorably with the values for **10** and **11** (7.7 Hz and 5.9 Hz, respectively). The *anti*-isomer **16** was not isolated in pure form as it contained traces of an impurity (assigned from the <sup>1</sup>H NMR spectrum of the mixture as the 5-substituted pyrazolidine) inseparable by chromatography.


When the symmetrical 1,2-disubstituted dipolarophile, dimethyl fumarate was used, the *anti*, *anti*-product **17** (38%; stereochemistry determined by n.O.e. experiments) was isolated along with an inseparable mixture of further **17** (21%) and an isomer tentatively assigned from the <sup>1</sup>H NMR spectrum as the *syn*, *anti*-product **18** (6%). Dimethyl maleate as dipolarophile afforded

only one pure isolated product (11%), all-*syn* **19**, assigned on the basis of n.O.e. experiments. A mixture of further unidentified cycloaddition products (22%) was also obtained. Only two diastereoisomers, the all-*syn* **19** and the *anti*, *syn*-isomer (having both ester substituents *anti*- to the phenyl substituent) can arise from concerted cycloaddition. However, since at least three different cycloadducts appeared to be formed, this suggests some conversion of dimethyl maleate into the more stable fumarate before cycloaddition. Reaction of the dipole with *N*-phenylmaleimide, usually a very reactive dipolarophile, resulted in isolation of one cycloadduct, the *anti*, *syn*-isomer **20** (stereochemistry again determined by n.O.e. studies) in a disappointing yield of 4%.




**Scheme 4.** Reagents: i,  $\text{CH}_2=\text{CHCN}$ ; ii, *E*- $\text{MeO}_2\text{CCH}=\text{CHCO}_2\text{Me}$ ; iii, *Z*- $\text{MeO}_2\text{CCH}=\text{CHCO}_2\text{Me}$ ; iv, *N*-phenylmaleimide (PhMe reflux, Dean-Stark trap).

We next examined the effect of altering the aldehyde. Replacing benzaldehyde by 2-methylpropanal (heating only for 16 h) gave the 4-substituted pyrazolidines **21** and **22**, in 25% and 11% yield, respectively (Scheme 5). The regiochemistry was determined by 2D NMR spectroscopy, the stereochemistry by comparison with data for **10** and **11**.<sup>14</sup> Also isolated was the 5-substituted isomer **23** in 1% yield (stereochemistry undetermined).



**Scheme 5.** Reagent (i),  $\text{CH}_2=\text{CHCO}_2\text{Me}$ , PhMe reflux, Dean-Stark trap.


Cycloaddition is thus possible when the aldehyde bears a simple alkyl or aryl group. Next we examined the use of more functionalized aldehydes, particularly those that would offer the exocyclic amino functionality present in the tetrahydropyrazoles **3** and **4**. Subjecting the Boc-protected  $\alpha$ -amino aldehyde **24** (synthesized from *S*-alanine<sup>1a,15</sup>) to the normal cycloaddition conditions with methyl acrylate did not give the expected adducts, but instead gave cyclization involving the carbamate functionality to afford the dihydrooxazole **25** (27%) (Scheme 6). The *N,N*-dibenzylated aldehyde **26** (prepared by tribenylation of *S*-alanine with PhCH<sub>2</sub>Br–K<sub>2</sub>CO<sub>3</sub>, reduction of the benzyl ester using LiAlH<sub>4</sub>, and Swern oxidation of the resulting 2-dibenzylaminopropan-1-ol<sup>16</sup>) was employed to prevent this, but instead gave rise to the 2,3-dihydro-1,3,4-oxadiazole **27** from an alternative cyclization involving the hydrazine *N*-acetyl group (Scheme 6). 2-Benzyloxypropanal **28** (prepared by benzylation of ethyl lactate using PhCH<sub>2</sub>Br–NaH and reduction of the ester with DIBAL at –50°C) gave the analogous product **29**. Products **25**, **27**, and **29** were incompletely characterized. The presence of an electronegative atom  $\alpha$ - to the dipole carbon atom thus seems to encourage formation of charge-neutral molecules **27** and **29** over reaction with the dipolarophile; this may be because of a destabilization of the dipole. Attempts to carry out cycloadditions using 2-oxopropanal (methyl glyoxal) or ethyl glyoxylate as aldehyde component were unsuccessful, the only product isolated in each case being the conjugate addition product **13** (see above).



**Scheme 6.** Reagent: i, MeNHNHCOCH<sub>3</sub> (**9**), PhMe reflux, Dean-Stark trap.


We next turned our attention to two further hydrazines. The *N*<sup>1</sup>-acetyl-*N*<sup>2</sup>-benzylhydrazine **30**, formed *via* reductive alkylation of *N*-acetylhydrazine using benzaldehyde,<sup>17</sup> afforded cycloadduct **31** in only 4% yield upon reaction with 2-methylpropanal and methyl propenoate (Scheme 7) in a sealed tube for 72 h, with molecular sieves being used to remove water. The increased steric bulk of the hydrazine substituent seemed to hinder the cycloaddition. We assume this adduct to be a 4-methoxycarbonylpyrazolidine **31**, based on the precedent above, although it

proved impossible to determine unequivocally either the regio- or stereo-chemistry of this product as the C-4 and C-5 methine protons were indistinguishable by NMR spectroscopy.



**Scheme 7.** Reagents: i, PhCHO, PhMe reflux; ii, NaBH<sub>3</sub>CN, *p*-TsOH; 1M NaOH; iii, *i*-PrCHO, CH<sub>2</sub>=CHCO<sub>2</sub>Me, molecular sieves, sealed tube.

Similarly, *N*<sup>1</sup>-*tert*-butoxycarbonyl-*N*<sup>2</sup>-methylhydrazine **32**<sup>18</sup> afforded the tetrahydropyrazole **33** (18%) as the only cycloadduct from reaction with benzaldehyde and methyl propenoate (Scheme 8). Once again the structure of this product is assumed to be that shown, although the signals for C-4(H) and one of the methylene protons could not be distinguished by NMR spectroscopy, hindering elucidation of the regio- and stereochemistry.



**Scheme 8.** Reagents: i, PhMe, reflux.

## Conclusions

The 1,3-dipolar cycloadditions of azomethine imine dipoles that have been described demonstrate that the reaction proceeds with electron-deficient alkene dipolarophiles to provide pyrazolidines; mono-substituted dipolarophiles afford predominantly 4-substituted pyrazolidines.<sup>6-8</sup> A variety of mono- and 1,2-disubstituted dipolarophiles can be used, but the cycloaddition is more sensitive to structural variations in the hydrazine and aldehyde. No cycloadditions were observed with an aldehyde bearing an  $\alpha$ -heteroatom functional group, so the only pyrazolidines accessible using this approach are those with alkyl or aryl substituents at the C-3.

## Experimental Section

**General Procedures.** Melting points were determined using a Kofler hot-stage apparatus and are uncorrected. Combustion analyses were performed by MEDAC Ltd. (Englefield Green, Surrey). Accurate mass measurements were carried out by the EPSRC National Mass Spectrometry Service Centre (University of Wales Swansea). Infrared spectra were recorded using a Perkin-Elmer 1710 FT-IR spectrometer.  $^1\text{H}$  NMR spectra were obtained at 400 MHz on a JEOL JNM-EX400 or at 300 MHz on a JEOL JNM-LA300 spectrometer.  $^{13}\text{C}$  NMR spectra were recorded at 100 MHz or at 75 MHz, respectively, on the same instruments. Low resolution mass spectra were recorded on a VG Micromass VG20-250 spectrometer or by the EPSRC National Mass Spectrometry Service Centre (University of Wales Swansea). Crystallographic measurements were carried out by the EPSRC X-Ray Crystallography Service (University of Southampton). All reagents were purified by distillation or recrystallization where appropriate, or according to standard procedures.<sup>19</sup> Anhydrous toluene was obtained commercially. Column chromatography was carried out using Fluka Silica Gel 60 (220-440 mesh). Preparative HPLC was performed using a Waters Delta Prep 3000 instrument fitted with a Jones Apex 2 silica column (5 micron, 25cm x 21mm i.d.).

***syn*- and *anti*-1-Acetyl-4-methoxycarbonyl-2-methyl-3-phenyltetrahydropyrazoles (10) and (11).** Benzaldehyde (1.20 g, 11.31 mmol), *N*<sup>1</sup>-acetyl-*N*<sup>2</sup>-methylhydrazine (**9**, 1.05 g, 11.92 mmol) and methyl propenoate (1.53 g, 17.77 mmol) in toluene (70 mL) were placed in a flask fitted with a Dean-Stark trap. After heating at reflux for 72 h, the solvent was removed using a rotary evaporator and the crude product purified by column chromatography (hexane:ethyl acetate 2:1 v/v) to give the *syn*-diastereoisomer **10** (0.50 g, 17%) as a white solid, m.p. 47-50° C. The other diastereoisomer was also isolated (0.63g, 22%) containing traces of an impurity (tentatively assigned as the 5-methoxycarbonyl isomer **12**). This was removed by further purification using normal-phase preparative HPLC (0.375% methanol in  $\text{CH}_2\text{Cl}_2$ , v/v) to give the *anti*-diastereoisomer **11** (0.17g, 6%) as a white solid, m.p. 79-81° C. Also isolated from the reaction mixture were *N*<sup>1</sup>-(2-methoxycarbonyl-ethyl)-*N*<sup>1</sup>-methyl-*N*<sup>2</sup>-acetylhydrazine **13**, as an oil (0.90 g, 33%), and *1,4*-diacetyl-2,5-dimethyl-3,6-diphenylhexahydrotetrazine **14** (0.53g, 27%), as a white solid, m.p. 80-82° C.

***syn*- Isomer (10):** Found: C, 63.99; H, 6.96; N, 10.68%;  $M^+$ , 262.1318.  $\text{C}_{14}\text{H}_{18}\text{N}_2\text{O}_3$  requires C, 64.11; H, 6.92; N, 10.67%;  $M$ , 262.1317;  $\nu_{\text{max}}$  (KBr)/ $\text{cm}^{-1}$  2955, 2380, 1728, 1646, 1437, 1382, 1316, 1213, 1134, 1094, 732 and 697;  $\delta_{\text{H}}$  (300 MHz;  $\text{CDCl}_3$ ) 2.16 (3H, s,  $\text{CH}_3\text{CO}$ ), 2.74 (3H, s,  $\text{CH}_3\text{N}$ ), 3.35 (3H, s,  $\text{CO}_2\text{CH}_3$ ), 3.76 (1H, dd,  $J$  9.7, 11.6 Hz, 5-*CHH*), 3.94 (1H, m,  $\text{CHCO}_2$ ), 4.35 (1H, d,  $J$  7.7 Hz,  $\text{C}_6\text{H}_5\text{CH}$ ), 4.45 (1H, dd,  $J$  7.5, 11.6 Hz, 5-*CHH*), 7.05-7.14 (2H, m, Ar-H) and 7.23-7.33 (3H, m, Ar-H);  $\delta_{\text{C}}$  (75 MHz;  $\text{CDCl}_3$ ) 21.08 ( $\text{CH}_3\text{CO}$ ), 44.45 (5- $\text{CH}_2$ ), 46.17 ( $\text{CH}_3\text{N}$ ), 47.84 ( $\text{CHCO}_2$ ), 51.77 ( $\text{CO}_2\text{CH}_3$ ), 72.36 ( $\text{C}_6\text{H}_5\text{CH}$ ), 126.70, 127.95 and 128.48 (Ar-CH), 137.90 (Ar-C), 170.06 and 171.45 (C=O);  $m/z$  (EI) 262 ( $M^+$ , 7%), 219 (100), 187, 159, 115, 91, 77, 57 and 43.



**anti- Isomer (11):** Found: C, 64.12; H, 6.94; N, 10.61%;  $MH^+$ , 263.1396.  $C_{14}H_{18}N_2O_3$  requires C, 64.11; H, 6.92; N, 10.67%;  $MH$ , 263.1396;  $\nu_{max}$  (KBr)/ $cm^{-1}$  3456, 2956, 2889, 1738, 1656, 1464, 1412, 1382, 1208, 1053, 726 and 701;  $\delta_H$  (300 MHz;  $CDCl_3$ ) 2.17 (3H, s,  $CH_3CO$ ), 2.63 (3H, s,  $CH_3N$ ), 3.45 (1H, m,  $CHCO_2$ ), 3.76 (3H, s,  $CO_2CH_3$ ), 3.86 (1H, m, 5-*CHH*), 4.43 (1H, m, 5-*CHH*), 4.47 (1H, d,  $J$  5.9 Hz,  $C_6H_5CH$ ) and 7.27-7.37 (5H, m, Ar-H);  $\delta_C$  (75 MHz;  $CDCl_3$ ) 20.83 ( $CH_3CO$ ), 44.51 ( $CH_3N$ ), 45.02 (5- $CH_2$ ), 50.87 ( $CHCO_2$ ), 52.62 ( $CO_2CH_3$ ), 73.51 ( $C_6H_5CH$ ), 126.68, 127.90 and 128.77 (Ar-CH), 139.86 (Ar-C), 171.05 and 172.72 (C=O);  $m/z$  (EI) 263 ( $MH^+$ , 10%), 219, 187, 159, 131, 117, 115, 91, 77, 57 and 43 (100).

**Hydrazine (13):** major rotational isomer:  $\delta_H$  (300 MHz;  $CDCl_3$ ) 2.06 (3H, s,  $CH_3CO$ ), 2.48-2.61 (2H, m,  $CH_2$ ), 2.59 (3H, s,  $CH_3N$ ), 2.86-3.07 (2H, m,  $CH_2$ ), 3.69 (3H, s,  $CO_2CH_3$ ) and 6.50 (1H, br. s, NH);  $\delta_C$  (75 MHz;  $CDCl_3$ ) 19.63 ( $CH_3CO$ ), 32.34 ( $CH_2$ ), 47.37 ( $CH_3N$ ), 51.84 ( $CH_2$ ), 55.53 ( $CO_2CH_3$ ), 172.36 and 174.84 (C=O); minor rotational isomer:  $\delta_H$  (300 MHz;  $CDCl_3$ ) 1.90 (3H, s,  $CH_3CO$ ), 2.48-2.61 (2H, m,  $CH_2$ ), 2.66 (3H, s,  $CH_3N$ ), 2.86-3.07 (2H, m,  $CH_2$ ), 3.69 (3H, s,  $CO_2CH_3$ ) and 6.54 (1H, br. s, NH);  $\delta_C$  (75 MHz;  $CDCl_3$ ) 21.48 ( $CH_3CO$ ), 32.40 ( $CH_2$ ), 46.17 ( $CH_3N$ ), 51.81 ( $CH_2$ ), 54.01 ( $CO_2CH_3$ ), 168.62 and 173.20 (C=O);  $m/z$  (EI) 174 ( $M^+$ , 1%), 143, 131, 116, 101, 59 (100) and 43.

**Tetrazine (14):** Found: C, 68.18; H, 6.87; N, 15.86%;  $MH^+$ , 353.1986.  $C_{20}H_{24}N_4O_2$  requires C, 68.16; H, 6.86; N, 15.89;  $MH$ , 353.1977;  $\nu_{max}$  (KBr)/ $cm^{-1}$  3020, 2975, 1682, 1602, 1573, 1462, 1411, 1398, 1142, 995, 762 and 698;  $\delta_H$  (300 MHz;  $CDCl_3$ ) 2.46 (6H, s,  $CH_3CO$ ), 3.34 (6H, s,  $CH_3N$ ), 7.34-7.44 (6H, m, Ar-H) and 7.59-7.72 (6H, m, Ar-H and NCHN);  $\delta_C$  (75 MHz;  $CDCl_3$ ) 21.59 ( $CH_3CO$ ), 27.47 ( $CH_3N$ ), 126.95, 128.74 and 129.62 (Ar-CH), 134.81 (Ar-C), 138.72 (NCHN) and 172.88 (C=O);  $m/z$  (EI) 176 ( $M/2^+$ , 5%), 133, 90, 89, 77, 63, 51 and 43 (100).

**Crystallographic data for (11):**  $C_{14}H_{18}N_2O_3$ ,  $M = 262.30$ , triclinic,  $a = 7.5758(3)$ ,  $b = 9.5502(4)$ ,  $c = 9.8576(3)$  Å,  $\alpha = 84.009(2)$ ,  $\beta = 86.769(2)$ ,  $\gamma = 74.377(2)^\circ$ ,  $U = 682.80(4)$  Å<sup>3</sup>,  $T = 150(2)$  K, space group  $P-1$ , monochromatic Mo-K $\alpha$  radiation,  $\lambda = 0.71073$  Å,  $Z = 2$ ,  $D_c = 1.276$  Mg m<sup>-3</sup>,  $F(000) = 280$ , colorless block, dimensions 0.4 x 0.3 x 0.3 mm,  $m(Mo-K\alpha) = 0.090$  mm<sup>-1</sup>,  $3.07 < 2\theta < 25.02^\circ$ , 6798 reflections measured, 2386 unique reflections. The structure was solved by direct methods and refined by full-matrix least-squares on  $F^2$ . The final cycle (for 245 parameters) converged with  $wR2 = 0.0912$  (for all data) and  $R1 = 0.0363$  [ $F^2 > 2s(F^2)$ ].

**syn- and anti-1-Acetyl-4-cyano-2-methyl-3-phenyltetrahydropyrazoles (15) and (16).** The procedure described for the synthesis of 1-acetyl-4-methoxycarbonyl-2-methyl-3-phenyltetrahydropyrazoles **10** and **11** was followed, but using benzaldehyde (0.94 g, 8.85 mmol),  $N^1$ -acetyl- $N^2$ -methylhydrazine **9** (0.76 g, 8.63 mmol), propenonitrile (0.93 g, 17.47 mmol) and toluene (40 mL). Column chromatography afforded the pure *syn*- diastereoisomer **15** (0.29 g, 15%), as a pale yellow solid, m.p. 150-153°C. The *anti*- diastereoisomer **16** was also isolated as a yellow solid (0.20g, 10%) containing traces of an impurity (tentatively assigned as the 5-cyano isomer).

**syn-Isomer (15):** Found: C, 68.09; H, 6.64; N, 18.06%;  $M^+$ , 229.1216.  $C_{13}H_{15}N_3O$  requires C, 68.10; H, 6.59; N, 18.32%;  $M$ , 229.1215;  $\nu_{max}$  (KBr)/ $cm^{-1}$  2998, 2964, 2895, 2246, 1656, 1470, 1453, 1444, 1414, 1373, 758 and 712;  $\delta_H$  (300 MHz;  $CDCl_3$ ) 2.21 (3H, s,  $CH_3CO$ ), 2.72 (3H, s,

CH<sub>3</sub>N), 3.78 (1H, ddd, *J* 5.6, 6.6, 8.2 Hz, CHCN), 3.87 (1H, dd, *J* 8.2, 11.4 Hz, 5-CHH), 4.32 (1H, d, *J* 6.6 Hz, C<sub>6</sub>H<sub>5</sub>CH), 4.41 (1H, dd, *J* 5.6, 11.4 Hz, 5-CHH), 7.25-7.27 (2H, m, Ar-H) and 7.31-7.43 (3H, m, Ar-H);  $\delta_{\text{C}}$  (75 MHz; CDCl<sub>3</sub>) 20.90 (CH<sub>3</sub>CO), 34.90 (CHCN), 45.83 (CH<sub>3</sub>N), 46.50 (5-CH<sub>2</sub>), 72.29 (C<sub>6</sub>H<sub>5</sub>CH), 117.07 (CN), 127.00, 128.88 and 128.94 (Ar-CH), 135.85 (Ar-C) and 172.09 (C=O); *m/z* (EI) 229 (M<sup>+</sup>, 5%), 186 (100), 159, 142, 133, 115, 91, 77, 57 and 43.

**anti-Isomer (16):** Found: C, 67.90; H, 6.57; N, 18.26%; M<sup>+</sup>, 229.1226. C<sub>13</sub>H<sub>15</sub>N<sub>3</sub>O requires C, 68.10; H, 6.59; N, 18.32%; M, 229.1215;  $\nu_{\text{max}}$  (KBr)/cm<sup>-1</sup> 3035, 2994, 2963, 2922, 2883, 2802, 2247, 1662, 1496, 1398, 1344, 1319, 1216, 1110, 768, 738 and 702;  $\delta_{\text{H}}$  (300 MHz; CDCl<sub>3</sub>) 2.14 (3H, s, CH<sub>3</sub>CO), 2.75 (3H, s, CH<sub>3</sub>N), 3.44 (1H, m, CHCN), 3.84 (1H, dd, *J* 7.1, 11.9 Hz, 5-CHH), 4.46 (1H, d, *J* 6.0 Hz, C<sub>6</sub>H<sub>5</sub>CH), 4.49 (1H, m, 5-CHH), 7.26-7.32 (2H, m, Ar-H) and 7.34-7.42 (3H, m, Ar-H);  $\delta_{\text{C}}$  (75 MHz; CDCl<sub>3</sub>) 20.82 (CH<sub>3</sub>CO), 34.96 (CHCN), 44.48 (CH<sub>3</sub>N), 46.04 (5-CH<sub>2</sub>), 74.21 (C<sub>6</sub>H<sub>5</sub>CH), 119.38 (CN), 126.52, 128.74 and 129.11 (Ar-CH), 137.26 (Ar-C) and 171.55 (C=O); *m/z* (EI) 229 (M<sup>+</sup>, 1%), 186, 143, 133, 115, 91, 77, 57 and 43 (100).

**anti, anti-1-Acetyl-4,5-bis(methoxycarbonyl)-2-methyl-3-phenyltetrahydropyrazole (17).**

The procedure described for the synthesis of 1-acetyl-4-methoxycarbonyl-2-methyl-3-phenyltetrahydropyrazoles **10** and **11** was followed, but using benzaldehyde (0.61 g, 5.71 mmol), *N*<sup>1</sup>-acetyl-*N*<sup>2</sup>-methylhydrazine **9** (0.51 g, 5.79 mmol), dimethyl fumarate (1.68 g, 11.66 mmol) and toluene (40 mL). Column chromatography afforded the *title compound 17* as a white solid (0.69 g, 38%), m.p. 152-154°C, along with a mixture (0.49 g, 27%) of the title compound (21%) and an isomer tentatively assigned as the *syn, anti*-diastereoisomer **18** (6%).

**anti, anti-Diastereoisomer (17):** Found: C, 59.89; H, 6.27; N, 8.71%; M<sup>+</sup>, 320.1368. C<sub>16</sub>H<sub>20</sub>N<sub>2</sub>O<sub>5</sub> requires C, 59.99; H, 6.29; N, 8.74%; M, 320.1372;  $\nu_{\text{max}}$  (KBr)/cm<sup>-1</sup> 2991, 2963, 2361, 1765, 1732, 1654, 1435, 1409, 1215, 764 and 708;  $\delta_{\text{H}}$  (300 MHz; CDCl<sub>3</sub>) 2.27 (6H, s, CH<sub>3</sub>CO and CH<sub>3</sub>N), 3.71 and 3.81 (each 3H, s, CO<sub>2</sub>CH<sub>3</sub>), 4.03 (1H, dd, *J* 8.3, 10.8 Hz, 4-CHCO<sub>2</sub>), 4.58 (1H, d, *J* 10.8 Hz, C<sub>6</sub>H<sub>5</sub>CH), 5.16 (1H, d, *J* 8.3 Hz, 5-CHCO<sub>2</sub>) and 7.33-7.43 (5H, m, Ar-H);  $\delta_{\text{C}}$  (75 MHz; CDCl<sub>3</sub>) 20.95 (CH<sub>3</sub>CO), 39.75 (CH<sub>3</sub>N), 49.79 (4-CHCO<sub>2</sub>), 52.84 and 52.97 (CO<sub>2</sub>CH<sub>3</sub>), 62.33 (5-CHCO<sub>2</sub>), 71.90 (C<sub>6</sub>H<sub>5</sub>CH), 128.82, 128.83 and 128.85 (Ar-CH), 132.85 (Ar-C), 170.93, 171.00 and 171.18 (C=O); *m/z* (EI) 320 (M<sup>+</sup>, 12%), 278, 245, 217, 185, 159, 115, 91, 77, 59 and 43 (100).

**syn, syn-1-Acetyl-4,5-bis(methoxycarbonyl)-2-methyl-3-phenyltetrahydropyrazole (19).**

The procedure described for the synthesis of 1-acetyl-4-methoxycarbonyl-2-methyl-3-phenyltetrahydropyrazoles **10** and **11** was followed, but using benzaldehyde (0.56 g, 5.31 mmol), *N*<sup>1</sup>-acetyl-*N*<sup>2</sup>-methylhydrazine **9** (0.47 g, 5.33 mmol), dimethyl maleate (1.66 g, 11.52 mmol) and toluene (40 mL). Column chromatography afforded an inseparable mixture of stereoisomeric tetrahydropyrazoles (0.43 g, 25%), and the *title compound 19* (0.18 g, 11%) as a white solid, m.p. 155-158°C; Found: C, 59.78; H, 6.32; N, 8.58%; M<sup>+</sup>, 320.1372. C<sub>16</sub>H<sub>20</sub>N<sub>2</sub>O<sub>5</sub> requires C, 59.99; H, 6.29; N, 8.74%; M, 320.1372;  $\nu_{\text{max}}$  (KBr)/cm<sup>-1</sup> 3007, 2953, 1765, 1747, 1641, 1494, 1455, 1433, 1377, 1356, 1342, 1198, 1168, 1110, 741 and 700;  $\delta_{\text{H}}$  (300 MHz; CDCl<sub>3</sub>) 2.33 (3H, s, CH<sub>3</sub>CO), 2.78 (3H, s, CH<sub>3</sub>N), 3.53 and 3.66 (each 3H, s, CO<sub>2</sub>CH<sub>3</sub>), 4.23 (1H, dd, *J* 7.9, 10.1 Hz, 4-CHCO<sub>2</sub>), 4.50 (1H, d, *J* 7.9 Hz, C<sub>6</sub>H<sub>5</sub>CH), 4.85 (1H, d, *J* 10.1 Hz, 5-CHCO<sub>2</sub>) and 7.24-

7.38 (5H, m, Ar-H);  $\delta_C$  (75 MHz; CDCl<sub>3</sub>) 21.18 (CH<sub>3</sub>CO), 45.23 (CH<sub>3</sub>N), 51.18 (4-CHCO<sub>2</sub>), 51.98 and 52.24 (CO<sub>2</sub>CH<sub>3</sub>), 59.00 (5-CHCO<sub>2</sub>), 71.65 (C<sub>6</sub>H<sub>5</sub>CH), 127.97, 128.01 and 128.09 (Ar-CH), 136.08 (Ar-C), 167.93, 168.74 and 169.57 (C=O);  $m/z$  (EI) 321 (MH<sup>+</sup>, 2%), 278, 219, 159, 133, 115, 91, 77, 59 and 43 (100).

**anti, syn-1-Acetyl-2-methyl-3,5-diphenyloctahydropyrrolo[3,4-c]pyrazole-4,6-dione (20).**

The procedure described for the synthesis of 1-acetyl-4-methoxycarbonyl-2-methyl-3-phenyltetrahydropyrazoles **10** and **11** was followed, but using benzaldehyde (0.63 g, 5.90 mmol), *N*<sup>1</sup>-acetyl-*N*<sup>2</sup>-methylhydrazine **9** (0.55 g, 6.24 mmol), *N*-phenylmaleimide (2.05 g, 11.84 mmol) and toluene (40 mL). Column chromatography (hexane: ethyl acetate 3:2 v/v) afforded the *title compound* **20** (0.08 g, 4%) as a yellow solid, m.p. 239-241°C; Found: C, 68.43; H, 5.44; N, 11.58%; M<sup>+</sup>, 349.1422. C<sub>20</sub>H<sub>19</sub>N<sub>3</sub>O<sub>3</sub> requires C, 68.75; H, 5.44; N, 12.02%; M, 349.1426;  $\nu_{\max}$  (KBr)/cm<sup>-1</sup> 1719, 1653, 1499, 1386, 1199, 1158, 746 and 696;  $\delta_H$  (300 MHz; CDCl<sub>3</sub>) 2.03 (3H, s, CH<sub>3</sub>CO), 2.87 (3H, s, CH<sub>3</sub>N), 4.02 (1H, d, *J* 8.6 Hz, C<sub>6</sub>H<sub>5</sub>CHCHCO), 4.73 (1H, s, C<sub>6</sub>H<sub>5</sub>CH), 5.80 (1H, d, *J* 8.6 Hz, NCHCO) and 7.18-7.53 (10H, m, Ar-H);  $\delta_C$  (75 MHz; CDCl<sub>3</sub>) 20.88 (CH<sub>3</sub>CO), 47.55 (CH<sub>3</sub>N), 53.11 (C<sub>6</sub>H<sub>5</sub>CHCHCO) 58.42 (NCHCO), 73.07 (C<sub>6</sub>H<sub>5</sub>CH), 125.82, 125.99 128.28, 129.07, 129.12 and 129.39 (Ar-CH), 131.51 and 139.67 (Ar-C), 172.99 (2 x C=O) and 175.80 (C=O);  $m/z$  (EI) 350 (MH<sup>+</sup>, 82%), 349 (100), 307, 159, 133, 115, 91, 77 and 43.

**syn- and anti-1-Acetyl-4-methoxycarbonyl-2-methyl-3-(2-propyl)tetrahydropyrazoles (21 and 22).**

The procedure described for the synthesis of 1-acetyl-4-methoxycarbonyl-2-methyl-3-phenyltetrahydropyrazoles **10** and **11** was followed, but using 2-methylpropanal (0.24 g, 3.30 mmol), *N*<sup>1</sup>-acetyl-*N*<sup>2</sup>-methylhydrazine **9** (0.50 g, 5.67 mmol), methyl propenoate (1.00 g, 11.66 mmol) and toluene (50 mL), and heating at reflux for 16 h. Column chromatography (hexane:ethyl acetate 1:1 v/v) yielded *syn-diastereoisomer* **21** as a yellow oil (0.19 g, 25%), and a fraction that was purified by normal-phase preparative HPLC (CH<sub>3</sub>CN: water 30% v/v) to afford the *anti-diastereoisomer* **22** (0.09 g, 11%) and the *5-methoxycarbonyl isomer* **23** (0.03 g, 1%), both as yellow oils. *N*<sup>1</sup>-(2-Methoxycarbonylethyl)-*N*<sup>1</sup>-methyl-*N*<sup>2</sup>-acetylhydrazine **13** was also isolated (0.45 g, 46%).

**syn-Diastereoisomer (21).** Found: M<sup>+</sup>, 228.1474. C<sub>11</sub>H<sub>20</sub>N<sub>2</sub>O<sub>3</sub> requires M, 228.1474;  $\nu_{\max}$  (CHCl<sub>3</sub>)/cm<sup>-1</sup> 3670, 3441, 3013, 2925, 1740, 1647, 1439, 1373, 1245, 1185 and 1124;  $\delta_H$  (300 MHz; CDCl<sub>3</sub>) 0.83 and 0.94 (each 3H, d, *J* 6.6 Hz, CH<sub>3</sub>CH), 1.70 (1H, m, CHMe<sub>2</sub>), 2.20 (3H, s, CH<sub>3</sub>CON), 2.59 (3H, s, CH<sub>3</sub>N), 3.02 (1H, m, CHCHMe<sub>2</sub>), 3.67 (1H, m, 5-CHH), 3.72 (3H, s, CO<sub>2</sub>CH<sub>3</sub>), 3.76 (1H, m, CHCO<sub>2</sub>) and 4.20 (1H, m, 5-CHH);  $\delta_C$  (100 MHz; CDCl<sub>3</sub>) 18.02 (CH<sub>3</sub>CH), 20.75 (CH<sub>3</sub>CON), 21.09 (CH<sub>3</sub>CH), 29.23 CHMe<sub>2</sub>), 44.80 (5-CH<sub>2</sub>), 44.98 (CHCO<sub>2</sub>), 46.57 (CH<sub>3</sub>N), 52.00 (CO<sub>2</sub>CH<sub>3</sub>), 73.69 (CHCHMe<sub>2</sub>, 170.51 and 171.06 (C=O);  $m/z$  (EI) 228 (M<sup>+</sup>, 13%), 197, 185 (100), 143, 125, 83 and 43.

**anti-Diastereoisomer (22):** Found: M<sup>+</sup>, 228.1474. C<sub>11</sub>H<sub>20</sub>N<sub>2</sub>O<sub>3</sub> requires M, 228.1474;  $\nu_{\max}$  (CHCl<sub>3</sub>)/cm<sup>-1</sup> 2963, 1734, 1637, 1498, 1369, 1257 and 1185;  $\delta_H$  (400 MHz; CDCl<sub>3</sub>) 0.93 (3H, d, *J* 6.4 Hz, CH<sub>3</sub>CH), 0.99 (3H, d, *J* 6.8 Hz, CH<sub>3</sub>CH), 1.50 (1H, m, CHMe<sub>2</sub>), 2.16 (3H, s, CH<sub>3</sub>CON), 2.60 (3H, s, CH<sub>3</sub>N), 2.94 (1H, dd, *J* 3.9, 8.8 Hz, CHCHMe<sub>2</sub>, 3.05 (1H, m, CHCO<sub>2</sub>),

3.68 (1H, m, 5-*CHH*), 3.75 (3H, s, CO<sub>2</sub>CH<sub>3</sub>) and 4.37 (1H, m, 5-*CHH*);  $\delta_C$  (100 MHz; CDCl<sub>3</sub>) 19.11 and 19.81 (CH<sub>3</sub>CH), 20.57 (CH<sub>3</sub>CON), 32.61 (CHMe<sub>2</sub>), 44.22 (5-CH<sub>2</sub>), 45.79 (CH<sub>3</sub>N), 47.46 (CHCO<sub>2</sub>), 52.49 (CO<sub>2</sub>CH<sub>3</sub>), 77.07 (CHCHMe<sub>2</sub>), 170.88 and 173.99 (C=O); *m/z* (EI) 228 (M<sup>+</sup>, 24%), 197, 185 (100), 169, 143, 125, 83 and 43.

**5-Methoxycarbonyl isomer (23):** Found: C, 56.73; H, 8.74; N, 12.00%; M<sup>+</sup>, 228.1474. C<sub>11</sub>H<sub>20</sub>N<sub>2</sub>O<sub>3</sub>·0.25 H<sub>2</sub>O requires C, 56.75; H, 8.87; N, 12.03%; C<sub>11</sub>H<sub>20</sub>N<sub>2</sub>O<sub>3</sub> requires M, 228.1474;  $\nu_{\max}$  (CHCl<sub>3</sub>)/cm<sup>-1</sup> 2960, 1749, 1646, 1438, 1420 and 1370; major rotational isomer:  $\delta_H$  (400 MHz; CDCl<sub>3</sub>) 0.92 (3H, d, *J* 6.4 Hz, CH<sub>3</sub>CH), 1.05 (3H, d, *J* 6.3 Hz, CH<sub>3</sub>CH), 1.73 (1H, m, CHMe<sub>2</sub>), 2.06 (1H, m, 4-*CHH*), 2.18 (3H, s, CH<sub>3</sub>CON), 2.50 (3H, s, CH<sub>3</sub>N), 2.53 (1H, m, 4-*CHH*), 2.66 (1H, m, CHCHMe<sub>2</sub>), 3.74 (3H, s, CO<sub>2</sub>CH<sub>3</sub>) and 4.61 (1H, m, CHCO<sub>2</sub>);  $\delta_C$  (100 MHz; CDCl<sub>3</sub>) 20.28 (CH<sub>3</sub>CH), 20.87 (CH<sub>3</sub>CON), 20.96 (CH<sub>3</sub>CH), 28.73 (CHMe<sub>2</sub>), 32.61 (4-CH<sub>2</sub>), 46.25 (CH<sub>3</sub>N), 52.40 (CO<sub>2</sub>CH<sub>3</sub>), 57.63 (CHCO<sub>2</sub>), 72.33 or 73.52 (CHCHMe<sub>2</sub>), 170.55 and 172.36 (C=O); minor rotational isomer:  $\delta_H$  (400 MHz; CDCl<sub>3</sub>) 0.88 and 0.96 (each 3H, d, *J* 6.8 Hz, CH<sub>3</sub>CH), 1.33 (1H, m, CHMe<sub>2</sub>), 2.06 (1H, m, 4-*CHH*), 2.20 (3H, s, CH<sub>3</sub>CON), 2.50 (3H, s, CH<sub>3</sub>N), 2.53 (1H, m, 4-*CHH*), 2.66 (1H, m, CHCHMe<sub>2</sub>), 3.76 (3H, s, CO<sub>2</sub>CH<sub>3</sub>) and 4.61 (1H, m, CHCO<sub>2</sub>);  $\delta_C$  (100 MHz; CDCl<sub>3</sub>) 19.44 (CH<sub>3</sub>CH), 20.87 (CH<sub>3</sub>CON), 20.96 (CH<sub>3</sub>CH), 30.36 (CHMe<sub>2</sub>), 31.27 (4-CH<sub>2</sub>), 46.25 (CH<sub>3</sub>N), 52.40 (CO<sub>2</sub>CH<sub>3</sub>), 58.16 (CHCO<sub>2</sub>), 72.33 or 73.52 (CHCHMe<sub>2</sub>), 171.18 and 172.86 (C=O); *m/z* (EI) 228 (M<sup>+</sup>, 24%), 197, 185 (100), 169, 143, 125, 83 and 43 (100).

**2-*tert*-Butoxy-5-(1-methyl-2-acetyl)hydrazino-4-methyl-4,5-dihydrooxazole (25).** The procedure described for the synthesis of 1-acetyl-4-methoxycarbonyl-2-methyl-3-phenyltetrahydropyrazoles **10** and **11** was followed, but using (*S*)-*N-tert*-butoxycarbonylalaninal **24** (0.50 g, 2.89 mmol), *N*<sup>1</sup>-acetyl-*N*<sup>2</sup>-methylhydrazine **9** (0.38 g, 4.31 mmol), methyl propenoate (0.75 g, 8.66 mmol) and toluene (50 mL), and heating at reflux for 16 h. Column chromatography (hexane:ethyl acetate 1:1 v/v) followed by further purification by normal phase HPLC (30% CH<sub>3</sub>CN in water v/v) afforded only the title compound **25** (0.19 g, 27%), which was incompletely characterized;  $\delta_H$  (400 MHz; CDCl<sub>3</sub>) 1.36 (3H, d, *J* 7.3 Hz, CH<sub>3</sub>CH), 1.46 (9H, s, Me<sub>3</sub>C), 2.32 (3H, s, CH<sub>3</sub>CO), 3.19 (3H, s, CH<sub>3</sub>N), 4.45 (1H, br., OCHN), 5.39 (1H, br., CHNCH<sub>3</sub>) and 7.03 (1H, br. s, NH);  $\delta_C$  (100 MHz; CDCl<sub>3</sub>) 18.86 (CH<sub>3</sub>CH), 21.43 (CH<sub>3</sub>CO), 27.32 (CH<sub>3</sub>N), 28.57 (Me<sub>3</sub>C), 47.93 (CHCH<sub>3</sub>), 79.55 (CMe<sub>3</sub>), 141.97 (OCHN), 155.32 (C=N) and 172.60 (C=O); *m/z* (EI) 244 (MH<sup>+</sup>, 75%), 204, 201, 144, 128, 99 (100), 59, 57, 43 and 41.

**(*S*)-5-(1-Dibenzylaminoethyl)-2,4-dimethyl-4,5-dihydro-1,3,4-oxadiazole (27).** (*S*)-*N,N*-Dibenzylalaninal **26** (0.12 g, 0.47 mmol), *N*<sup>1</sup>-acetyl-*N*<sup>2</sup>-methylhydrazine **9** (0.05 g, 0.56 mmol) and methyl propenoate (0.33 g, 3.89 mmol) were dissolved in toluene (50 mL) and heated at reflux in the presence of 4Å molecular sieves for 16 h. Removal of the solvent under reduced pressure and column chromatography of the residue (hexane: ethyl acetate 1:1 v/v) afforded the *title compound 27*, a yellow oil (0.01g, 7%) as the only product isolated pure, although incompletely characterized;  $\delta_H$  (400 MHz; CDCl<sub>3</sub>) 1.29 (3H, d, *J* 7.4 Hz, CH<sub>3</sub>CH), 2.30 (3H, s, CH<sub>3</sub>C=N), 3.07 (3H, s, CH<sub>3</sub>N), 3.63 (2H, d, *J* 10.3 Hz, C<sub>6</sub>H<sub>5</sub>CHH), 3.67 (1H, m, CHCH<sub>3</sub>), 3.71 (2H, d, *J* 10.3 Hz, C<sub>6</sub>H<sub>5</sub>CHH), 6.94 (1H, d, *J* 5.0 Hz, NCHO) and 7.22-7.40 (10H, m, Ar-H).

**2-Benzyloxypropanal (28).** Ethyl lactate (7.40 g, 0.063 mol) in dry THF (20 mL) was added dropwise over 1 h to sodium hydride (5.00 g of a 60% suspension in oil, 0.125 mol) suspended in dry THF (120 mL). After stirring the mixture for 2 h, benzyl bromide (21.27 g, 0.126 mol) was added and the mixture stirred for a further 48 h, before dropwise addition of methanol: diethyl ether (1:1 v/v) to destroy excess sodium hydride. Saturated ammonium chloride solution was added and the resulting precipitate was filtered and washed with diethyl ether. The combined organic extracts were washed with saturated brine, dried ( $\text{MgSO}_4$ ) and purified by column chromatography (hexane: ether 9:1 v/v) to afford methyl 2-benzyloxypropanoate (4.31 g, 36%) that was used directly. To the foregoing ester (1.98 g, 10.19 mmol) stirred in anhydrous toluene (40 mL) at  $-50^\circ\text{C}$  was added dropwise DIBAL (10.16 mL of a 1.2M solution in toluene, 12.40 mmol) and the solution stirred for 2.5 h. A saturated solution of sodium metabisulfite (60 mL) was added and the mixture allowed to warm to  $20^\circ\text{C}$ . The layers were separated and the organic phase was further extracted with saturated sodium metabisulfite solution (3 x 80 mL). The combined aqueous layers were washed with diethyl ether (3 x 60 mL) then cooled to  $0^\circ\text{C}$ , the pH was raised to 11 with sodium hydroxide solution (8M) and the solution extracted with diethyl ether (3 x 50 mL), washed with saturated brine (75 mL), dried ( $\text{MgSO}_4$ ) and the solvent removed under reduced pressure. Purification by column chromatography (hexane: ethyl acetate 9:1 v/v) gave the *title compound 28* as a colorless oil (0.41 g, 25%); Found:  $\text{MNH}_4^+$ , 182.1183.  $\text{C}_{10}\text{H}_{12}\text{O}_2$  requires  $\text{MNH}_4$ , 182.1181;  $\nu_{\text{max}}$  (neat)/ $\text{cm}^{-1}$  3465, 3032, 2982, 2869, 2361, 1736, 1497, 1455, 1375 and 1094;  $\delta_{\text{H}}$  (300 MHz;  $\text{CDCl}_3$ ) 1.31 (3H, d,  $J$  7.0 Hz,  $\text{CH}_3$ ), 3.87 (1H, dq,  $J$  1.7 and 7.0 Hz,  $\text{CHCH}_3$ ), 4.57 and 4.64 (each 1H, d,  $J$  11.7 Hz,  $\text{CH}_2$ ), 7.28-7.36 (5H, m, Ar-H) and 9.65 (1H, d,  $J$  1.7 Hz, CHO);  $\delta_{\text{C}}$  (75 MHz;  $\text{CDCl}_3$ ) 15.26 ( $\text{CH}_3$ ), 71.96 ( $\text{CH}_2$ ), 79.41 ( $\text{CHCH}_3$ ), 127.91, 128.04 and 128.54 (Ar-CH), 137.37 (Ar-C) and 203.32 (CHO);  $m/z$  (CI) 182 ( $\text{MNH}_4^+$ , 100%), 125, 108, 106, 91 and 61.

**5-(1-Benzyloxyethyl)-2,4-dimethyl-4,5-dihydro-1,3,4-oxadiazole (29).** The procedure described for the synthesis of 1-acetyl-4-methoxycarbonyl-2-methyl-3-phenyltetrahydropyrazoles **10** and **11** was followed, but using 2-benzyloxypropanal **28** (0.37 g, 2.25 mmol),  $N^1$ -acetyl- $N^2$ -methylhydrazine **9** (0.20 g, 2.27 mmol), methyl propenoate (0.40 g, 4.66 mmol) and toluene (25 mL), and heating at reflux for 24 h. Column chromatography (hexane: ethyl acetate 1:1 v/v) afforded the *title compound 29* as a yellow oil (0.06 g, 11%) as the only product isolated pure;  $\delta_{\text{H}}$  (300 MHz;  $\text{CDCl}_3$ ) 1.40 (3H, d,  $J$  6.4 Hz,  $\text{CH}_3\text{CH}$ ), 2.32 (3H, s,  $\text{CH}_3\text{C}=\text{N}$ ), 3.13 (3H, s,  $\text{CH}_3\text{N}$ ), 4.18 (1H, quintet,  $J$  6.4 Hz,  $\text{CHCH}_3$ ), 4.55 (2H, s,  $\text{C}_6\text{H}_5\text{CH}_2$ ), 6.84 (1H, d,  $J$  6.4 Hz, NCHO) and 7.19-7.37 (5H, m, Ar-H).

**1-Acetyl-2-benzyl-4-methoxycarbonyl-3-(2-propyl)tetrahydropyrazole (31).**  $N^1$ -Acetyl- $N^2$ -benzylhydrazine **30** (0.32 g, 1.95 mmol), 2-methylpropanal (0.15 g, 2.09 mmol), methyl propenoate (0.52 g, 6.00 mmol), toluene (10 mL) and 4Å molecular sieves were placed in a glass tube, which was then sealed with a screw-cap. The mixture was heated for 72 h, after which the solvent was removed under reduced pressure. Column chromatography (hexane: ethyl acetate 2:1 v/v) of the residue gave the *title compound 31* (0.03 g, 4%) as a yellow oil; Found:  $\text{MH}^+$ , 305.1863.  $\text{C}_{17}\text{H}_{24}\text{N}_2\text{O}_3$  requires  $\text{MH}$ , 305.1865;  $\nu_{\text{max}}$  ( $\text{CHCl}_3$ )/ $\text{cm}^{-1}$  3021, 2963, 1733, 1641,

1437, 1369, 1216 and 700;  $\delta_{\text{H}}$  (300 MHz,  $\text{CDCl}_3$ ) 0.83 (3H, d,  $J$  6.9 Hz,  $\text{CH}_3\text{CH}$ ), 0.85 (3H, d,  $J$  7.1 Hz,  $\text{CH}_3\text{CH}$ ), 1.39 (1H, m,  $\text{CHMe}_2$ ), 1.92 (3H, s,  $\text{CH}_3\text{CON}$ ), 3.06-3.12 (2H, m,  $\text{CHCHMe}_2$  and  $\text{CHCO}_2$ ), 3.75 (1H, m, 5- $\text{CHH}$ ), 3.79 (3H, s,  $\text{CO}_2\text{CH}_3$ ), 3.83 (1H, d,  $J$  11.9 Hz,  $\text{C}_6\text{H}_5\text{CHH}$ ), 3.90 (1H, d,  $J$  11.9 Hz,  $\text{C}_6\text{H}_5\text{CHH}$ ), 4.52 (1H, dd,  $J$  9.8, 12 Hz, 5- $\text{CHH}$ ) and 7.26-7.38 (5H, m, Ar-H);  $\delta_{\text{C}}$  (75 MHz,  $\text{CDCl}_3$ ) 18.99 ( $\text{CH}_3\text{CH}$ ), 20.09 ( $\text{CH}_3\text{CH}$ ), 20.72 ( $\text{CH}_3\text{CON}$ ), 32.51 ( $\text{CHMe}_2$ ), 44.65 (5- $\text{CH}_2$ ), 47.56 ( $\text{CHCO}_2$ ), 52.54 ( $\text{CO}_2\text{CH}_3$ ), 61.84 ( $\text{C}_6\text{H}_5\text{CH}_2$ ), 74.86 ( $\text{CHCHMe}_2$ ), 127.93, 128.45 and 130.29 (Ar-CH), 136.29 (Ar-C), 171.59 and 174.37 (C=O);  $m/z$  (EI) 305 ( $\text{MH}^+$ , 2%), 261, 171, 111, 91 (100), 69, 65 and 43.

**1-tert-Butoxycarbonyl-4-methoxycarbonyl-2-methyl-3-phenyltetrahydropyrazole (33).** The procedure described for the synthesis of 1-acetyl-4-methoxycarbonyl-2-methyl-3-phenyltetrahydropyrazoles **10** and **11** was followed, but using benzaldehyde (0.26 g, 2.46 mmol),  $N^1$ -tert-butoxycarbonyl- $N^2$ -methylhydrazine **32** (0.35 g, 2.39 mmol), methyl propenoate (0.42 g, 4.89 mmol) and toluene (25 mL). Column chromatography (hexane: ethyl acetate 2:1 v/v) afforded the *title compound* **33** (0.14 g, 18%) as a colorless oil; Found:  $\text{M}^+$ , 320.1735.  $\text{C}_{17}\text{H}_{24}\text{N}_2\text{O}_4$  requires  $\text{M}$ , 320.1736;  $\nu_{\text{max}}$  ( $\text{CHCl}_3$ )/ $\text{cm}^{-1}$  2982, 1739, 1710, 1390, 1369, 1312, 1169 and 1121;  $\delta_{\text{H}}$  (300 MHz,  $\text{CDCl}_3$ ) 1.54 (3H, s,  $\text{Me}_3\text{C}$ ), 2.72 (3H, s,  $\text{CH}_3\text{N}$ ), 3.42 (3H, s,  $\text{CO}_2\text{CH}_3$ ), 3.72 (2H, m,  $\text{CHCO}_2$ , 5- $\text{CHH}$ ), 4.00 (1H, dd,  $J$  10.8, 14.3 Hz, 5- $\text{CHH}$ ), 4.32 (1H, d,  $J$  6.4 Hz,  $\text{C}_6\text{H}_5\text{CH}$ ) and 7.17-7.30 (5H, m, Ar-H);  $\delta_{\text{C}}$  (75 MHz,  $\text{CDCl}_3$ ) 28.52 ( $\text{Me}_3\text{C}$ ), 45.54 ( $\text{CH}_3\text{N}$ ), 46.36 (5- $\text{CH}_2$ ), 48.33 ( $\text{CHCO}_2$ ), 51.65 ( $\text{CO}_2\text{CH}_3$ ), 71.61 ( $\text{C}_6\text{H}_5\text{CH}$ ), 80.39 ( $\text{CMe}_3$ ), 126.70, 127.69 and 128.27 (Ar-CH), 137.85 (Ar-C), 154.79 (CON) and 170.19 ( $\text{CO}_2\text{CH}_3$ );  $m/z$  (EI) 320 ( $\text{M}^+$ , 3%), 265, 219, 187, 159, 133, 115, 91, 77, 57 (100) and 41.

## Acknowledgements

We wish to acknowledge the Open University for a studentship (S. J. H), the EPSRC National Mass Spectrometry Service Centre (University of Wales Swansea) for some mass spectral data, and the EPSRC X-Ray Crystallography Service (University of Southampton) for crystal structure determinations.

## References and Footnotes

1. (a) Jones, R. C. F.; Hollis S. J.; Iley, J. N. *Tetrahedron: Asymmetry*, **2000**, *11*, 3273. (b) Jones, R. C. F.; Gilbert, I. H.; Rees D. C.; Crockett, A. K. *Tetrahedron* **1995**, *51*, 6315.
2. *cf.*: Dutta, A. S.; Morley, J. S. *J. Chem. Soc., Perkin Trans. 1* **1975**, 1712. Mish, M. R.; Guerra, F. M.; Carreira, E. M. *J. Am. Chem. Soc.* **1997**, *119*, 8379.
3. See, for example: *Advances in Amino Acid Mimetics and Peptidomimetics*, Abell, A., Ed.; Jai Press Inc, Stamford (USA), 1999; Vol. 2.

4. Grashey, R. in *1,3-Dipolar Cycloaddition Chemistry*; Padwa, A., Ed.; John Wiley: New York, **1984**, p. 733.
5. See, for example: Sun, B.; Adachi, K.; Noguchi, M. *Tetrahedron* **1996** *52*, 901. Jacobi, P. A.; Martinelli, M. J.; Polanc, S. *J. Am. Chem. Soc.* **1984**, *106*, 5594. Oppolzer, W. *Tetrahedron Lett.* **1970**, 3091.
6. Oppolzer, W. *Tetrahedron Lett.* **1970**, 2199.
7. Huisgen, R.; Grashey, R.; Sauer, J. *The Chemistry of Alkenes*, Patai, S., Ed.; Wiley-Interscience, New York, 1964; p. 849 *et seq.*
8. Roussi, F.; Chauveau, A.; Bonin, M.; Micouin, L.; Husson, H.-P. *Synthesis* **2000**, 1170.
9. Chauveau, A.; Martens, T.; Bonin, M.; Micouin, L.; Husson, H.-P. *Synthesis* **2002**, 1885.
10. Condon, F. E. *J. Org. Chem.* **1972**, *37*, 3608.
11. Experiments carried out in an NMR tube with *d*<sub>8</sub>-toluene showed that heating for 72 h was required for complete reaction of the dipole to occur.
12. Dorn, H. *Tetrahedron Lett.* **1985**, *26*, 5123.
13. Kanemasa, S.; Tomoshige, N.; Wada, E.; Tsuge, O. *Bull. Chem. Soc. Jpn.* **1989**, *62*, 3944.
14. In all cycloadditions carried out with mono-substituted dipolarophiles, where two 4-substituted pyrazolidines were observed, the isomer identified as *syn*- was less polar in TLC analysis on silica than the *anti*- isomer. The chemical shift of the methine proton at C-4 was also found to be greater for the *syn*-isomer than for the *anti*-isomer ( $\delta$  3.94 ppm for **10** and  $\delta$  3.45 for **11**). Values for this proton in **21** and **22** are  $\delta$  3.76 and  $\delta$  3.05, respectively.
15. Chung, Y. J.; Ryu, E. J.; Keum, G.; Kim, B. H. *Bioorg. Med. Chem.* **1996**, *4*, 209.
16. Reetz, M. T.; Drewes, M. W.; Schmitz, A. *Angew. Chem., Int. Ed. Engl.* **1987**, *26*, 1141.
17. Calabretta, R.; Gallina, C.; Giordano, C. *Synthesis* **1991**, 536.
18. Lenman, M. M.; Lewis, A.; Gani, D. *J. Chem. Soc., Perkin Trans. 1* **1997**, 2297.
19. See: Perrin, D. D.; Armarego, W. L. F. *Purification of Laboratory Chemicals*, 3<sup>rd</sup> Edn.; Pergamon Press: Oxford, 1988.