

Structural correlations among aromatic azine derivatives

Theunis G.D. van Schalkwyk and Alistair M. Stephen

Department of Chemistry, University of Cape Town, Rondebosch 7701, South Africa

E-mail: stepham@science.uct.ac.za

(received 19 Jan 05; accepted 22 Mar 05; published on the web 20 Apr 05)

Abstract

This paper is a sequel to the *ARKIVOC* publication [2002 (iii) 103-111] describing the spontaneous conversion of the methochloride (**A**) of salicylaldazine to the azine (**B**), which is considered to take place *via* the oxidation of an *N*-alkylhydrazone. X-Ray crystallographic aspects of (**A**), (**B**) and three related compounds are presented, and a survey given of the conformations of a number of derivatives of the parent compound benzaldazine. Three categories of structure are discerned, one essentially planar, another twisted as a consequence of steric interactions in the neighborhood of the azine system, and a third group having further distinctive features including extreme deviations from planarity.

Keywords: Aromatic azines, conformation, configuration, X-ray crystallography

Introduction

The set of aromatic azines and their derivatives described is of interest on account of the facile de-*N*-alkylation of the methochlorides (and related compounds)^{1,cf.2}, and their interesting physical properties, but it is the conformational aspect that is presently under review. The literature has been scanned in order to make structural comparisons among substituted benzalazines, as has formerly been done *eg.* by Glaser and co-workers³ and others^{cf.4}; these compounds have been grouped according to the degree of distortion brought about by interactions within the azine system.

Results and Discussion

Prismatic yellow crystals of salicylaldazine methochloride (**A**) from methanol or other low-molecular weight alcohols, were usually threaded with needles of salicylaldazine (**B**), and furthermore the crystals were twinned; accordingly the structure determined by x-ray diffraction using direct methods of phasing⁴ did not have an R-factor of <0.145. However, the *Z,E*

configuration is unequivocal, by contrast with the expected *E,E* of (**B**). The near-planar structure of (**A**) and the juxtaposition of the ring planes of the molecules are described in the Experimental section, which gives also the crystal features of compounds (**C**)-(**E**). The structure of **B** was solved by direct methods⁴, final R-factor 0.041 (cf. Ref. 10). The methochloride of *p*-hydroxybenzaldehyde azine (**C**) and the azine (**D**) itself, likewise have the *Z,E* and *E,E* configurations respectively. Crystalline acetophenone azine (**E**), from which a metho-derivative could not be prepared, showed the *E,E* configuration, but was not centro-symmetric, and the principal residues (aromatic rings) tended to be orthogonal (cf. Ref. 15).

A CSD search, undertaken to find what structural variations existed among aromatic azines in general, yielded over 200 examples, from which 46 structures with formulae related to (**A**) – (**E**) were selected for comparison purposes. Eleven contained the system $[\text{ArCH}=\text{N-}]_2$, and a large set $[\text{ArCMe}=\text{N-}]_2$; some were related compounds with differently-substituted aromatic rings, and others were of the form $[\text{ArCH}=\text{N-N}=\text{CMeAr}']$. Seven examples contained other groups X, in the system $[\text{ArCX}=\text{N-}]_2$. The effects of steric hindrance on the preferred *E,E* structures are clearly seen; only one apart from those of (**A**) and (**C**) has the *Z,E* configuration. The most marked conformational changes result from substitution of various bulky groups for H on the azine carbon atoms; bizarre 3-dimensional structures develop where there is exceptional steric hindrance. The essential molecular formula showing torsional angles T1 to T5 and bond angles A1 and A2 is given below as Diagram 1, and should be used in conjunction with a model.




Diagram 1. Each molecule has two aromatic ring planes, designated A and B. Between lies a chain of four atoms that controls the conformation.

Of the torsional angles, if T3 is 180° as in the diagram, conjugation of two halves (A and B) of the azine molecule is at a maximum, i.e. *may* cross the N-N barrier³, with the N non-bonded electron pairs not being implicated. Values for T2 and T4 about the C=N double bond are relatively constant, except for some of the $[\text{ArCX}=\text{N-}]_2$ examples. Lack of conjugation of C=N with the aromatic pi-systems is suggested by the T1 and T2 values, which are often large. The relationship between the planes A and B follows from the magnitude and sign of T1 and of T2, and is influenced in *ca.* 15 examples by significant values of T3.

Analysis of molecular conformations (NB for *E,E* T3 is 180°, but in order to demonstrate deviations from planarity, the *E,E* angle T3 is recorded as 0°, and deviations as + or – , so to be

compatible with T1 and T5. This applies also to the T2 and T4 values where tabulated. For this reason the Greek tau symbol is not used).

Set (i). Molecules [ArCH=N-]₂

The prototype **1** (benzalazine) is, but for a 2.4° rotation of the ring planes, a planar molecule overall. **2** shows the effect of monoprotection of **1** at an N atom, which gives a T3 value of 5°, in addition to a 7° angle between the benzene ring planes. The twist of the benzene rings relative to one another enlarges to nearly 24° in the *p,p'*-dibromo derivative **3**, T3 here being restored to zero (an antiperiplanar arrangement of N=C groups). Overall planarity of structure characterizes **4**, in spite of the seven-carbon terminal chains, and similarly the mixed *p*-cyano/*p*-methoxyl derivative **5**, apart from the 7° angle between the benzene ring planes. **6** has been described in detail above (as **D**), exhibiting a considerable (14°) T3 and inter-benzene plane angle (17°). **7** is the important example described in full above (**B**). There is virtual overall planarity, and the structure uniquely allows for strong H-bonding between *ortho*-OH groups and N atoms. **8**, the Me ether corresponding to **6** (**D**), has a 4° twist associated with the benzene rings. **9** (4-pyridinium rings instead of phenyl) is unexceptionally planar; **10** and **11** show 11° and 7°, conrotatory, benzene-ring twists, and necessarily out-of-plane situations for F substituents in the CH₂F and CF₃ ethers respectively.

The average N-N bond length (D3) for examples **1-11**, all of which are *E,E* and show minimal effects on the azine system by the *p,p*-substituents, is 1.409 ± 0.003 Å, whereas the universal average for N-N is 1.383-1.403, and for N=N 1.255. The variation is within uncertainty limits, except for **8** and **9** (N-N 1.418 and 1.400 respectively), the *termini* of which are the most electron-releasing and electron-attracting. Details in Table 1. In this and subsequent tables centrosymmetric structures are starred. Concerning T3, the central rotational parameter, the policy was adopted in drawing up the tables that follow to express the torsional angle as a deviation from the 180° which allows maximum conjugation (because of coplanarity) of the C=N double bonds. This is based on recognition of the fact that the sign of T3 is assigned arbitrarily in the publications cited; signs of the other torsion angles are coherent with the decision made for T3 in particular cases. The signs for the angles recorded here (excepting T3) are as taken from the literature, and values are throughout rounded off to integral numbers.

Set (ii). Molecules [ArCMe=N-]₂**Table 1.** Asterisks following molecule numbers indicate *centrosymmetric*

Molecule ^{Ref}	N-N/Å	T3	T1	T5	Remarks	R1	R2	A1/°	A2/°
1* ⁵	1.412	0	2	-2		H	H	122	113
2 ⁶	1.402	5	4	-3	NH ⁺ (cf. A)	H	H	125	120
3 ⁷	1.408	0	21	-2		Br	Br	119	111
4* ⁸	1.406	0	-4	4		C ₆ H ₁₃ O	C ₆ H ₁₃ O	121	111
5 ⁹	1.406	-4	-2	4	<i>a</i>	MeO	CN	121	112
6 ⁴	1.412	-14	-4	13	= D	HO	HO	124	113
7* ¹⁰	1.410	0	-3	3	= B	H	H	121	113
8 ⁸	1.418	1	4	0	<i>b</i>	MeO	MeO	121	111
9* ¹¹	1.400	0	-	-	<i>bc</i>	NH ⁺	NH ⁺	119	112
10* ¹²	1.413	0	-11	11		F ₂ CHO	F ₂ CHO	121	115
11* ¹³	1.417	0	7	-7		F ₃ CO	F ₃ CO	121	122

a Contrast **5** with **29**, which has C_{azine}Me; T3 is 62° for **29**, and there are other distortions.

b Largest and smallest N-N bond lengths.

c CH in *p* position replaced by NH⁺.

Table 2. Asterisks following molecule numbers indicate *centrosymmetric*

Molecule ^{Ref}	N-N/Å	T3	T1	T5	Remarks	R1	R2	A1/°	A2/°
12* ¹⁴	1.390	0	-5	5	<i>a</i>	<i>o</i> -HO <i>p</i> -EtO	<i>o</i> -HO <i>p</i> -EtO	117	116
13* ¹⁵	1.397	0	-1	1		CN	CN	115	114
14* ¹⁶	1.404	0	12	-12		CO ₂ Et	CO ₂ Et	116	114
15* ¹⁶	1.395	0	3	-3		NHAc	NHAc	115	114
16* ¹⁶	1.402	0	-14	14		NMe ₂	NMe ₂	117	114
17 ¹⁷	1.402	0	11	-11	<i>b</i>	OCOEt	OCOEt	114	114
18 ¹⁶	1.409	0	-9	11		MeO	MeO	116	114
19* ¹⁸	1.405	0	-24	24	<i>c</i>	Me	Me	116	114

a Special case (cf.**7**) having *ortho*-HO, H-bonded to N, for each aryl ring. This promotes planarity of the azine system.

b One of two independent sets of molecules.

c Having the same formula as **26**, which is *not centrosymmetric*.

This group of compounds [ArCMe=N-]₂ is markedly different structurally from that described above as Set (i), in that steric hindrance causes distortion about the N-N bond, with the result that T3 tends in some of the cases to be much larger, and the general form of the molecule approximates rather to an orthogonal than a planar structure. There is, however, an observable gradation in this

trend. **12** is a special case, being almost planar, but this no doubt follows from the strong H-bonding of the *o*-HO substituent to N as observed in (**7**). **13** and **14** are relatively flat molecules, perhaps due to the effect of electron-withdrawing *p*-substituents which embody C-N triple bonds and trigonal C, respectively (and influence stacking). In **14** the opposite signs of T1 and T5 result in the aromatic rings being twisted conrotationally with respect to the azine system, losing conjugation, which however is gained by incorporation of the unsaturated substituent moieties. Compounds **15** to **19** all contain a planar system C=N-N=C, but of these **16** to **19** show marked twisting (T1 and T5: -24°, +24° for **19**).

The average bond length N-N for **12** to **19** (all centrosymmetric molecules) is 1.401 Å, with a large scatter (1.390 to 1.409), these being values for **12** (the special case; see above) and **18** (end groups electron-releasing, as for Set (i)). If anything this shows that Me replacing H in the azine moiety, releasing electrons on to N, increases the charge and shortens the N-N bond. Another way of regarding the effect of the methyl group is to assume hyperconjugation (the old "no-bond resonance") with H₂C=N imparting some double-bond character to N-N. Contrariwise, the electron density enhancement by conjugation of *p*-RO might lengthen N-N. More on the effects of electronegativities of *p*-groups is given below (see Table 3).

For **20** the twist about N-N is large, and disrotation of the benzene rings is large as well; in the case of **21** one ring only is skewed, in spite of the *p*- substituents being the same. The structural prototype of Set (ii) is acetophenone azine **22**, equal to (**E**), which has T3 41° and one benzene ring coplanar with C=N, but not the other ring with respect to the adjacent C=N, the latter being (T5) 20° out of plane. Thus for the two separate halves of the molecule conjugation is preserved in one but not in the other.

Table 3

Molecule Ref	CSD Code	N-N/Å	T3	T1	T5	Remarks	R1	R2	T2+ T4
20 ¹⁹	BITTIH	1.419	-31.6	18.6	12.8	<i>a</i>	HO	HO	-6.9
21 ¹⁶	ZEHIJUR	1.398	-28	1.2	14.1		NO ₂	NO ₂	-5.4
22 ¹⁵	LIKHIW	1.403	41.3	0.4	19.7	(=E)	H	H	2.9
23 ¹⁵	LIKHOH	1.396	-42	1.9	-10.6	<i>b</i>	F	F	-4.9
24 ¹⁵	LIKHUI	1.398	-45.3	-29.3	-30.5	<i>b</i>	Cl	Cl	-7.8
25 ¹⁵	LIKJEU	1.383	-55.4	-27.3	-21	<i>b</i>	Br	Br	-9.6
26 ¹⁸	PIYYAX	1.407	37.2	-0.5	19.9	(cf. E)	Me	Me	8.1
27 ¹⁶	ZEHIJIF	1.409	-48.9	-15.1	-0.2	<i>a</i>	NH ₂	NH ₂	-3.3
28 ²⁰	VIJVOH	1.406	40.7	13.9	6.3	<i>c</i>	O(CH ₂) ₅ _CO ₂ H	O(CH ₂) ₅ _CO ₂ H	1.6
29 ²¹	ZIFCAS	1.390	62.1	16.3	14.1	<i>d</i>	CN	OMe	7.2
30 ²²	BIWZIQ	1.414	56.4	29.6	-8.4	<i>a</i>	F	NH ₂	2.8
31 ²¹	ZIFBUL	1.374	-43.9	-17.9	-3.1	<i>e</i>	Br	OMe	-2.7
32 ²¹	ZIFCEW	1.396	-45.9	-1.4	-5.2	<i>f</i>	NO ₂	OMe	-9.1

a Very high N-N; cf. 6, Table 1.

b The torsion angles T3 follow a pattern reflecting the various electronegativities of R1= R2, as interpreted by Glaser and his co-workers (Ref. 15). Data for the I,I derivative given in Ref.23.

c For **28** complex (identical) end groups impart irregular profile.


d T3 exceptionally high. T1 and T5 nearly equal although the conjugative effects of electron-releasing OMe and electron-attracting CN differ.

e One of two independent sets of molecules, showing very different parameters. One only is quoted. T2 and T4 are small and nearly equal.

f Although **32** twisted about N-N, exterior groups R1 and R2 highly conjugated with aryl rings.

Set (iii). Molecules [ArCMe=N-]₂, all non-centrosymmetric

The group that follows, **20** to **32** forms a relatively coherent set and because of many resemblances to **22** (=E) are tabulated together, with comments as footnotes. The two halves of each azine are independent (T3 finite and large), and although the substitution patterns from **20** to **28** are the same in each molecule, the molecular conformations (T1 vs T5) are different. R1 and R2 differ in **29** to **32**. Inter-plane angles for pairs of aryl rings are very large for **22** to **32**. Table 3 includes first decimal, and summation of the torsion angles (including T2 and T4 which indicate twisting about double bonds) gives a value for the inter-plane angle in each case. These values agree well with those measured on the screen, when the molecule is appropriately displayed. Code letters are tabulated for **20** to **32**, and certain molecules are formulated in Diagram 1a.



28 R1 = R2 = O(CH₂)₅COOH

30 R1 = F, R2 = NH₂

31 R1 = Br, R2 = OMe

32 R1 = NO₂, R2 = OMe

Diagram 1a

Set (iv). Special set: various azine systems

Table 4. Asterisks following molecule numbers indicate *centrosymmetric*

Molecule ^{Ref}	N-N/Å	T3	T1	T5	Remarks	R1	R2	A1/ ^o	A2/ ^o
33 ⁹	1.405	-4	12	-2	<i>ab</i>	MeO	OCN	116	116
34 ²⁴	1.405	18	-11	-8	<i>a</i>	NO ₂	NMe ₂	121	112
35 ^{+ 9}	1.400	29	20	6	<i>a</i>	CN	OMe	121	114
36 ^{* 25}	1.396	0	-5	5	C _{azine} CN	H	H	121	113
37 ^{* 26}	1.407	0	6	-6	C _{az} (CO ₂ Me)	H	H	121	112
38 ^{* 27}	1.418	0	-6	6	C _{az} (CO ₂ Et)	H	H	121	112
39 ²⁸	1.410	<i>c</i>	<i>c</i>	<i>c</i>	C _{az} H,NH ₂	MeO	<i>d</i>	123	114
40 ²⁹	1.425	-6	<i>c</i>	<i>c</i>	C _{az} NH ₂ ,NH ₂	<i>e</i>	<i>e</i>	117	112

+ One of two independent sets of molecules.

a CazMe, CazH system in **33**; and CazH, CazMe in **34** and **35**.

b One ring plane skewed, minimal T3. This is one of four independent sets of molecules.

c Figures not available, but values are small.

d 2-Thiopyrazolyl in place of phenyl ring. A1 not=corresponding A4, A2 not=corresponding A3.

e *o*-Pyridyl in place of each phenyl ring.

The structures **33** to **40** form a sub-set resembling those of set (i), only **35** having the aryl rings twisted to a large extent, but here the variety of substituents on the azine carbon results in unusual features. The [PhC(CN)=N-]₂ system of **36** shows a conrotatory twist of 5° for each aryl ring, but N-N (1.396 Å) is close to that of the distantly related molecule **13**. **37** and **38** are carboxylate esters corresponding to the nitrile **36**, and are essentially planar, but the large ester groups lie orthogonally

to the plane; the phenyl rings are each twisted by 6° in a conrotatory mode. **39** and **40** are unique with primary amino groups on azine carbons, two such in **40**. For **40** T3 is 6° and the *o*-pyridyl rings are twisted with respect to each other. Certain of these molecules are formulated in Diagram 1b.




Diagram 1b

Set (v). Miscellaneous, centrosymmetric--> highly distorted

41 to **46** form a miscellaneous group of distorted structures with further distinctive features. Structure **41** falls into the Set (i) formula category, and T3 shows a planar inner azine moiety, but T1 and T5 are both large (-34° , 34°) and conrotatory, and the *o*-nitro substituents are out of plane. The appearance is unique; contrast **B**. **42** (the C_{azine}Me analogue of **41**) and **43** are configurational isomers, and form an *E,E* and *E,Z* pair; steric interaction of the nitro groups with N lone-pairs causes aryl-ring twist in **42**, and results in configurational change in **43**. The major consequence is a 180° rotational difference (T4). The other parameters of **42** and **43** differ little apart from T5 (-38° and 66° respectively). Structures **44** (benzophenone azine) to **46** resemble 4-bladed propellers on account of the exceptionally bulky nature of X in the systems [PhCX=N-]₂. For **44** A1 and A4 are large (124°), while for **46** they are different (112° and 123°). The T4 values recorded for **44** to **46** are 10° , 173° and -3° which are identical (or nearly so) to T2. **41**, **45** and **46** are formulated in Diagram 1c.

Table 5

Molecule ^{Ref}	N-N/Å	T3	T1	T5	Remarks	R1	R2	A1/°	A2/°
41 * ³⁰	1.424	0	-34	34	<i>a</i>	<i>o</i> -NO ₂	<i>o</i> -NO ₂	120	111
42 * ³⁰	1.420	0	38	-38	<i>b</i>	"	"	115	112
43 ³⁰	1.418	-14	37	66	<i>c</i>	"	"	116	114
44 ³¹	1.402	47	58	58	<i>d</i>	H	H	124	117
45 ³²	1.384	57	29	29	<i>e</i>	f	f	118	118
46 ²⁶	1.395	-53	-50	-84	<i>g</i>	C _{az} C Cl ₂ Me	C _{az} C Cl ₂ Me	112	118

a Azine carbons ArCH=.

b Azine carbons ArCMe=. The Me group in place of H has a minimal effect on T1 and T5.

c Configurational isomer of **42**, being *E,Z*; the *o*-NO₂ conflicting with the N lone pair is sufficient, in one half of the molecule, to cause exchange of the Me and Ar positions. T5 very large and A1not=A4 (126°).

d Large bulk of phenyl groups on azine carbons contributes to large torsion angles as seen, including T2=T4 at 10°, and inter-bond angles A1=A4=124°. Result: four non-parallel ring planes distributed in space.

e Resembles **44** in overall shape. T2=T4=7°.

f 2-Pyridyl groups instead of phenyl.

g A1 and A4 differ, 112° and 123°. R1=R2=H.




Diagram 1c

Summary

The factors influencing the sometimes very large deviations from planarity of azines were given at length, but generalisations regarding steric hindrance and conjugative effects are not easy, nor can the magnitude of the packing effect be quantified, in any particular case. It is difficult to assess the role of *p*-substituents whenever opposing effects may operate, typically when -OR groups are involved. Whereas there is an e-attracting influence due to the electronegativity of O relative to the ring carbon, the chemical, as opposed to static, effect is e-releasing into the ring by the resonance effect. Similarly with -NR₂, and in both instances e-withdrawing or e-releasing R complicates the issue.

T3, which indicates the extent of independence (cf. Ref.15) of the two halves, and the values of T1 and T5 which show torsion about the bond from azine carbon to aromatic ring, are the most critical determinants of structure of an azine molecule in the crystalline state. As to the interbond angles A1 and A2 the overall averages are 118° and 114° but the scatter is $\pm 6^\circ$, with occasional differences between these values and those for the other half of the azine molecule (A4 and A3). An extreme case of this effect appears in **33**, A1 and its counterpart A4 measuring 115° and 123° (unsymmetrical *p*-substitution, OMe and OCN, and azine CMe and CH, respectively); compare **34** and **35**. Torsion angles T2 and T4 are generally very similar, indicating co-planarity at the double bonds. A divergence in this respect is shown in the molecule **31**, an X=Me case, T4 being equal to 6° adjacent to the *p*-OMe substituted aromatic ring, and T2=1° where the substituent is Br. Curiously, when **18** and **25** are compared, it is the *dibromo* compound that has T4 distorted (see Tables 2 and 3). Of course when X=Ph, T2 and T4 values are both much larger, 10° for **44** and 7° for **45**; with X being CCl₂Me, T2 is -5° and T4 is -3°. Melting points, and a comparison of structures with those of di-aryl butadienes, might be topics that correlate with these findings. NMR correlations are complementary¹⁵, and, being solution-based, offset crystal packing effects. Six diagrams of the 46 printed in full in Ref. 4 are appended as Diagram 2 above to illustrate structural diversity among azines; these, given in order of increasing complexity, show in addition to the number given to the molecule (**7**, **22**, **29** etc.), its code letters (SALIAZ, etc.), formula, R factor, and numbered references (7, 5, 8, etc.). The views of the molecules in boxes lettered A, B and C in each lower left-hand corner show (A) conventional formulae, (B) stereo diagrams with atomic nomenclature, and (C) stereo diagrams orthogonal to those in B.




Diagram 2

Key to lettering and numbering are given in text

Experimental Section

The method of preparation and x-ray crystallographic examination of **A** and **B** were reported briefly, and stereo diagrams indicating their *Z,E* and *E,E* conformations were given¹. Details for **A**, **C**, **D** and **E** follow.

Structure A. $[C_{15}H_{15}N_2O_2]^+.Cl^-$ $P2_1/c$, $Z=4$, $a=7.967(1)$ $b=6.044(1)$ $c=29.407(2)$ Å, $\beta=95.60(4)^\circ$

X-ray diffraction showed that the crystal belonged to the monoclinic system (2/m Laue symmetry). The space group is $P2_1/c$, from the following conditions: hkl : none; $h0l$: $l = 2n$; $0k0$: $k = 2n$.

Density and unit cell volume indicated four formula units in the unit cell; direct methods were used to solve the structure. Careful inspection of the x-ray reflections of the crystal, mounted about the *b*-axis, showed that the crystal was twinned, and the structure obtained was an average one. Refinement did not yield an *R* factor of better than 0.145. (Evidence of twinning was obtained also for the fluoride, bromide and iodide analogues of the chloride **A**). Figure 1 displays (a) the perspective view, with atomic nomenclature, of **A**; (b) the stereo diagram; and (c) a view of the cation with r.m.s.(root mean square) ring plane A perpendicular. Figure 2 shows (a) a packing diagram for **A** viewed along $[010]$, and (b) its stereo diagram





Figure 1

**Figure 2**

The dihedral angle between the r.m.s. planes of rings A and B is $4.1(2)^\circ$; the r.m.s. deviation of the ring atoms is 0.01 \AA ; the angles between the r.m.s.

Planes of the azine system and those of A and B are $5.3(2)^\circ$ and $7.5(3)^\circ$, respectively; and the torsion angle T3 is 180° . Table 6 gives parameters for the two intermolecular hydrogen bonds that alternate between two oxygen and two chlorine atoms as the Figure shows. The two (nearly) coplanar phenyl rings are kept in place, parallel to the *c* axis, by two hydrogen bonds. Alternate planes of the rings are inclined at angles of approximately 60° .

Table 6

Bond			Distance (\AA)			Angle ($^\circ$)
D	H	A	D-H	H.....A	D.....A	D-H.....A
O(1)	H(1O)	Cl(1)	(a) 0.97	2.25	2.950(7)	124(1)
O(2)	H(2O)	Cl(1)	(b) 0.97	2.03	2.979(8)	164(1)

(D=donor, A=acceptor)

Close contacts:

Cl(1).....H(152)	(c)	2.62
H(2O).....H(152)	(d)	2.29

Equivalent positions:

x-1, *y*+1, *z* (b) *x*, *y*-1, *z* (c) -*x*+1, -*y*, -*z* (d) -*x*+1, -*y*, -*z*

Structure C. $[C_{15}H_{15}N_2O_2]^+.Cl^-$ $P2_1/n$, $Z=4$, $a=6.657(1)$ $b=16.134(1)$ $c=13.608(2)\text{\AA}$, $\beta=99.94(3)^\circ$

X-ray diffraction showed that the crystal belonged to the monoclinic system (2/m Laue symmetry). The space group is $P2_1/n$, from the conditions limiting reflections

$$hkl : \text{none}$$

$$h0l : h + l = 2n$$

$$0k0 : k = 2n$$

Unit cell and density measurements indicated $Z=4$, and hence the formula unit was placed in a general position. Direct methods were used to solve the crystal structure; R factor 0.093. Figure 3 gives (a) the perspective view of **C**, (b) its stereo diagram and (c) a view of the cation with r.m.s. plane A perpendicular. Figure 4 shows (a) the packing diagram of **C**, with hydrogen bonds, along $[100]$; and (b) its stereo diagram along $[100]$. Figure 5 displays stereo diagrams of **C** (a) along $[010]$ and (b) $[001]$.

The general structure of **C** resembles that of **A**, both having the atypical Z,E configuration. The dihedral angle between the least-squares planes through **A** and **B** is $8.4(3)^\circ$, and the r.m.s. deviations of the ring atoms is 0.01\AA . Dihedral angles between the mean plane of the azine system and planes A and B are $7.5(2)^\circ$ and $5.0(3)^\circ$, and the torsion angle T3 is $179.0(7)^\circ$. The chelated ring formed by the relatively weak hydrogen bond $C(5)-H(5)\cdots N(2)$ is nearly flat [maximum deviation 0.22\AA , for $N(2)$]. One relatively strong inter-ionic H bond [Table 7, entry (a)] gives stability to the structure.




Figure 3



**Figure 4****Figure 5**

Table 7

Bond			Distance (Å)			Angle (A°)
D	H	A	D-H	H.....A	D.....A	D-H.....A
C(5)	H(5)	N(2)	1.10	2.19	2.907(12)	121(3)
O(1)	H(1O)	Cl(1)	(a) 0.97	2.07	3.039(8)	176(5)

(D=donor, A=acceptor)

Close contacts:

Cl(1).....H(10)	(b)	2.07
Cl(1).....H(2)	(d)	2.78

Equivalent positions:

$-x+1, -y, -z-1$ (b) $-x+1, -y, -z-1$ (c) $-x+1, -y, -z-1$ note a no tb

Structure D. $C_{14}H_{12}N_2O_2$ $P2_1/n$, $Z=4$, $a=12.605(1)$, $b=8.670(1)$, $c=12.005(1)$ Å, $\beta=115.16(2)^\circ$

X-ray diffraction showed that the crystal belonged to the monoclinic system (2/m Laue symmetry). The space group is $P2_1/n$, from the conditions limiting reflections

$$hkl : \text{none}$$

$$h0l : h + l = 2n$$

$$0k0 : k = 2n$$

Unit cell and density measurements indicated $Z=4$, and hence the formula unit was placed in a general position. Direct methods were used to solve the crystal structure (R factor=0.043). These data resemble those for **C**. Figure 6 gives (a) the perspective view of **D**, (b) the stereo diagram and (c) a view of the cation with r.m.s. plane A perpendicular. Figure 7 shows (a) the packing diagram of **D**, with one intermolecular hydrogen bond, along [001]; and (b) its stereo diagram. Figure 8 displays stereo packing diagrams of **D** along (a) [100] and (b) [010].




Figure 6





Figure 7

**Figure 8**

The formula unit of **D** resembles that of **B** in having the E configuration at both C=N bonds in the azine chain. **D**, however, lacks centro-symmetry. The dihedral angle between the r.m.s. planes (A and B) of the two aryl rings is $9.5(1)^\circ$. The angles between the r.m.s. plane of the azine system and those of A and B are $-4.5(1)^\circ$ and $13.5(1)^\circ$ respectively. Due to noticeable kinks in the azine system (Figure 6 (c)) the torsion angle about N-N is $165.6(2)^\circ$. Among other non-symmetric structures having relatively small angular differences between A and B planes, yet showing very diverse molecular formulae, are **2**, **5**, **8**, **20**, **33** and **34**.

From the packing diagram (Figure 8 (a)) it is evident that the planes of molecules of **D** lie approximately *perpendicular* to the *b/c* plane, and in alternate directions approximately *parallel* to the two diagonals of the *b-c* plane. Distances between the molecular planes alternate between 3.3 and 3.6 Å. An intermolecular hydrogen bond O(2)-H(2O).....O(1) stabilises the structure by interconnecting the molecules along their linear axes. Table 8 records data for the H-bond.

Table 8

Bond			Distance (Å)			Angle (A°)
D	H	A	D-H	H....A	D.....A	D-H.....A
O(2)-H(2O)....O(1)			0.97	1.78	2.764(3)	173(2)

(D=donor, A=acceptor)

No close contacts observed

Equivalent positions:

(a) $x-1, y-1, z-1$

Structure E. $C_{16}H_6N_2$ $P2_1/n$, $Z=4$, $a=11.709(2)$, $b=7.552(1)$, $c=14.874(4)$ Å, $\beta=97.76(3)^\circ$ X-ray diffraction indicated that the crystal belonged to the monoclinic system (2/m Laue symmetry). The space group is $P2_1/n$, from the conditions limiting reflections

$$hkl : \text{none}$$

$$h0l : h + l = 2n$$

$$0k0 : k = 2n$$

Unit cell and density determinations fixed $Z=4$, and the formula unit of **E** is placed at a general position. Direct methods for structure determination gave an E-map showing some of the non-hydrogen atoms of two formula units overlapping, with one unit shifted by a distance of one half of a phenyl ring along the axis of the azine (Figure 9; projection down b axis). Refinement procedures did not help, but the structure was eventually solved by using the *average* positions of the two displaced molecules (R factor 0.046). Figure 10 shows (a) perspective and (b) stereo diagrams, and two views (c) and (d) of **E**, drawn with each of its r.m.s. ring planes A and B perpendicular. Figure 11 shows (a) the packing diagram of **E** viewed along $[010]$, and (b) its stereo diagram. Figure 12 shows stereo views of the packing diagram along (a) $[100]$ and (b) $[001]$.






Figure 9

**Figure 10****Figure 11**




Figure 12

The molecule (*E,E*) does not contain a centre of symmetry, but has the torsion angle $C(7)=N(1)-N(2)=C(8)$ equal to $-139.2(2)^\circ$. In this and other structural parameters there is good correlation with a published result⁵.

Acknowledgments

This paper is largely based on the Ph.D. thesis of the late Professor T.G.D.van Schalkwyk. Thanks are due to the J. Murray Hofmeyr Charitable Trust for providing financial assistance, and to Professors L.R.Nassimbeni and M.R.Caira for many helpful discussions. Dr Neil Ravenscroft and Mr Vincent Smith helped in the presentation of this paper.

References

1. Stephen, A. M.; vanSchalkwyk, T. G. D.; Ravenscroft, N. *ARKIVOC* **2002**, 103.
2. Teoh, S.-G.; Teo, S.-B.; Yeap, G.-Y.; Fun, H.-K. *Main Group Met. Chem.* **1994**, *17*, 595; *Chem. Abstr.* **1994**, *122*, 80804.
3. Lewis, M.; Glaser, R. *J. Org. Chem.* **2002**, *67*, 1441.
4. van Schalkwyk, T. G. D. *Ph.D. Thesis*, University of Cape Town, 2001.
5. Burke-Laing, M. and Laing, M. *Acta Cryst. B* **1976**, *32*, 3216.
6. Noltemeyer, M.; Schmidt, M.; Sheldrick, G. M. *Z. Kristallogr.* **1983**, *163*, 129.
7. Marignan, J.; Galigne, J. L.; Falgueirettes, J. *Acta Cryst. B* **1972**, *28*, 93.
8. Astheimer, H.; Walz, L.; Haase, W.; Loub, J.; Muller, H. J.; Gallardo, M. *Mol. Cryst. Liq. Cryst.* **1985**, *131*, 343.
9. Centore, R.; Panunzi, B.; Tuzi, A. *Z. Kristallografiya* **1996**, *211*, 31.

10. Arcovito, G.; Bonamico, M.; Domenicano, A.; Vaciago, A. *J. Chem. Soc. B* **1969**, 733.
11. Chen, W.; Liu, C.-M.; Li, D.-G.; You, X.-Z. *Acta Cryst. C* **1997**, 53, 1499.
12. Sereda, S. V.; Antipin, M. Yu.; Timofeeva, T. V.; Struchkov, Yu. T. *Kristallografiya* **1988**, 33, 118.
13. Sereda, S. V.; Shelyazhenko, S. V.; Fialkov, Yu. A.; Pivovarova, N. S.; Mikhaliuk, V. A. *Kristallografiya* **1992**, 37, 894.
14. Fayos, J.; Martinez-Ripoll, M.; Garcia-Mina, M. C.; Gonzalez-Martinez, J.; Arrese, F. *Acta Cryst. B* **1980**, 36, 1952.
15. Chen, G. S.; Anthamatten, M.; Barnes, C. L.; Glaser, R. *J. Org. Chem.* **1994**, 59, 4336.
16. Glaser, R.; Chen, G. S.; Anthamatten, M.; Barnes, C. L. *J. Chem. Soc., Perkin Trans. 2* **1995**, 1449.
17. Ciajolo, M. R.; Sirigu, A.; Tuzi, A. *Acta Cryst. C* **1985**, 41, 483.
18. Chen, G. S.; Anthamatten, M.; Barnes, C. L.; Glaser, R. *Angew. Chem., Int. Ed.* **1994**, 33, 1081.
19. Garcia-Mina, M. C.; Arrese, F.; Martinez-Ripoll, M.; Garcia-Blanco, S.; Serrano, J. L. *Acta Cryst. B* **1982**, 38, 2726.
20. Centore, R.; Ciajolo, M. R.; Roviello, A.; Sirigu, A.; Tuzi, A. *Mol. Cryst. Liq. Cryst.* **1990**, 165, 99.
21. Chen, G. S.; Wilbur, J. K.; Barnes, C. L.; Glaser, R. *J. Chem. Soc. Perkin Trans. 2* **1995**, 2311.
22. Lewis, M.; Barnes, C. L.; Glaser, R. *Can. J. Chem.* **1998**, 76, 1371.
23. Lewis, M.; Barnes, C. L.; Glaser, R. *J. Chem. Crystallog.* **1999**, 29, 1043.
24. Centore, R.; Garzillo, C. *J. Chem. Soc., Perkin Trans. 2* **1997**, 79.
25. de Meester, P.; Chu, S. S. C.; Johnson, J. E. *Acta Cryst. C* **1986**, 42, 1657.
26. Lai, E. C. K.; Mackay, D.; Taylor, N. J.; Watson, K. N. *Can. J. Chem.* **1988**, 66, 2839.
27. Glaser, R.; Chen, G. S.; Barnes, C. L. *J. Org. Chem.* **1993**, 58, 7446.
28. Dehne, H.; Drexler, K.; Scheunemann, A.; Reinke, H.; Michalik, M. *Phosphorus, Sulfur, Silicon, Related Elem.* **1996**, 118, 79.
29. Xu, Z.; Thompson, L. K.; Miller, D. O. *Inorg. Chem.* **1997**, 36, 3985.
30. Hsu, L.-Y.; Nordman, C. E.; Kenny, D. H. *Acta Cryst. C* **1993**, 49, 398.
31. Saha, A. K.; Hossain, M. M.; Grubisha, D. S.; Bennett, D. W. *J. Chem. Cryst.* **1995**, 25, 383.
32. Amadei, S.; Garcelli, M.; Ianelli, S.; Cozzini, P.; Pelagatti, P.; Pelizzi, C. *J. Chem. Soc. Dalton Trans.* **1998**, 1025.