

Interaction of plant amine oxidases with diaminoethers

Marek Šebela^{a,*}, Karla Jarkovská^a, René Lenobel^b, Rosaria Medda^c, Alessandra Padiglia^c, Giovanni Floris,^c and Pavel Pečá^a

^aDepartment of Biochemistry, Faculty of Science, Palacký University, Šlechtitelů 11, CZ-783 71 Olomouc, Czech Republic; ^bLaboratory of Growth Regulators, Palacký University and Institute of Experimental Botany - Czech Academy of Sciences, Šlechtitelů 11, 783 71 Olomouc, Czech Republic; ^cDepartment of Applied Sciences in Biosystems, University of Cagliari, I-09042 Monserrato (CA), Italy

E-mail: marek.sebela@upol.cz

Dedicated to Prof. Atta-ur-Rahman on the occasion of his 65th birthday

Abstract

Polyamines are ubiquitous compounds, which are involved in crucial physiological events including cell growth and differentiation. The catabolic oxidative degradation of polyamines is catalyzed by quinoprotein copper-containing amine oxidases (CAOs) and flavoprotein polyamine oxidases (PAOs). Various synthetic polyamine analogs and polyamine derivatives have been reported, which represent important tools (substrates or inhibitors) in the study of catalytic properties of the enzymes. In this work, two related compounds were studied in the reactions with plant amine oxidases: 1,8-diamino-3,6-dioxaoctane (DADO) and 1,10-bis(2-pyridinylmethyl)-4,7-dioxa-1,10-diazadecane (BPDD). Based on activity and stoichiometry assays together with spectrophotometric measurements, DADO can be considered a good substrate for grass pea, lentil and *E. characias* CAOs with K_m values in the range 10^{-4} – 10^{-3} M. Its oxidative degradation produces the corresponding aminoaldehyde 8-amino-3,6-dioxo-octanal, which does not undergo spontaneous cyclization (as it is known for the oxidation products of natural substrates putrescine, cadaverine and spermidine) or polymerization in the reaction mixture. Conversely, oat PAO does not oxidize DADO and is only weakly inhibited by the compound ($K_i = 1.6$ mM towards putrescine). BPDD was found to be a competitive inhibitor of both CAOs and PAOs with K_i values of 10^{-4} M. DADO could be suggested as a potential affinity ligand for CAOs.

Keywords: Diamine oxidase, diaminoether, inhibition, polyamine oxidase, substrate

Abbreviations: BPDD, 1,10-bis(2-pyridinylmethyl)-4,7-dioxa-1,10-diazadecane; CAO, copper amine oxidase; DADO, 1,8-diamino-3,6-dioxaoctane; ELAO; *Euphorbia characias* amine

oxidase; GPAO, grass pea amine oxidase; LSAO, lentil amine oxidase; OPAO, oat polyamine oxidase; PAO, polyamine oxidase

Introduction

Polyamines (i.e. putrescine, spermidine and spermine) are ubiquitous cationic compounds, which are involved in crucial physiological events including cell growth and differentiation. Polyamine levels are controlled through biosynthetic and biodegradation pathways¹. The oxidative degradation of polyamines is catalyzed by quinoprotein copper-containing amine oxidases (CAOs, EC 1.4.3.6) and flavoprotein polyamine oxidases (PAOs, EC 1.5.3.11)¹. As it is considered in the classification, CAOs attack primary amino groups of substrates, whereas PAOs attack secondary amino groups. In plants, CAOs preferentially oxidize diamines like putrescine or cadaverine and from that reason they are often called diamine oxidases². The oxidative deamination of CAO substrates results in the corresponding aldehydes, ammonium ions and hydrogen peroxide². PAOs catalyze the oxidative cleavage of spermine and spermidine at their secondary amino groups. Plant PAOs produce 4-aminobutanal (from spermidine) or *N*-(3-aminopropyl)-4-aminobutanal (from spermine), hydrogen peroxide and 1,3-propanediamine³. Representative members of the enzymes were crystallized and thoroughly characterized with respect to structure-functional relationships⁴.




Figure 1. Chemical structures of the studied compounds. (A) 1,8-diamino-3,6-dioxaoctane (DADO); (B) 1,10-bis(2-pyridinylmethyl)-4,7-dioxa-1,10-diazadecane (BPDD).

Various synthetic polyamine analogs and polyamine derivatives have been reported as substrates or inhibitors of amine oxidases⁵. Such compounds represent important tools in the study of catalytic properties of the enzymes and they often find applications in physiological research⁶. We synthesized two polyamine analogs comprising ether groups: 1,8-diamino-3,6-dioxaoctane (DADO) and 1,10-bis(2-pyridinylmethyl)-4,7-dioxa-1,10-diazadecane (BPDD); see Figure 1. DADO is often used as a bifunctional reactant in condensation reactions producing

macrocyclic compounds⁷. Similar applications in organic syntheses have been described for various analogs of BPDD, where the 2-pyridylmethyl moieties are replaced by benzyls, furfuryls, 2-methoxyethyls etc. Such substituents are converted into side arms in macrorings formed by condensation with 1,2-bis(2-iodoethoxy)ethane⁸. In this work, DADO and BPPD were tested for their interaction with plant amine oxidases. If enzymatic oxidation to the corresponding aldehydes would occur, it might initiate intramolecular or intermolecular cyclizations resulting in heterocyclic compounds. It should be noted in this context that the diaminoether 1,5-diamino-3-oxapentane is known as a good substrate of plant CAOs, which is converted to a dihydrooxazine⁹. Indeed, DADO was readily oxidized by plant CAOs. However, mass spectrometric analysis of the reaction mixture demonstrated that no cyclic product was formed. BPDD was not oxidized at all, but it provided a competitive inhibition. In case of plant PAO, both studied compounds were found to be competitive inhibitors showing only weak inhibition potency.

Results and Discussion

Amine oxidase assay used in this study is based on the formation of hydrogen peroxide by substrate oxidation, which is immediately utilized by peroxidase in the coupled detecting reaction. In this way, DADO was recognized as a substrate of plant CAOs. Relative rate of its oxidation (at a saturating concentration of 2.5 mM) by grass pea amine oxidase (GPAO) was 20% when that of putrescine oxidation was taken as 100%. This ratio was similar to those for 1,6-hexanediamine (29%), 1,7-heptanediamine (14%) and 1,8-octanediamine (10%). For the enzymes from lentil seedlings (LSAO) and *Euphorbia characias* latex (ELAO), the relative rates of DADO oxidation were 15 and 8%, respectively. Stoichiometry of the reaction was determined by oxygen consumption. In all three cases, 1 mol of oxygen was consumed per 1 mol of substrate, which is in agreement with a general scheme of CAO reaction². The following Michaelis-Menten constant (K_m) values for DADO of GPAO, LSAO and ELAO were determined: 0.15, 0.36 and 1.05 mM, respectively. Such values correspond well with those for the best natural diamine substrates of plant CAOs - putrescine, cadaverine and spermidine¹⁰. Using Britton-Robinson buffers for activity assay (pH range 5.0-9.0), an optimal pH value of 7.5 was found for DADO oxidation. Based on the above characteristics, DADO can be considered a good substrate of the enzymes, which have long been known for their broad substrate specificity.




Figure 2. Absorption spectroscopy of GPAO reaction with DADO. Figure shows difference absorption spectra of the interaction of native GPAO (20 μ M) with DADO (1.0 mM) in air saturated 20 mM potassium phosphate buffer, pH 7.0. The spectra were recorded immediately after mixing the reagents in 100-ms intervals for 10 s at 30°C .

Visible spectra of plant CAOs show a broad absorption band centered at 500 nm reflecting the presence of the topaquinone cofactor¹¹. Aerobic addition of a substrate to the enzyme is followed by rapid disappearance of the cofactor absorption, which slowly reappears after the substrate exhaustion. A stable yellow intermediate, the Cu(I)–semiquinolamine radical, is observable in anaerobiosis, when the reaction is initiated by substrate addition. By means of rapid scanning, the radical spectrum can be recorded also under admission of air¹². Figure 2 presents absorption spectra recorded in aerobiosis within first 10 s after rapid mixing of GPAO with DADO in potassium phosphate buffer, pH 7.0. The time delay of each successive spectrum was 100 ms. Absorption maxima at 365, 436 and 465 nm are clearly distinguished, which are characteristic for substrate-reduced CAOs. They indicate fast formation of the cofactor radical¹¹. The peak at 315 nm can be assigned to the Cu(II)–aminoresorcinol form of the cofactor, which is in equilibrium with the Cu(I)–semiquinolamine radical in the presence of air¹². Similar results were also obtained for LSAO and ELAO.




Figure 3. Tandem mass spectrometric (MS/MS) analysis of DADO oxidation product. DADO oxidation by an excess of GPAO was performed in 50 mM ammonium bicarbonate at 37 °C for 12h. After removing the enzyme by ultrafiltration, the reaction mixture was analyzed by electrospray-ionization mass spectrometry. The spectrum shown was recorded after collision-induced fragmentation of the parent ion belonging to the DADO oxidation product (m/z 148.1; $[M+H]^+$ - $C_6H_{14}NO_3^+$). The observed fragment ions: m/z 130.1 ($C_6H_{12}NO_2^+$), m/z 106.1 ($C_4H_{12}NO_2^+$), m/z 86.1 ($C_4H_8NO^+$).

The aminoaldehyde products of putrescine, cadaverine and spermidine oxidation by plant CAOs spontaneously cyclize to Δ^1 -pyrroline, Δ^1 -piperidine and 1,5-diaza-bicyclononane, respectively¹¹. Mass spectrometric analysis of GPAO/DADO reaction mixture revealed the presence of a reaction product with m/z 148, whose MS/MS fragmentation pattern (Figure 3) was consistent with the structure of 8-amino-3,6-dioxaoctanal. No peaks, which would indicate spontaneous cyclization or polymerization of the aldehyde were found in the mass spectrum. This is in contrast to the oxidation of 1,5-diamino-3-oxapentane as a related diaminoether compound, where the corresponding oxidation product undergoes intramolecular cyclization and loses water to form 3,6-2*H*-1,4-dihydrooxazine⁹. It seems that an intramolecular cyclization of 8-amino-3,6-dioxaoctanal to a dioxazonine derivative or condensations to larger polymers are unfavorable. Cyclization did not occur even after heating of the reaction mixture (after 12 h incubation at 37 °C) at 50 °C for 30 min.




Figure 4. Double reciprocal plot of competitive inhibition of GPAO by BPDD. Putrescine (Put) was a substrate. Initial rates were determined by the guaiacol spectrophotometric method²⁰. The assay was carried out in 0.1 M potassium phosphate buffer, pH 7.0, at 30 °C. BPDD concentrations were as follows: ▲ - 0, ● - 1.0 and ■ - 2.0 mM, $K_i = 0.28$ mM; y-axis represents the reciprocal absorbance values. The inset shows secondary plot of slopes against the inhibitor concentration.

BPDD was not oxidized at all by GPAO. In order to look for another effects on the enzyme, the compound was tested as an inhibitor towards the substrate putrescine. Figure 4 illustrates an experiment for the determination of inhibitory properties of the compound. As can be seen from the Lineweaver-Burk plot, BPDD functions as a weak competitive inhibitor of the enzyme and it is characterized by an inhibition constant (K_i) value of 0.28 mM. Inhibitors of plant CAOs have been recently reviewed¹³. The most potent competitive inhibitors show K_i values of 10^{-7} - 10^{-6} M (diaminoketones).

Plant PAOs are more specific than CAOs. Usually they oxidize only spermine and spermidine. For some enzymes (e. g. maize PAO), a limited number of other substrates (polyamine derivatives) is known¹⁴. Oat seedling PAO (OPAO) did not utilize DADO and BPDD as substrates. OPAO interaction with DADO was found to produce a weak competitive inhibition towards the substrate spermidine. The inhibition was characterized by a relatively high K_i value of 1.6 mM. This is surprising when we realize that diamines with 6 to 10 carbon atoms

are known as powerful competitive inhibitors of maize PAO with K_i values of 10^{-7} - 10^{-6} M¹⁵. As can be seen in the crystal structure of maize PAO, the active site consists of a „U-shaped“ tunnel, which passes through the protein molecule. Polyamine substrates or efficient PAO inhibitors bind at the active site and their molecules are bent to accommodate the tunnel shape¹⁵. Probably DADO is not flexible enough to compete with the natural substrate spermidine used for the inhibition assay. BPDD also inhibited OPAO in a competitive manner. The inhibition was moderate ($K_i = 0.58$ mM) though the terminal pyridinyl substituents could mediate a hydrophobic or ionic interaction facilitating the binding. Interestingly, *N,N'*-bis(benzyl)polyamine analogs derived by binding benzylaminopropyl groups to nitrogens of C4-C10 diamines have already been described as substrates for a mammalian PAO¹⁶.

In conclusion, we describe interactions of two synthetic diaminoether compounds with representatives of plant CAOs and PAOs. Whereas BPPD was recognized as a weak competitive inhibitor of the enzymes, the second studied compound – DADO – provided a different result. Plant CAOs accept DADO as a good substrate, plant PAOs are weakly inhibited by the compound. From that reason, we suggest possible DADO application as a ligand for affinity separation, which might selectively discriminate CAOs from PAOs in plant extracts. This suggestion comes from the fact that similarly based 6-aminoethyl-Sepharose has been successfully used for the isolation of LSAO and ELAO¹⁷.

Experimental Section

Chemicals. 2-Pyridinecarboxaldehyde was from Acros (Geel, Belgium). 1,2-Bis(chloroethoxy)ethane, deuterium oxide (99.96%), deuterated dimethyl sulfoxide (99.96%), DMF, potassium phthalimide and sodium borohydride were from Sigma-Aldrich Chemie (Steinheim, Germany). A hydrogen chloride solution in isopropanol was provided by Dr. Zatloukal, Laboratory of Growth Regulators, Faculty of Science, Palacký University. All other chemicals including ethanol (96%) were of analytical purity grade.

Enzymes. Grass pea (*Lathyrus sativus*) seedling amine oxidase (GPAO) was obtained as already described¹⁰. Amine oxidases from lentil seedlings (LSAO) and *Euphorbia characias* latex (ELAO) were purified according to previously published protocols¹⁷. The purified enzymes were stored frozen at -80 °C. Oat seedling polyamine oxidase (OPA) was isolated as recently published¹⁸. Horseradish (*Armoracia rusticana*) peroxidase was purchased from Sigma-Aldrich Chemie.

Instruments. NMR measurements were performed at 25 °C on a Bruker AVANCE 300 NMR spectrometer (Bruker BioSpin, Rheinstetten, Germany), operating at a magnetic field strength of 7.05 T. Mass spectra were acquired on a Quattro microTM triple-stage quadrupole mass spectrometer (Micromass, Manchester, UK) equipped with an electrospray ion source. Positive ionization mode was used producing protonated quasimolecular ions ($[M+H]^+$). All samples were directly introduced to the mass spectrometer by a syringe (flow rate of 10 μ l min⁻¹).

Parameters of the electrospray were as follows: capillary voltage 3.0 kV, cone voltage 30.0 V, desolvation gas flow 550 L h⁻¹, desolvation temperature 350 °C, source temperature 100 °C. Full scan spectra were obtained by scanning mass range 50-400 amu (0.7 s per scan). Tandem mass spectra (daughter ion scan mode) were obtained by scanning mass range from 50 to *m/z* of appropriate parent ion. Collision cell pressure was kept between 3.0 and 3.5 mbar with argon gas and collision cell energy was 25 eV. Elemental analyses were carried out using a combustion elemental analyzer Finnigan Flash EA1112 (Thermo Electron Corporation, San Jose, CA, USA). Melting points were measured using an SMP 3 apparatus for glass sample capillaries (Stuart Scientific, Redhill, Surrey, UK).

Compound characterization. 1,8-Diamino-3,6-dioxaoctane (DADO) was prepared by the Gabriel synthesis. According to a common procedure¹⁹, 1,2-bis(chloroethoxy)ethane (9.4 g, 50 mmol) was added dropwise with stirring at 0-5 °C to a suspension of potassium phthalimide (20.8 g, 110 mmol) in 70 mL of DMF over 15 min. The suspension was then heated for 3 h at 100°C, cooled, and poured into 1 L of ice-cold water. Filtration of the white solid and washing with water gave the intermediate bisphthalimide. A solution of this material (melting point 178-179 °C) in 50 mL of acetic acid and 50 mL of concentrated HCl was heated under reflux for 65 h. The precipitated phthalic acid was removed by filtration, the filtrate was concentrated to a volume of 10 mL in vacuo, and 250 mL of warm ethanol was added. Upon cooling, DADO dihydrochloride crystallized (7.8 g, 71%). The compound was recrystallized by dissolving in hot ethanol, filtration with charcoal and precipitation in refrigerator. ¹H-NMR (300 MHz, D₂O): 3.15 (t, 4H, *J*=5.1, CH₂N), 3.66 (s, 4H, OCH₂CH₂O), 3.70 (t, 4H, *J*=5.1, OCH₂CH₂N) ppm. ¹³C-NMR (75 MHz, D₂O): 39.1 (CH₂N), 66.4 (OCH₂CH₂N), 69.6 (OCH₂CH₂O) ppm. Elemental composition for C₆H₁₆N₂O₂·2HCl: C 32.80%, H 7.97%, N 12.21% (experimental values), C 32.59%, H 8.20%, N 12.67% (calculated values). ESI-MS: *m/z* 149 ([M+H]⁺ - C₆H₁₇N₂O₂⁺); MS/MS: *m/z* 106 (C₄H₁₂NO₂⁺), 88 (C₄H₁₀NO⁺), 70 (C₄H₆O⁺). Melting point was 114-115 °C. Free base of DADO (a transparent oil) was prepared by hydrazinolysis¹⁹ of the above bisphthalimide reaction intermediate.

1,10-Bis(2-pyridinylmethyl)-4,7-dioxa-1,10-diazadecane (BPDD) was synthesized by reductive alkylation of DADO. 2-Pyridinecarboxaldehyde (5.6 g, 52.5 mmol) was added dropwise to a solution of DADO (3.7 g of free base, 25 mmol) in 25 mL of methanol and the mixture was heated under reflux for 6 h. After cooling to 25 °C, NaBH₄ (2.7 g, 70 mmol) was added in small portions and the mixture was stirred for 16 h. The reaction was quenched by careful addition of 4M HCl until a strongly acidic pH was reached and the mixture was stirred for an additional 30 min. Then the solution was made basic using ammonia (12.5% in water), extracted with CH₂Cl₂ (3 x 50 ml) and the combined organic layers were dried (Na₂SO₄). Crude product was obtained as a yellow oil after evaporation of the solvent under reduced pressure. The product oil was purified by column chromatography on silica gel as described for a similar dibenzyl derivative⁷. BPDD tetrahydrochloride was obtained by adding an excess of hydrogen chloride in isopropanol and recovered by filtration (6.9 g, 58%). Recrystallization of the final product was performed by dissolving in hot ethanol, filtration with charcoal, cooling on ice and careful adding of ice-cold

diethylether. BPDD solution in water showed an absorption maximum at 260 nm revealing the presence of pyridine moiety. $^1\text{H-NMR}$ (300 MHz, d_6 -DMSO): 3.09 (t, 4H, $J=5.0$ Hz, CH_2N), 3.61 (m, 8H, $\text{CH}_2\text{OCH}_2\text{CH}_2\text{OCH}_2$), 3.82 (s, 4H, 2-pyridinylmethyl CH_2), 7.41 (m, 2H, CH-Py), 7.60 (d, 2H, $J=7.9$ Hz, CH-Py), 7.89 (m, 2H, CH-Py), 8.59 (d, 2H, $J=4.8$ Hz, CH-Py) ppm. $^{13}\text{C-NMR}$ (75 MHz, d_6 -DMSO): 48.6 (CH_2N), 59.1 (2-pyridinylmethyl CH_2), 65.5 ($\text{OCH}_2\text{CH}_2\text{N}$), 69.5 ($\text{OCH}_2\text{CH}_2\text{O}$), 125.4 (CH-Py), 125.6 (CH-Py), 140.9 (CH-Py), 149.8 (CH-Py), 156.8 (C-Py) ppm. Elemental composition for $\text{C}_{18}\text{H}_{26}\text{N}_4\text{O}_{2.4} \text{HCl}$: C 45.22%, H 6.30%, N 12.01% (experimental values), C 45.39%, H 6.35%, N 11.76% (calculated values). ESI-MS: m/z 331 ($[\text{M}+\text{H}]^+ - \text{C}_{18}\text{H}_{27}\text{N}_4\text{O}_2^+$); MS/MS: m/z 197 ($\text{C}_{10}\text{H}_{17}\text{N}_2\text{O}_2^+$), 179 ($\text{C}_{10}\text{H}_{15}\text{N}_2\text{O}^+$), 135 ($\text{C}_8\text{H}_{11}\text{N}_2^+$), 92 ($\text{C}_6\text{H}_6\text{N}^+$). Melting with decomposition was at 183 °C.

Enzyme assays. Amine oxidase activity was determined by spectrophotometer using a coupled reaction with horseradish peroxidase and guaiacol; putrescine served as a substrate²⁰. The method was also applied for polyamine oxidase; spermidine was a substrate in this case (1 mM final concentration). Both assays were performed at 30 °C in a thermostated cell. For inhibition measurements, the reaction mixture was pipetted as follows: 1.50 mL of 0.117 M potassium phosphate buffer, pH 7.0, 25 μL of 35 mM guaiacol, 25 μL of 5 μM horseradish peroxidase (2500 nkat mg^{-1}), 10 μL of diluted enzyme and 140 μL of inhibitor (in a proper dilution to get desired concentration). The reaction was initiated by the addition of substrate (50 μL). The mixture was continually stirred and monitored by increasing absorbance at 436 nm. Inhibition constants were calculated from Lineweaver-Burk plots constructed by linear regression of the measured data (three different measurements) using the computer program Microsoft Excel 2002. Protein content was determined using the spectrophotometric method with bicinchoninic acid; BSA served as a standard²¹. Rapid-scan absorption spectroscopy of the reaction mixture in the early phase of GPAO reaction with DADO was performed according to described methodology¹². For mass spectrometric analysis of oxidation products, 20 mM DADO in 50 mM ammonium bicarbonate was incubated with an excess of GPAO or LSAO at 37 °C for 12h. Then the mixture was ultrafiltrated using a 10 kDa cut-off Centricon filter units (Millipore, Bedford, MA, USA) and the filtrate injected for analysis. Reaction stoichiometry was analyzed as described²².

Acknowledgements

The authors gratefully acknowledge financial help of the grant MSM 6198959216 from the Ministry of Education, Youth and Sports, Czech Republic. Dr. Igor Popa from Laboratory of Growth Regulators, Palacký University, is thanked for performing NMR analyses.

References

1. (a) Bouchereau, A.; Aziz, A.; Larher, F.; Martin-Tanguy, J. *Plant Sci.* **1999**, *140*, 103. (b) Cona, A.; Rea, G.; Angelini, R.; Federico, R.; Tavladoraki, P. *Trends Plant Sci.* **2006**, *11*, 80.
2. Frébort, I.; Adachi, O. *J. Ferment. Bioeng.* **1995**, *80*, 625.
3. Šebela, M.; Radová, A.; Angelini, R.; Tavladoraki, P.; Frébort, I.; Peč, P. *Plant Sci.* **2001**, *160*, 197.
4. (a) Kumar, V.; Dooley, D. M.; Freeman, H. C.; Guss, J. M.; Harvey, I.; McGuirl, M. A.; Wilce, M. C. J.; Zubak, V. M. *Structure* **1996**, *4*, 943. (b) Binda, C.; Coda, A.; Angelini, R.; Federico, R.; Ascenzi, P.; Mattevi, A. *Structure* **1999**, *7*, 265.
5. (a) Peč, P.; Frébort, I. *Eur. J. Biochem.* **1992**, *209*, 661. (b) Shepard, E. M.; Smith, J.; Elmore, B. O.; Kuchar, J. A.; Sayre, L. M.; Dooley, D. M. *Eur. J. Biochem.* **2002**, *269*, 3645. (c) Bianchi, M.; Polticelli, F.; Ascenzi, P.; Botta, M.; Federico, R.; Mariottini, P.; Cona, A. *FEBS J.* **2006**, *273*, 1115.
6. (a) Ha, H. C.; Woster, P. M.; Yager, J. D.; Casero, R. A. Jr. *Proc. Natl. Acad. Sci. USA* **1997**, *94*, 11557. (b) Burns, M. R.; LaTurner, S.; Ziemer, J.; McVean, M.; Devens, B.; Carlson, C. L.; Graminski, G. F.; Vanderwerf, S. M.; Weeks, R. S.; Carreon J. *Bioorg. Med. Chem. Lett.* **2002**, *12*, 1263.
7. Teysstot, M.-L.; Fayolle, M.; Philouze, C.; Dupuy, C. *Eur. J. Org. Chem.* **2003**, *1*, 54.
8. Gatto, V. J.; Arnold, K. A.; Viscariello, A. M.; Miller, S. R.; Morgan, C. R.; Gokel, G. W. *J. Org. Chem.* **1986**, *51*, 5373.
9. (a) Cragg, J. E.; Herbert, R. B.; Kgaphola, M. *Tetrahedron Lett.* **1990**, *47*, 6907. (b) Šebela, M.; Lamplot, Z.; Petřivalský, M.; Kopečný, D.; Lemr, K.; Frébort, I.; Peč, P. *Biochim. Biophys. Acta* **2003**, *1647*, 355.
10. Šebela, M.; Luhová, L.; Frébort, I.; Faulhammer, H. G.; Hirota, S.; Zajoncová, L.; Stučka, V.; Peč, P. *Phytochem. Anal.* **1998**, *9*, 211.
11. Medda, R.; Padiglia, A.; Floris, G. *Phytochemistry* **1995**, *39*, 1.
12. Šebela, M.; Frébort, I.; Lemr, K.; Brauner, F.; Peč, P. *Arch. Biochem. Biophys.* **2000**, *384*, 88.
13. Padiglia, A.; Medda, R.; Pedersen, J. Z.; Lorrain, A.; Peč, P.; Frébort, I.; Floris, G. *J. Enzym. Inhib.* **1998**, *13*, 311.
14. Federico, R.; Ercolini, L.; Laurenzi, M.; Angelini, R. *Phytochemistry* **1996**, *43*, 339.
15. Cona, A.; Manetti, F.; Leone, R.; Corelli, F.; Tavladoraki, P.; Polticelli, F.; Botta, M. *Biochemistry* **2004**, *43*, 3426.
16. Bitonti, A. J.; Dumont, J. A.; Bush, T. L.; Stemerick, D. M.; Edwards, M. L.; McCann, P. P. *J. Biol. Chem.* **1990**, *265*, 382-388.
17. (a) Floris, G.; Giartosio, A.; Rinaldi, A. *Phytochemistry* **1983**, *22*, 1871. (b) Padiglia, A.; Medda, R.; Lorrain, A.; Murgia, B.; Pedersen, J. Z.; Finazzi Agró, A.; Floris, G. *Plant Physiol.* **1998**, *117*, 1363.

18. Stránská, J.; Šebela, M.; Tarkowski, P.; Řehulka, P.; Chmelík, J.; Popa, I.; Peč, P. *Biochimie* **2006**, in press.
19. He, Z.; Nadkarni, D. V.; Sayre, L. M.; Greenaway, F. T. *Biochim. Biophys. Acta* **1995**, *1253*, 117.
20. (a) Frébort, I.; Haviger, A.; Peč, P. *Biológia (Bratislava)* **1989**, *44*, 729. (b) Smith, T. A. *Methods Enzymol.* **1983**, *94*, 311.
21. Smith, P. K.; Krohn, R. I.; Hermanson, G. T.; Mallia, A. K.; Gartner, F. H.; Provenzano, M. D.; Fujimoto, E. K.; Goeke, N. M.; Olson, B. J.; Klenk, D. C. *Anal. Biochem.* **1985**, *150*, 76.
22. Lamplot, Z.; Šebela, M.; Fryčák, P.; Longu, S.; Padiglia, A.; Medda, R.; Floris, G.; Peč, P. *J. Enzym. Inhib. Med. Chem.* **2005**, *20*, 143.