

Supporting Information

First preparation and isolation of a stable ferrocenylketene

Nanyan Fu, Annette D. Allen, and Thomas T. Tidwell*

Department of Chemistry, University of Toronto, Toronto, Ontario M5S 3H6 Canada

Email: ttidwell@chem.utoronto.ca

File 1 (RTF file). Crystal data and structure refinement for **6** (k0665).

Identification code	k0665	
Empirical formula	C ₂₁ H ₃₀ Fe O Si	
Formula weight	382.39	
Temperature	150(1) K	
Wavelength	0.71073 Å	
Crystal system	Monoclinic	
Space group	P 2 ₁ /n	
Unit cell dimensions	a = 7.9843(2) Å	a = 90°.
	b = 7.7592(3) Å	b = 93.920(2)°.
	c = 31.5333(10) Å	g = 90°.
Volume	1948.97(11) Å ³	
Z	4	
Density (calculated)	1.303 Mg/m ³	
Absorption coefficient	0.840 mm ⁻¹	
F(000)	816	
Crystal size	0.24 x 0.20 x 0.08 mm ³	
Theta range for data collection	2.59 to 27.53°.	
Index ranges	-10<=h<=10, -9<=k<=10, -36<=l<=40	
Reflections collected	13213	
Independent reflections	4308 [R(int) = 0.0704]	
Completeness to theta = 27.53°	98.6 %	
Absorption correction	Semi-empirical from equivalents	
Max. and min. transmission	0.937 and 0.657	
Refinement method	Full-matrix least-squares on F ²	
Data / restraints / parameters	4308 / 0 / 223	
Goodness-of-fit on F ²	1.048	
Final R indices [I>2sigma(I)]	R1 = 0.0453, wR2 = 0.1032	
R indices (all data)	R1 = 0.0731, wR2 = 0.1153	
Largest diff. peak and hole	0.402 and -0.790 e.Å ⁻³	

Table 2. Atomic coordinates ($\times 10^4$) and equivalent isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for **6** (k0665). $U(\text{eq})$ is defined as one third of the trace of the orthogonalized U^{ij} tensor

	x	y	z	U(eq)
Fe(1)	5147(1)	2785(1)	2995(1)	21(1)
Si(1)	2219(1)	5688(1)	4141(1)	20(1)
O(1)	6342(3)	3548(3)	4299(1)	45(1)
C(1)	3832(3)	4360(3)	3376(1)	22(1)
C(2)	2719(3)	3288(3)	3119(1)	24(1)
C(3)	2971(3)	3610(3)	2690(1)	27(1)
C(4)	4237(3)	4883(3)	2669(1)	27(1)
C(5)	4770(3)	5346(3)	3093(1)	25(1)
C(6)	6749(4)	1224(4)	3357(1)	32(1)
C(7)	5594(4)	211(4)	3096(1)	34(1)
C(8)	5807(4)	664(4)	2665(1)	33(1)
C(9)	7076(3)	1944(4)	2659(1)	31(1)
C(10)	7655(3)	2284(4)	3086(1)	32(1)
C(11)	3883(3)	4563(3)	3845(1)	23(1)
C(12)	5197(3)	4013(4)	4075(1)	29(1)
C(13)	631(3)	6577(3)	3724(1)	25(1)
C(14)	1266(4)	8043(4)	3457(1)	36(1)
C(15)	-1040(3)	7091(4)	3905(1)	36(1)
C(16)	3335(3)	7309(3)	4507(1)	27(1)
C(17)	4448(4)	8603(4)	4291(1)	39(1)
C(18)	2153(4)	8245(4)	4791(1)	34(1)
C(19)	1157(3)	4038(3)	4470(1)	23(1)
C(20)	2324(4)	3295(4)	4830(1)	31(1)
C(21)	396(4)	2579(3)	4187(1)	30(1)

Table 3. Bond lengths [Å] and angles [°] for **6**

Fe(1)-C(3)	2.032(3)
Fe(1)-C(4)	2.033(3)
Fe(1)-C(8)	2.036(3)
Fe(1)-C(9)	2.036(3)
Fe(1)-C(5)	2.036(3)
Fe(1)-C(10)	2.040(3)
Fe(1)-C(2)	2.042(3)
Fe(1)-C(7)	2.050(3)
Fe(1)-C(6)	2.050(3)
Fe(1)-C(1)	2.051(3)
Si(1)-C(19)	1.885(3)
Si(1)-C(11)	1.888(3)
Si(1)-C(16)	1.888(3)
Si(1)-C(13)	1.893(3)
O(1)-C(12)	1.174(3)
C(1)-C(5)	1.425(4)
C(1)-C(2)	1.427(4)
C(1)-C(11)	1.488(4)
C(2)-C(3)	1.405(4)
C(3)-C(4)	1.418(4)
C(4)-C(5)	1.421(4)
C(6)-C(10)	1.419(4)
C(6)-C(7)	1.429(4)
C(7)-C(8)	1.425(4)
C(8)-C(9)	1.420(4)
C(9)-C(10)	1.417(4)
C(11)-C(12)	1.304(4)
C(13)-C(14)	1.523(4)
C(13)-C(15)	1.540(4)
C(16)-C(18)	1.528(4)
C(16)-C(17)	1.532(4)
C(19)-C(20)	1.532(4)
C(19)-C(21)	1.539(4)
C(3)-Fe(1)-C(4)	40.83(11)
C(3)-Fe(1)-C(8)	104.75(11)
C(4)-Fe(1)-C(8)	118.97(12)
C(3)-Fe(1)-C(9)	120.46(12)

C(4)-Fe(1)-C(9)	104.49(12)
C(8)-Fe(1)-C(9)	40.82(11)
C(3)-Fe(1)-C(5)	68.52(11)
C(4)-Fe(1)-C(5)	40.88(11)
C(8)-Fe(1)-C(5)	155.83(12)
C(9)-Fe(1)-C(5)	121.11(11)
C(3)-Fe(1)-C(10)	157.76(12)
C(4)-Fe(1)-C(10)	122.47(12)
C(8)-Fe(1)-C(10)	68.40(11)
C(9)-Fe(1)-C(10)	40.69(12)
C(5)-Fe(1)-C(10)	108.60(11)
C(3)-Fe(1)-C(2)	40.35(11)
C(4)-Fe(1)-C(2)	68.44(11)
C(8)-Fe(1)-C(2)	122.30(11)
C(9)-Fe(1)-C(2)	157.36(11)
C(5)-Fe(1)-C(2)	68.43(11)
C(10)-Fe(1)-C(2)	160.98(12)
C(3)-Fe(1)-C(7)	120.94(12)
C(4)-Fe(1)-C(7)	155.59(12)
C(8)-Fe(1)-C(7)	40.81(12)
C(9)-Fe(1)-C(7)	68.80(12)
C(5)-Fe(1)-C(7)	162.44(12)
C(10)-Fe(1)-C(7)	68.54(12)
C(2)-Fe(1)-C(7)	108.29(11)
C(3)-Fe(1)-C(6)	158.51(12)
C(4)-Fe(1)-C(6)	160.43(12)
C(8)-Fe(1)-C(6)	68.49(12)
C(9)-Fe(1)-C(6)	68.57(12)
C(5)-Fe(1)-C(6)	125.86(11)
C(10)-Fe(1)-C(6)	40.59(12)
C(2)-Fe(1)-C(6)	124.83(12)
C(7)-Fe(1)-C(6)	40.80(12)
C(3)-Fe(1)-C(1)	68.52(10)
C(4)-Fe(1)-C(1)	68.84(11)
C(8)-Fe(1)-C(1)	160.28(11)
C(9)-Fe(1)-C(1)	158.64(11)
C(5)-Fe(1)-C(1)	40.80(11)
C(10)-Fe(1)-C(1)	124.67(11)
C(2)-Fe(1)-C(1)	40.82(10)
C(7)-Fe(1)-C(1)	125.52(12)

C(6)-Fe(1)-C(1)	110.63(11)
C(19)-Si(1)-C(11)	108.40(12)
C(19)-Si(1)-C(16)	109.11(12)
C(11)-Si(1)-C(16)	106.93(12)
C(19)-Si(1)-C(13)	108.92(12)
C(11)-Si(1)-C(13)	106.62(12)
C(16)-Si(1)-C(13)	116.57(12)
C(5)-C(1)-C(2)	107.0(2)
C(5)-C(1)-C(11)	125.8(2)
C(2)-C(1)-C(11)	126.8(2)
C(5)-C(1)-Fe(1)	69.05(15)
C(2)-C(1)-Fe(1)	69.24(14)
C(11)-C(1)-Fe(1)	131.67(18)
C(3)-C(2)-C(1)	108.5(2)
C(3)-C(2)-Fe(1)	69.44(15)
C(1)-C(2)-Fe(1)	69.93(14)
C(2)-C(3)-C(4)	108.6(2)
C(2)-C(3)-Fe(1)	70.22(14)
C(4)-C(3)-Fe(1)	69.66(15)
C(3)-C(4)-C(5)	107.5(2)
C(3)-C(4)-Fe(1)	69.51(15)
C(5)-C(4)-Fe(1)	69.67(15)
C(4)-C(5)-C(1)	108.4(2)
C(4)-C(5)-Fe(1)	69.45(15)
C(1)-C(5)-Fe(1)	70.15(14)
C(10)-C(6)-C(7)	107.9(3)
C(10)-C(6)-Fe(1)	69.33(16)
C(7)-C(6)-Fe(1)	69.59(16)
C(8)-C(7)-C(6)	107.4(3)
C(8)-C(7)-Fe(1)	69.09(16)
C(6)-C(7)-Fe(1)	69.62(16)
C(9)-C(8)-C(7)	108.5(3)
C(9)-C(8)-Fe(1)	69.59(16)
C(7)-C(8)-Fe(1)	70.10(16)
C(10)-C(9)-C(8)	107.7(3)
C(10)-C(9)-Fe(1)	69.80(16)
C(8)-C(9)-Fe(1)	69.59(16)
C(9)-C(10)-C(6)	108.5(3)
C(9)-C(10)-Fe(1)	69.51(15)
C(6)-C(10)-Fe(1)	70.08(16)

C(12)-C(11)-C(1)	118.9(2)
C(12)-C(11)-Si(1)	116.4(2)
C(1)-C(11)-Si(1)	124.62(18)
O(1)-C(12)-C(11)	176.6(3)
C(14)-C(13)-C(15)	110.0(2)
C(14)-C(13)-Si(1)	115.09(19)
C(15)-C(13)-Si(1)	113.16(19)
C(18)-C(16)-C(17)	110.5(2)
C(18)-C(16)-Si(1)	112.99(19)
C(17)-C(16)-Si(1)	115.3(2)
C(20)-C(19)-C(21)	110.4(2)
C(20)-C(19)-Si(1)	112.94(18)
C(21)-C(19)-Si(1)	110.86(18)

Symmetry transformations used to generate equivalent atoms:

Table 4. Anisotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for k0665. The anisotropic displacement factor exponent takes the form: $-2p^2[h^2 a^{*2}U^{11} + \dots + 2 h k a^* b^* U^{12}]$

	U^{11}	U^{22}	U^{33}	U^{23}	U^{13}	U^{12}
Fe(1)	20(1)	18(1)	26(1)	-2(1)	3(1)	1(1)
Si(1)	19(1)	20(1)	22(1)	-1(1)	1(1)	1(1)
O(1)	31(1)	69(2)	35(1)	0(1)	-5(1)	16(1)
C(1)	22(1)	20(1)	23(1)	-3(1)	2(1)	6(1)
C(2)	17(1)	24(1)	32(2)	-3(1)	2(1)	3(1)
C(3)	25(1)	29(1)	26(2)	-4(1)	-2(1)	5(1)
C(4)	33(2)	26(1)	23(1)	2(1)	4(1)	7(1)
C(5)	27(1)	18(1)	32(2)	1(1)	5(1)	1(1)
C(6)	32(2)	30(2)	33(2)	4(1)	0(1)	11(1)
C(7)	32(2)	19(1)	53(2)	3(1)	11(1)	7(1)
C(8)	30(2)	25(1)	45(2)	-13(1)	2(1)	6(1)
C(9)	24(1)	29(2)	41(2)	-6(1)	11(1)	5(1)
C(10)	19(1)	32(2)	44(2)	-6(1)	0(1)	1(1)
C(11)	22(1)	24(1)	24(1)	-2(1)	3(1)	2(1)
C(12)	26(1)	35(2)	28(2)	-6(1)	5(1)	4(1)
C(13)	25(1)	25(1)	25(1)	1(1)	1(1)	3(1)
C(14)	36(2)	38(2)	35(2)	12(1)	2(1)	5(1)
C(15)	26(2)	45(2)	37(2)	7(1)	2(1)	7(1)
C(16)	25(1)	25(1)	30(2)	-3(1)	-2(1)	0(1)
C(17)	37(2)	33(2)	45(2)	-7(2)	3(1)	-11(1)
C(18)	41(2)	28(2)	33(2)	-7(1)	3(1)	3(1)
C(19)	21(1)	22(1)	25(1)	-1(1)	5(1)	2(1)
C(20)	34(2)	30(2)	29(2)	4(1)	1(1)	2(1)
C(21)	31(2)	27(2)	31(2)	0(1)	2(1)	-2(1)

Table 5. Hydrogen coordinates ($\times 10^4$) and isotropic displacement parameters ($\text{\AA}^2 \times 10^{-3}$) for **6**.

	x	y	z	U(eq)
H(2A)	1908	2437	3227	29
H(3A)	2365	3033	2441	32
H(4A)	4673	5369	2404	33
H(5A)	5659	6209	3179	30
H(6A)	6886	1200	3674	38
H(7A)	4781	-652	3197	41
H(8A)	5164	171	2410	40
H(9A)	7490	2503	2400	37
H(10A)	8542	3137	3180	38
H(13A)	359	5608	3522	30
H(14A)	449	8274	3217	54
H(14B)	2344	7718	3349	54
H(14C)	1414	9081	3632	54
H(15A)	-1840	7450	3672	53
H(15B)	-848	8047	4105	53
H(15C)	-1497	6103	4052	53
H(16A)	4111	6623	4704	32
H(17A)	5131	9242	4507	58
H(17B)	3740	9410	4119	58
H(17C)	5185	7985	4106	58
H(18A)	2810	8983	4992	51
H(18B)	1528	7397	4948	51
H(18C)	1364	8954	4615	51
H(19A)	210	4629	4603	27
H(20A)	1699	2476	4996	47
H(20B)	2753	4232	5015	47
H(20C)	3266	2702	4710	47
H(21A)	-180	1757	4363	44
H(21B)	1293	1985	4048	44
H(21C)	-409	3065	3971	44

File 2. CIF file

data_k0665

_audit_creation_method SHELXL-97
_chemical_name_systematic
;
?
;
_chemical_name_common ?
_chemical_melting_point ?
_chemical_formula_moiety 'C21 H30 Fe O Si'
_chemical_formula_sum
'C21 H30 Fe O Si'
_chemical_formula_weight 382.39

loop_

_atom_type_symbol
_atom_type_description
_atom_type_scatter_dispersion_real
_atom_type_scatter_dispersion_imag
_atom_type_scatter_source
'C' 'C' 0.0033 0.0016
'International Tables Vol C Tables 4.2.6.8 and 6.1.1.4'
'H' 'H' 0.0000 0.0000
'International Tables Vol C Tables 4.2.6.8 and 6.1.1.4'
'O' 'O' 0.0106 0.0060
'International Tables Vol C Tables 4.2.6.8 and 6.1.1.4'
'Si' 'Si' 0.0817 0.0704
'International Tables Vol C Tables 4.2.6.8 and 6.1.1.4'
'Fe' 'Fe' 0.3463 0.8444
'International Tables Vol C Tables 4.2.6.8 and 6.1.1.4'

_symmetry_cell_setting Monoclinic
_symmetry_space_group_name_H-M 'P 21/n'
_symmetry_space_group_name_Hall '-P 2yn'

loop_

_symmetry_equiv_pos_as_xyz
'x, y, z'
'-x+1/2, y+1/2, -z+1/2'
'-x, -y, -z'
'x-1/2, -y-1/2, z-1/2'

```

_cell_length_a      7.9843(2)
_cell_length_b      7.7592(3)
_cell_length_c      31.5333(10)
_cell_angle_alpha   90.00
_cell_angle_beta    93.920(2)
_cell_angle_gamma    90.00
_cell_volume        1948.97(11)
_cell_formula_units_Z  4
_cell_measurement_temperature 150(1)
_cell_measurement_reflns_used 13213
_cell_measurement_theta_min 2.6
_cell_measurement_theta_max 27.5

_exptl_crystal_description 'plate'
_exptl_crystal_colour 'orange'
_exptl_crystal_size_max 0.24
_exptl_crystal_size_mid 0.20
_exptl_crystal_size_min 0.08
_exptl_crystal_density_meas ?
_exptl_crystal_density_diffn 1.303
_exptl_crystal_density_method 'not measured'
_exptl_crystal_F_000 816
_exptl_absorpt_coefficient_mu 0.840
# Absorption correction

_exptl_absorpt_correction_type 'multi-scan'
_exptl_absorpt_correction_T_min 0.657
_exptl_absorpt_correction_T_max 0.937
_exptl_absorpt_process_details
;
multi-scan from symmetry-related measurements
Sortav (Blessing 1995)
;

_exptl_special_details
;
?
;

_diffn_ambient_temperature 150(1)

```

```

_diffrn_radiation_wavelength  0.71073
_diffrn_radiation_type        MoK\alpha
_diffrn_radiation_source      'fine-focus sealed tube'
_diffrn_radiation_monochromator  graphite
_diffrn_measurement_device_type 'Bruker-Nonius KappaCCD'
_diffrn_measurement_method    '\f scans and \w scans with \k offsets'
_diffrn_detector_area_resol_mean 9
_diffrn_standards_number      0
_diffrn_standards_interval_count 0
_diffrn_standards_interval_time 0
_diffrn_standards_decay_%     0
_diffrn_reflns_number         13213
_diffrn_reflns_av_R_equivalents 0.0704
_diffrn_reflns_av_sigmaI/netI  0.0679
_diffrn_reflns_limit_h_min     -10
_diffrn_reflns_limit_h_max     10
_diffrn_reflns_limit_k_min     -9
_diffrn_reflns_limit_k_max     10
_diffrn_reflns_limit_l_min     -36
_diffrn_reflns_limit_l_max     40
_diffrn_reflns_theta_min       2.59
_diffrn_reflns_theta_max       27.53
_reflns_number_total           4308
_reflns_number_gt              3178
_reflns_threshold_expression    'I>2\sigma(I)'

_computing_data_collection     'Collect (Nonius BV, 1997-2002)'
_computing_cell_refinement     'Denzo-SMN (Otwinowski & Minor, 1997)'
_computing_data_reduction      'Denzo-SMN'
_computing_structure_solution   'SIR-92 (Altomare et al., 1994)'
_computing_structure_refinement 'SHELXTL V6.1(Sheldrick, 2001)'
_computing_molecular_graphics  'SHELXTL V6.1'
_computing_publication_material 'SHELXTL V6.1'

```

```
_refine_special_details
```

```
;
```

Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R-factors(gt) etc. and is

not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R-factors based on ALL data will be even larger.

;

```
_refine_ls_structure_factor_coef Fsqd
_refine_ls_matrix_type      full
_refine_ls_weighting_scheme  calc
_refine_ls_weighting_details
'calc w=1/[s^2*(Fo^2)+(0.0480P)^2+1.0926P] where P=(Fo^2+2Fc^2)/3'
_atom_sites_solution_primary direct
_atom_sites_solution_secondary difmap
_atom_sites_solution_hydrogens geom
_refine_ls_hydrogen_treatment constr
_refine_ls_extinction_method none
_refine_ls_extinction_coef  ?
_refine_ls_number_reflns    4308
_refine_ls_number_parameters 223
_refine_ls_number_restraints 0
_refine_ls_R_factor_all     0.0731
_refine_ls_R_factor_gt      0.0453
_refine_ls_wR_factor_ref    0.1153
_refine_ls_wR_factor_gt     0.1032
_refine_ls_goodness_of_fit_ref 1.048
_refine_ls_restrained_S_all  1.048
_refine_ls_shift/su_max     0.001
_refine_ls_shift/su_mean    0.000
```

loop_

```
_atom_site_label
_atom_site_type_symbol
_atom_site_fract_x
_atom_site_fract_y
_atom_site_fract_z
_atom_site_U_iso_or_equiv
_atom_site_adp_type
_atom_site_occupancy
_atom_site_symmetry_multiplicity
_atom_site_calc_flag
_atom_site_refinement_flags
```

_atom_site_disorder_assembly

_atom_site_disorder_group

Fe1 Fe 0.51474(4) 0.27851(5) 0.299539(12) 0.02117(13) Uani 1 1 d . . .
Si1 Si 0.22191(9) 0.56883(9) 0.41409(2) 0.02040(17) Uani 1 1 d . . .
O1 O 0.6342(3) 0.3548(3) 0.42988(7) 0.0455(6) Uani 1 1 d . . .
C1 C 0.3832(3) 0.4360(3) 0.33755(8) 0.0217(6) Uani 1 1 d . . .
C2 C 0.2719(3) 0.3288(3) 0.31193(9) 0.0243(6) Uani 1 1 d . . .
H2A H 0.1908 0.2437 0.3227 0.029 Uiso 1 1 calc R . .
C3 C 0.2971(3) 0.3610(3) 0.26900(9) 0.0267(6) Uani 1 1 d . . .
H3A H 0.2365 0.3033 0.2441 0.032 Uiso 1 1 calc R . .
C4 C 0.4237(3) 0.4883(3) 0.26691(9) 0.0272(6) Uani 1 1 d . . .
H4A H 0.4673 0.5369 0.2404 0.033 Uiso 1 1 calc R . .
C5 C 0.4770(3) 0.5346(3) 0.30932(9) 0.0253(6) Uani 1 1 d . . .
H5A H 0.5659 0.6209 0.3179 0.030 Uiso 1 1 calc R . .
C6 C 0.6749(4) 0.1224(4) 0.33567(10) 0.0317(7) Uani 1 1 d . . .
H6A H 0.6886 0.1200 0.3674 0.038 Uiso 1 1 calc R . .
C7 C 0.5594(4) 0.0211(4) 0.30958(10) 0.0344(7) Uani 1 1 d . . .
H7A H 0.4781 -0.0652 0.3197 0.041 Uiso 1 1 calc R . .
C8 C 0.5807(4) 0.0664(4) 0.26650(10) 0.0334(7) Uani 1 1 d . . .
H8A H 0.5164 0.0171 0.2410 0.040 Uiso 1 1 calc R . .
C9 C 0.7076(3) 0.1944(4) 0.26590(10) 0.0310(7) Uani 1 1 d . . .
H9A H 0.7490 0.2503 0.2400 0.037 Uiso 1 1 calc R . .
C10 C 0.7655(3) 0.2284(4) 0.30858(10) 0.0320(7) Uani 1 1 d . . .
H10A H 0.8542 0.3137 0.3180 0.038 Uiso 1 1 calc R . .
C11 C 0.3883(3) 0.4563(3) 0.38454(8) 0.0232(6) Uani 1 1 d . . .
C12 C 0.5197(3) 0.4013(4) 0.40746(9) 0.0292(6) Uani 1 1 d . . .
C13 C 0.0631(3) 0.6577(3) 0.37239(9) 0.0252(6) Uani 1 1 d . . .
H13A H 0.0359 0.5608 0.3522 0.030 Uiso 1 1 calc R . .
C14 C 0.1266(4) 0.8043(4) 0.34567(10) 0.0362(7) Uani 1 1 d . . .
H14A H 0.0449 0.8274 0.3217 0.054 Uiso 1 1 calc R . .
H14B H 0.2344 0.7718 0.3349 0.054 Uiso 1 1 calc R . .
H14C H 0.1414 0.9081 0.3632 0.054 Uiso 1 1 calc R . .
C15 C -0.1040(3) 0.7091(4) 0.39048(10) 0.0356(7) Uani 1 1 d . . .
H15A H -0.1840 0.7450 0.3672 0.053 Uiso 1 1 calc R . .
H15B H -0.0848 0.8047 0.4105 0.053 Uiso 1 1 calc R . .
H15C H -0.1497 0.6103 0.4052 0.053 Uiso 1 1 calc R . .
C16 C 0.3335(3) 0.7309(3) 0.45069(9) 0.0267(6) Uani 1 1 d . . .
H16A H 0.4111 0.6623 0.4704 0.032 Uiso 1 1 calc R . .
C17 C 0.4448(4) 0.8603(4) 0.42905(10) 0.0385(7) Uani 1 1 d . . .
H17A H 0.5131 0.9242 0.4507 0.058 Uiso 1 1 calc R . .

H17B H 0.3740 0.9410 0.4119 0.058 Uiso 1 1 calc R . .
H17C H 0.5185 0.7985 0.4106 0.058 Uiso 1 1 calc R . .
C18 C 0.2153(4) 0.8245(4) 0.47909(10) 0.0340(7) Uani 1 1 d . . .
H18A H 0.2810 0.8983 0.4992 0.051 Uiso 1 1 calc R . .
H18B H 0.1528 0.7397 0.4948 0.051 Uiso 1 1 calc R . .
H18C H 0.1364 0.8954 0.4615 0.051 Uiso 1 1 calc R . .
C19 C 0.1157(3) 0.4038(3) 0.44697(8) 0.0225(6) Uani 1 1 d . . .
H19A H 0.0210 0.4629 0.4603 0.027 Uiso 1 1 calc R . .
C20 C 0.2324(4) 0.3295(4) 0.48301(9) 0.0312(7) Uani 1 1 d . . .
H20A H 0.1699 0.2476 0.4996 0.047 Uiso 1 1 calc R . .
H20B H 0.2753 0.4232 0.5015 0.047 Uiso 1 1 calc R . .
H20C H 0.3266 0.2702 0.4710 0.047 Uiso 1 1 calc R . .
C21 C 0.0396(4) 0.2579(3) 0.41873(9) 0.0297(6) Uani 1 1 d . . .
H21A H -0.0180 0.1757 0.4363 0.044 Uiso 1 1 calc R . .
H21B H 0.1293 0.1985 0.4048 0.044 Uiso 1 1 calc R . .
H21C H -0.0409 0.3065 0.3971 0.044 Uiso 1 1 calc R . .

loop_

_atom_site_aniso_label
_atom_site_aniso_U_11
_atom_site_aniso_U_22
_atom_site_aniso_U_33
_atom_site_aniso_U_23
_atom_site_aniso_U_13
_atom_site_aniso_U_12

Fe1 0.0197(2) 0.0180(2) 0.0260(2) -0.00238(15) 0.00288(15) 0.00063(15)
Si1 0.0192(4) 0.0200(4) 0.0219(4) -0.0009(3) 0.0005(3) 0.0008(3)
O1 0.0309(11) 0.0693(16) 0.0354(12) -0.0003(11) -0.0053(10) 0.0162(11)
C1 0.0217(13) 0.0204(13) 0.0231(14) -0.0026(10) 0.0023(11) 0.0059(11)
C2 0.0165(12) 0.0244(13) 0.0320(16) -0.0026(11) 0.0021(11) 0.0028(11)
C3 0.0246(14) 0.0293(14) 0.0258(15) -0.0043(12) -0.0021(11) 0.0048(12)
C4 0.0333(15) 0.0262(14) 0.0225(14) 0.0022(11) 0.0038(12) 0.0068(12)
C5 0.0267(14) 0.0177(13) 0.0319(16) 0.0013(11) 0.0046(12) 0.0005(11)
C6 0.0323(15) 0.0299(15) 0.0325(16) 0.0038(13) -0.0003(13) 0.0106(13)
C7 0.0318(15) 0.0192(14) 0.053(2) 0.0030(13) 0.0110(14) 0.0067(12)
C8 0.0303(15) 0.0250(14) 0.0447(19) -0.0128(13) 0.0018(13) 0.0064(12)
C9 0.0238(14) 0.0293(15) 0.0410(18) -0.0060(13) 0.0112(12) 0.0046(12)
C10 0.0192(13) 0.0321(15) 0.0442(19) -0.0058(13) -0.0004(12) 0.0012(12)
C11 0.0218(13) 0.0244(13) 0.0237(14) -0.0020(11) 0.0027(11) 0.0020(11)
C12 0.0258(14) 0.0348(15) 0.0276(15) -0.0059(12) 0.0053(12) 0.0039(13)

C13 0.0251(14) 0.0253(13) 0.0250(14) 0.0007(11) 0.0007(11) 0.0033(11)
C14 0.0363(17) 0.0375(17) 0.0347(17) 0.0116(14) 0.0023(13) 0.0053(14)
C15 0.0258(15) 0.0445(18) 0.0366(18) 0.0068(14) 0.0019(13) 0.0074(14)
C16 0.0247(14) 0.0252(14) 0.0295(15) -0.0028(12) -0.0022(11) -0.0003(11)
C17 0.0369(17) 0.0333(16) 0.0454(19) -0.0074(15) 0.0033(14) -0.0113(14)
C18 0.0411(17) 0.0284(15) 0.0325(17) -0.0072(13) 0.0025(13) 0.0031(13)
C19 0.0207(13) 0.0217(13) 0.0254(14) -0.0010(11) 0.0045(11) 0.0016(11)
C20 0.0339(16) 0.0304(15) 0.0292(16) 0.0042(12) 0.0011(12) 0.0023(13)
C21 0.0312(15) 0.0272(15) 0.0305(16) 0.0002(12) 0.0017(12) -0.0024(12)

_geom_special_details

;

All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

;

loop_

_geom_bond_atom_site_label_1

_geom_bond_atom_site_label_2

_geom_bond_distance

_geom_bond_site_symmetry_2

_geom_bond_publ_flag

Fe1 C3 2.032(3) . ?

Fe1 C4 2.033(3) . ?

Fe1 C8 2.036(3) . ?

Fe1 C9 2.036(3) . ?

Fe1 C5 2.036(3) . ?

Fe1 C10 2.040(3) . ?

Fe1 C2 2.042(3) . ?

Fe1 C7 2.050(3) . ?

Fe1 C6 2.050(3) . ?

Fe1 C1 2.051(3) . ?

Si1 C19 1.885(3) . ?

Si1 C11 1.888(3) . ?

Si1 C16 1.888(3) . ?

Si1 C13 1.893(3) . ?

O1 C12 1.174(3) . ?
C1 C5 1.425(4) . ?
C1 C2 1.427(4) . ?
C1 C11 1.488(4) . ?
C2 C3 1.405(4) . ?
C3 C4 1.418(4) . ?
C4 C5 1.421(4) . ?
C6 C10 1.419(4) . ?
C6 C7 1.429(4) . ?
C7 C8 1.425(4) . ?
C8 C9 1.420(4) . ?
C9 C10 1.417(4) . ?
C11 C12 1.304(4) . ?
C13 C14 1.523(4) . ?
C13 C15 1.540(4) . ?
C16 C18 1.528(4) . ?
C16 C17 1.532(4) . ?
C19 C20 1.532(4) . ?
C19 C21 1.539(4) . ?

loop_

_geom_angle_atom_site_label_1
_geom_angle_atom_site_label_2
_geom_angle_atom_site_label_3
_geom_angle
_geom_angle_site_symmetry_1
_geom_angle_site_symmetry_3
_geom_angle_publ_flag
C3 Fe1 C4 40.83(11) . . ?
C3 Fe1 C8 104.75(11) . . ?
C4 Fe1 C8 118.97(12) . . ?
C3 Fe1 C9 120.46(12) . . ?
C4 Fe1 C9 104.49(12) . . ?
C8 Fe1 C9 40.82(11) . . ?
C3 Fe1 C5 68.52(11) . . ?
C4 Fe1 C5 40.88(11) . . ?
C8 Fe1 C5 155.83(12) . . ?
C9 Fe1 C5 121.11(11) . . ?
C3 Fe1 C10 157.76(12) . . ?
C4 Fe1 C10 122.47(12) . . ?

C8 Fe1 C10 68.40(11) . . ?
C9 Fe1 C10 40.69(12) . . ?
C5 Fe1 C10 108.60(11) . . ?
C3 Fe1 C2 40.35(11) . . ?
C4 Fe1 C2 68.44(11) . . ?
C8 Fe1 C2 122.30(11) . . ?
C9 Fe1 C2 157.36(11) . . ?
C5 Fe1 C2 68.43(11) . . ?
C10 Fe1 C2 160.98(12) . . ?
C3 Fe1 C7 120.94(12) . . ?
C4 Fe1 C7 155.59(12) . . ?
C8 Fe1 C7 40.81(12) . . ?
C9 Fe1 C7 68.80(12) . . ?
C5 Fe1 C7 162.44(12) . . ?
C10 Fe1 C7 68.54(12) . . ?
C2 Fe1 C7 108.29(11) . . ?
C3 Fe1 C6 158.51(12) . . ?
C4 Fe1 C6 160.43(12) . . ?
C8 Fe1 C6 68.49(12) . . ?
C9 Fe1 C6 68.57(12) . . ?
C5 Fe1 C6 125.86(11) . . ?
C10 Fe1 C6 40.59(12) . . ?
C2 Fe1 C6 124.83(12) . . ?
C7 Fe1 C6 40.80(12) . . ?
C3 Fe1 C1 68.52(10) . . ?
C4 Fe1 C1 68.84(11) . . ?
C8 Fe1 C1 160.28(11) . . ?
C9 Fe1 C1 158.64(11) . . ?
C5 Fe1 C1 40.80(11) . . ?
C10 Fe1 C1 124.67(11) . . ?
C2 Fe1 C1 40.82(10) . . ?
C7 Fe1 C1 125.52(12) . . ?
C6 Fe1 C1 110.63(11) . . ?
C19 Si1 C11 108.40(12) . . ?
C19 Si1 C16 109.11(12) . . ?
C11 Si1 C16 106.93(12) . . ?
C19 Si1 C13 108.92(12) . . ?
C11 Si1 C13 106.62(12) . . ?
C16 Si1 C13 116.57(12) . . ?
C5 C1 C2 107.0(2) . . ?

C5 C1 C11 125.8(2) . . ?
C2 C1 C11 126.8(2) . . ?
C5 C1 Fe1 69.05(15) . . ?
C2 C1 Fe1 69.24(14) . . ?
C11 C1 Fe1 131.67(18) . . ?
C3 C2 C1 108.5(2) . . ?
C3 C2 Fe1 69.44(15) . . ?
C1 C2 Fe1 69.93(14) . . ?
C2 C3 C4 108.6(2) . . ?
C2 C3 Fe1 70.22(14) . . ?
C4 C3 Fe1 69.66(15) . . ?
C3 C4 C5 107.5(2) . . ?
C3 C4 Fe1 69.51(15) . . ?
C5 C4 Fe1 69.67(15) . . ?
C4 C5 C1 108.4(2) . . ?
C4 C5 Fe1 69.45(15) . . ?
C1 C5 Fe1 70.15(14) . . ?
C10 C6 C7 107.9(3) . . ?
C10 C6 Fe1 69.33(16) . . ?
C7 C6 Fe1 69.59(16) . . ?
C8 C7 C6 107.4(3) . . ?
C8 C7 Fe1 69.09(16) . . ?
C6 C7 Fe1 69.62(16) . . ?
C9 C8 C7 108.5(3) . . ?
C9 C8 Fe1 69.59(16) . . ?
C7 C8 Fe1 70.10(16) . . ?
C10 C9 C8 107.7(3) . . ?
C10 C9 Fe1 69.80(16) . . ?
C8 C9 Fe1 69.59(16) . . ?
C9 C10 C6 108.5(3) . . ?
C9 C10 Fe1 69.51(15) . . ?
C6 C10 Fe1 70.08(16) . . ?
C12 C11 C1 118.9(2) . . ?
C12 C11 Si1 116.4(2) . . ?
C1 C11 Si1 124.62(18) . . ?
O1 C12 C11 176.6(3) . . ?
C14 C13 C15 110.0(2) . . ?
C14 C13 Si1 115.09(19) . . ?
C15 C13 Si1 113.16(19) . . ?
C18 C16 C17 110.5(2) . . ?

C18 C16 Si1 112.99(19) . . ?
C17 C16 Si1 115.3(2) . . ?
C20 C19 C21 110.4(2) . . ?
C20 C19 Si1 112.94(18) . . ?
C21 C19 Si1 110.86(18) . . ?

_diffn_measured_fraction_theta_max 0.960
_diffn_reflns_theta_full 27.53
_diffn_measured_fraction_theta_full 0.986
_refine_diff_density_max 0.402
_refine_diff_density_min -0.790
_refine_diff_density_rms 0.096