

Thermal cyclisation reactions of methyl 2-(pyrrol-1-yl)cinnamate and methyl 3-(1-phenylpyrrol-2-yl) propenoate

Hamish McNab,^{*†} David Reed, Iain D. Tipping, and Richard G. Tyas

School of Chemistry, The University of Edinburgh, West Mains Road, Edinburgh EH9 3JJ, UK

E-mail: H.McNab@ed.ac.uk


Abstract

Flash vacuum pyrolysis (FVP) of the pyrrolylcinnamate derivative **3** gives the pyrrolo[1,2-*a*]quinoline **10**, the pyrrolo[2,1-*a*]isoindole **13** (R = Me) and the benzindoles **14** and **15**, all in low yield, whose structures were determined by NMR methods. The isomeric precursor **4** cyclises efficiently under the same conditions to give the pyrrolizin-3-one **16**.

Keywords: Flash vacuum pyrolysis, heterocyclisations, pyrroles, electrocyclisation

Introduction

In previous work we have established a versatile elimination-cyclisation sequence for the formation of fused pyrrol-2-one units (*e.g.* pyrrolizin-3-one **1**) under flash vacuum pyrolysis (FVP) conditions (Scheme 1).¹




Scheme 1

Although the general strategy was shown to be applicable to the synthesis of a variety of ring systems, attempted formation of the azepinone **2** by FVP of a vinylogous precursor was not


[†] Hamish McNab was Secretary and Treasurer of the RSC Heterocyclic Group during the period 1989-1992

successful.¹ This disappointing result may be due to the rotational degrees of freedom in the diene precursor at the high pyrolysis temperatures (Scheme 1). In order to probe this issue further, we reasoned that the incorporation of a fused benzene ring as part of the diene system would reduce these degrees of freedom and so might favour cyclisation to the larger ring size. For the synthesis of benzannulated pyrrolizin-3-ones, we have shown that the corresponding 1-arylpyrroles are useful starting materials, since they undergo a 1,5-aryl shift under FVP conditions prior to the elimination and cyclisation sequence (Scheme 2).²



Scheme 2

The cinnamate ester **3** was therefore selected as the initial target for the work reported in this paper. In the event, this precursor did not lead to seven-membered rings but instead has resulted in the discovery of a new type of electrocyclication-elimination process. The contrasting thermal behaviour of the isomeric precursor **4** is also reported.




Results and Discussion

The precursor **3** was made from 2-nitrocinnamic acid **5** by sequential formation of the ester **6** (64%), selective reduction to the amino compound **7** by zinc dust in the presence of ammonium chloride³ (51%) and formation of the pyrrole ring by condensation with 2,5-dimethoxytetrahydrofuran (54%). The crude precursor **3** was purified by dry-flash chromatography and characterised by standard methods. Its ¹H NMR spectrum confirms that the *E*-configuration of the alkene unit has been maintained throughout the reaction sequence (³*J* 16.0 Hz). A phenyl-substituted analogue of **3**, prepared by a different route, has been reported.⁴ The

isomeric precursor **4** was made by formylation of 1-phenylpyrrole **8** and separation of the 2-carboxaldehyde isomer **9** (91% of the mixture), followed by Wittig reaction.

FVP of the cinnamate **3** was carried out over a range of furnace temperatures, and it was found that 850 °C was required for complete conversion to products. Four significant products were isolated in low yield by dry-flash chromatography of the pyrolysate (Scheme 3). The first product (8%) showed a molecular ion at m/z 167 and was identified as pyrrolo[1,2-*a*]quinoline **10** by comparison with literature data.^{5,6} These data serve to exclude the isomeric pyrrolo[2,1-*a*]isoquinoline **11**,⁷ which might have been formed by loss of CO from the anticipated pyrroloazepinone product **12**. The connectivity of **10** is the same as in the starting material and so it follows that a novel direct cyclisation mechanism must take place which is of lower (or at least comparable) activation energy than the initial 1,5-aryl shift of the expected cascade process.



Scheme 3

The second product (m/z 169) was obtained in low yield (<11%) as the major component of an impure fraction. The most significant features of its ¹H NMR spectrum were a mutually coupled doublet (δ_{H} 1.68) and quartet (δ_{H} 5.08) in 3:1 integral ratio. The ¹³C NMR spectrum showed one methyl group and eight CH resonances, one of which was in the aliphatic region (δ_{C} 56.9), apparently attached to a nitrogen atom. A NOESY spectrum indicated that the methyl group was close in space to an aromatic proton as well as the aliphatic CH to which it is coupled. The pyrrolo[2,1-*a*]isoindole structure **13** (R = Me) is consistent with these data. Although this is not a known compound, the methylene protons of the parent pyrrolo[2,1-*a*]isoindole **13** (R = H) resonate at δ_{H} 4.92,^{8,9} very close to the value reported above for **13** (R = Me), and the remainder of the spectra are very similar [**13** (R = H), aromatic protons δ_{H} 7.1-7.3, pyrrole-type protons δ_{H}

6.3-7.0;^{8,9} **13** (R = Me), aromatic protons δ_{H} 7.2-7.5, pyrrole-type protons, δ_{H} 6.3-7.0]. Although present in low yield, this product shows skeletal connectivity derived from an initial 1,5-aryl shift and must therefore be formed in preference to products of the expected elimination-cyclisation sequence (e.g. **11**) shown in Scheme 3.

The final two products (14% in total) were closely related; although they co-eluted from the column, a pure sample of the component which was slightly more polar was obtained. Both products showed the presence of an NH signal in their ^1H NMR spectra, together with two other signals due to pyrrole-type protons and six signals in the aromatic region. The product which was not obtained in pure form, was unambiguously identified by comparison of its spectra with those reported for 1*H*-benz[*g*]indole **14** (see Experimental section).¹⁰ In particular the ^{13}C NMR spectra are closely comparable. The remaining isomer (*m/z* 167), which was obtained in a pure state, was identified as 3*H*-benz[*e*]indole **15** by the following NMR experiments. (The reported spectra of **15** are not sufficiently clear to allow unambiguous identification.¹¹) NOESY (and COSY) correlations between the aromatic doublets at δ_{H} 7.57 and 7.64 define the 1,2-fusion of the naphthalene ring and a NOESY correlation between the pyrrole-type proton at δ_{H} 7.14 and the aromatic doublet at δ_{H} 8.27 defines the mode of fusion of the heterocycle. Further NOESY correlations allow a complete assignment of the ^1H NMR spectrum and subsequent ^{13}C assignments were gained as a result of an HSQC experiment (Figure 1).





Figure 1. ^1H and ^{13}C NMR assignment of **15**.


In view of the unexpected nature of the results of this pyrolysis experiment, it was of interest to study the thermal behaviour of the isomeric pyrrole **4**. In this case FVP at 850 °C gave a single main product which was identified as the pyrrolizin-3-one **16** (38% after chromatography) by comparison of its spectra with those of an authentic sample (Scheme 4).¹ In contrast to the pyrrole derivative **3**, therefore, in the case of **4** a 1,5-sigmatropic shift of the phenyl group takes place (regiospecifically to the 5-position) followed by elimination of methanol to generate the ketene **17** and electrocyclisation to the pyrrolizinone **16** (Scheme 4).

These results clearly demonstrate that the anticipated rearrangement-elimination-cyclisation cascade dominates the FVP reactions of **4**, though no products are formed by this sequence in the FVP reactions of **3**. Possible reasons for this are discussed below.



Scheme 4

In pyrolysis of **3**, the product **10** is obtained without skeletal rearrangement but with loss of the entire ester function; such a cyclisation mechanism appears to be unprecedented. Our suggested route involves an initial 10-electron (disrotatory) electrocyclic cyclisation to provide the extended dipolar species **18** (Scheme 5). This may take place either with the *E*-cinnamate **3a** or with the *Z*-isomer **3b**; *E*-*Z*-isomerisation of cinnamate esters is known to be essentially equilibrated at 850 °C under FVP conditions.¹² Aromaticity of both the pyrrole and the benzene ring may be restored prior to the elimination step, by a 1,8-hydrogen shift from **18** to provide the 4,5-dihydro compound **19** (Scheme 5). There is good evidence that this route is followed in related cases.¹³




Scheme 5

The oxidative elimination step which leads from **19** to **10** appears to be a novel thermal behaviour of esters, though the high temperature elimination of sulfonyl groups to generate stilbene sub-units has some similarity.¹⁴ The study of model compounds, to be published elsewhere, suggest that at high temperatures the CO₂Me group, when it is situated in a benzylic-type position, is able to function as a radical leaving group leading (after subsequent loss of H[•])


to the oxidative elimination reaction observed in the pyrolysis of **3**. It is also of interest that the thermal elimination of sulfinate from sulfones to give stilbene-type products, takes place by a radical mechanism.¹⁴ The most likely mechanism for oxidative elimination step in the formation of **10** from **3**, therefore involves radical cleavage of the ester group to form a radical stabilised by the adjacent pyrrole ring, which can lose a hydrogen atom to form **10** (Scheme 5).

Though a similar electrocycloisatation mechanism to that in Scheme 5 can be drawn for the FVP of **4**, in this case the 1,5-aryl shift (leading to the pyrrolizinone **16**) is lower in energy. Electronic effects are unlikely to account for this result; electron withdrawing groups generally migrate *better* in 1,5-sigmatropic shifts (though the effect may be masked for aryl migrations¹⁵) so, paradoxically, the 1,5-aryl shift might be expected to be relatively favoured for **3** rather than **4**. Though the experimental results are clear, the reasons for the contrasting behaviour of **3** and **4** remain unknown.




Scheme 6

It was thought initially that the benzindoles **14** and **15** could be derived from the pyrroloquinoline **10** by a sequence of 1,5-shifts *via* **20** as the key intermediate. (Scheme 6). However, control pyrolyses of **10** at 850 °C and even at 950 °C showed no trace of these potential rearrangement products and so it is likely that they are formed by a different mechanism. Thus, initial 1,5-shift(s) of the aryl group can generate the 2- and 3-arylpyrroles **21** and **22** respectively. Compounds **14** and **15** can then be formed by electrocycloisatation followed by 1,5-hydrogen shift and elimination of the ester moiety in a similar fashion to the formation of **10** (Scheme 7 – *c.f.* Scheme 5). Once the 1,5-aryl shift has taken place, therefore, we believe that the novel ester elimination route is favoured over the expected ketene formation by methanol elimination. These results suggest that such an elimination step, requiring an eight or nine-membered transition state, is simply not viable, allowing the other processes to intervene.



Scheme 7

The formation of the pyrroloisindole **13** is unusual (formal loss of $C_2H_2O_2$ from the starting material) but a possible mechanism is shown in Scheme 8. We propose that the initial 1,5-aryl shift is followed by a 1,9-H shift leading to the intermediate **23** which can collapse to 2-(pyrrol-2-yl)styrene **24** by loss of $CH_2=O$ and CO. A further 1,9-H shift leads to the *o*-quinone dimethide **25**, the precursor of the final product. It is again noteworthy that a route such as this can apparently compete successfully with the expected ketene formation by methanol elimination from **21** (*c.f.* previous paragraph).



Scheme 8

In conclusion, it is clear from the results of the pyrolyses of **3** that the sequence of 1,5-aryl shift, *E/Z* isomerisation, thermal elimination of methanol and electrocyclisation to provide a fused azepinone system, cannot compete with alternative thermal processes. Product **10** is

formed from **3** even prior to the expected sigmatropic shift of the *N*-aryl group. Products **13-15** are obtained after sigmatropic shift, but are formed in preference to the azepinone **12** or its decomposition products (*e.g.* **11**). We conclude that the methanol elimination step (required for the formation of **12** from **21**) is not feasible geometrically, even though the formation of five-membered rings (*via* a six- or seven-membered transition state) is highly efficient. This efficiency is illustrated by the formation of **16** exclusively, by pyrolysis of **4**.

The substrate **3** is however left with no clearly defined route to products and a number of different pathways take place. One set of products, **10**, **14** and **15**, has provided a prototype of a new thermal cyclisation method, in which a newly recognised oxidative elimination of an ester group, taking place by a free-radical mechanism, provides a key aromatisation step. Further applications of this methodology in the synthesis of polycyclic ring systems are under investigation and will be reported in the near future.¹³

Experimental Section

General Procedures. ¹H and ¹³C NMR spectra are recorded at 250 (or 200) and 63 (or 50) MHz respectively for solutions in [²H]chloroform unless otherwise stated. Coupling constants are quoted in Hz; ¹³C NMR signals refer to CH resonances unless otherwise stated. ¹H/¹³C HSQC experiments were performed at 360 MHz using a Bruker DPX360 NMR spectrometer with gradient selection using the INVIGPTP sequence as provided by Bruker. NOESY and COSY experiments were similarly carried out at 360 MHz. Mass spectra were obtained under electron impact conditions unless otherwise stated.

Methyl 2-nitrocinnamate (6). A solution of 2-nitrocinnamic acid **5** (4.84 g, 0.025 mol) and anhydrous potassium carbonate (3.82 g) in dimethylformamide (30 cm³) was stirred magnetically. Iodomethane (3.92 g, 0.028 mol) was added dropwise over 10 min, and the solution left to stir for 24 h. The yellow solution was diluted with water (50 cm³) and extracted with diethyl ether (3 × 50 cm³). The ether was washed with water (6 × 50 cm³) and dried over MgSO₄. The ether was removed under reduced pressure to give the solid product **6** (3.31 g, 64%), mp 71-73 °C (lit.,¹⁶ 72-73 °C), δ_H 8.03 (1H, d, ³J 15.9), 7.98 (1H, d, ³J 10.0), 7.62-7.43 (3H, m), 6.30 (1H, d, ³J 15.9) and 3.76 (3H, s).

Methyl 2-aminocinnamate (7)³. Methyl 2-nitrocinnamate **6** (3.31 g, 0.016 mol) was dissolved in methanol (60 cm³). Water (7 cm³) and ammonium chloride (1.35 g, 0.025 mol) were added. The solution was stirred and zinc dust (9.00 g) was added over 15 min. After the initial reaction had subsided the solution was heated under reflux for 2 h and filtered hot through celite. The solution was diluted with water (100 cm³) and cooled in ice to separate the solid. The solid was filtered off and the solution extracted with dichloromethane (3 × 75 cm³). The dichloromethane layer was separated from the aqueous layer and concentrated to give **7** as a yellow solid³ (1.43 g, 51%) δ_H 7.77 (1H, d, ³J 15.7), 7.31 (1H, dd, ³J 8.1, ⁴J 2.8), 7.08 (1H, d, ³J 8.1), 6.73-6.61 (2H, m), 6.26 (1H, d, ³J 15.8), 3.91 [2H, br.s, (NH₂)] and 3.72 (3H, s).

Methyl 2-(pyrrol-1-yl)cinnamate (3). Methyl 2-aminocinnamate **7** (0.92 g, 5 mmol) was added to 2,5-dimethoxytetrahydrofuran (0.68 g, 5 mmol) with glacial acetic acid (1 cm³) and heated under reflux for 20 min. The acetic acid was removed under reduced pressure and the crude product distilled by Kugelrohr distillation. The product was then purified by dry flash chromatography on silica (15% ethyl acetate in hexane) to give *methyl 2-(pyrrol-1-yl)cinnamate 3* (0.64 g, 54%), mp 39-41 °C, bp 92 °C (0.2 Torr) [(FAB) Found MH⁺ 228.1024 C₁₄H₁₄NO₂ requires MH 228.1025], δ_H 7.69-7.25 (4H, m), 7.56 (2H, d, ³J 16.0), 6.81 (2H, s), 6.35 (1H, s), 6.32 (1H, d, ³J 16.0) and 3.76 (3H, s); δ_C 166.82 (quat), 140.53 (quat), 140.34, 130.49, 130.03 (quat), 127.54, 127.36, 126.55, 122.64 (2C), 119.90, 109.78 (2C) and 51.62 (CH₃); *m/z* (e.i.) 227 (M⁺, 0.6%), 169 (42), 168 (100), 167 (85), 166 (55), 140 (23), 139 (29), 128 (20), 115 (22), 83 (25), 75 (25), 63 (24) and 51 (29); *m/z* (FAB) 228 (MH⁺, 63%).

1-Phenylpyrrole-2-carbaldehyde (9). A solution of 1-phenylpyrrole **8** (9.99 g, 69.9 mmol) in DMF (100 cm³) was added to a solution of phosphoryl chloride (12.9 g, 84.0 mmol) in DMF (210 cm³) and stirred for 1 h. A further portion of phosphoryl chloride (12.9 g, 84.0 mmol) in DMF (210 cm³) was added and the solution was stirred overnight. The reaction mixture was poured over crushed ice and hydrolysed using 10 M sodium hydroxide (100 cm³) and neutralised with hydrochloric acid. The mixture was then extracted with ether (4 × 300 cm³) and washed with water (250 cm³), dried (MgSO₄) and concentrated under reduced pressure to give a brown oil. Distillation (Kugelrohr) of the crude product yielded the 1-phenylpyrrolecarbaldehyde as a mixture of the 2- and 3-isomers (11.07 g, 93%). A portion of the 1-phenylpyrrolecarbaldehyde mixture (2.68 g, 15.7 mmol) was separated into the two isomers using dry flash chromatography to yield 1-phenylpyrrole-2-carbaldehyde¹⁷ **9** (2.44 g, 91%) δ_H 9.65 (1H, s), 7.40–7.59 (5H, m), 7.24 (1H, dd, *J* 4.3 and 1.8), 7.15 (1H, t, *J* 1.8) and 6.48 (1H, dd, *J* 3.9 and 2.3).

Methyl 3-(1-phenylpyrrol-2-yl)propenoate (4). 1-Phenylpyrrole-2-carbaldehyde **9** (2.44 g, 14.3 mmol) was added to a solution of methyl (triphenylphosphoranylidene)acetate (4.80 g, 14.4 mmol) in toluene (50 cm³) and stirred under reflux for 6 h. The reaction product was concentrated under reduced pressure, absorbed onto silica and then separated using dry flash chromatography (10% ethyl acetate in *n*-hexane). The resulting solution was then distilled (Kugelrohr) to leave the product as orange crystals. The final product was recrystallised from hexane to give the *methyl 3-(1-phenylpyrrol-2-yl)propenoate 4* (0.91 g, 28%) mp 61 °C [Found, M⁺ 227.0951. C₁₄H₁₃NO₂ requires *M* 227.0946] δ_H 7.3-7.5 (6H, m) 6.97 (1H, m), 6.82 (1H, m), 6.35 (1H, m) 6.09 (1H, d, ³J 15.4) and 3.71 (3H, s); δ_C 167.98 (quat), 138.84 (quat), 133.37, 129.43 (2CH), 127.99, 126.72, 126.32 (2CH), 113.18, 112.34, 110.44 and 51.40 (CH₃) (one quaternary signal not apparent); *m/z* 227 (M⁺, 32%), 196 (27), 168 (84), 167 (100) and 51 (20).

Flash vacuum pyrolysis (FVP) experiments

The precursor was volatilised under rotary pump vacuum (pumping speed 100 dm³ min⁻¹) through an empty, electrically heated silica tube (35 × 2.5 cm) and the products were collected in a liquid nitrogen trap situated at the exit point of the furnace. The pressure was measured by a Pirani gauge situated between the product trap and the pump. The precursor, pyrolysis

conditions [quantity of precursor, furnace temperature (T_f), inlet temperature (T_i), pressure range (P) and pyrolysis time (t)] and products are given.

FVP of methyl 2-(pyrrol-1-yl)cinnamate (3). Methyl 2-(pyrrol-1-yl)cinnamate **3** (50 mg, 0.22 mmol) was distilled at 40-75 °C (0.05 Torr) into the furnace tube which was maintained at temperatures of 800, 850 or 900 °C. The optimum furnace temperature for complete conversion of starting material was 850 °C. A preparative pyrolysis of **3** (0.363 g, 1.6 mmol) was performed under the following conditions: T_f 850 °C, T_i 50-100 °C, P 0.07-0.01 Torr, t 100 min. Reaction products were separated by dry flash chromatography on silica using a hexane/ethyl acetate gradient as eluent (isolated products 0.146 g, 40%). The following components were identified, in order of elution: pyrrolo[1,2-*a*]quinoline **10** (21 mg, 8%) δ_H 7.88 (1H, d, 3J 8.8), 7.85 (1H, dd, 3J 2.9, 4J 1.4), 7.64 (1H, dd, 3J 7.8, 4J 1.5), 7.49 (1H, ddd, 3J 7.2 and 8.6, 4J 1.5), 7.34-7.26 (2H, m), 6.98 (1H, d, 3J 8.8), 6.78 (1H, dd, 3J 3.8 and 2.9) and 6.52 (1H, dd, 3J 3.8, 4J 1.4); δ_C 133.3 (quat), 130.9 (quat), 128.4, 127.5, 123.7 (quat), 123.3, 118.8, 118.5, 114.0, 112.5, 111.8 and 102.6; m/z 167 (M^+ , 100%), 166 (20), 139 (13), 123 (17), 105 (28), 69 (32), 57 (34), 43 (48) and 41 (45), spectra consistent with those in the literature.^{5,6} 5-methyl-5*H*-pyrrolo[2,1-*a*]isoindole **13** (R = Me) (30 mg, <11%), as the major component of an impure fraction (Found, M^+ 169.0893. $C_{12}H_{11}N$ requires M 169.0892) δ_H (360 MHz) 7.52 (1H, d, 3J 7.4), 7.34 (2H, m), 7.21 (1H, t, 3J ca. 7.4), 6.97 (1H, m), 6.41 (1H, dd, 3J 3.5 and 2.6), 6.33 (1H, dd, 3J 3.5, 4J 1.1), 5.08 (1H, q, 3J 6.9) and 1.68 (3H, d, 3J 6.9); δ_C (quaternary signals not quoted) 127.85, 124.73, 122.16, 118.38, 114.65, 112.51, 97.90, 56.90 and 20.89 (CH₃); m/z 169 (M^+ , 88%) and 154 (100): 1*H*-benz[*g*]indole **14** as the minor component in a mixture with **15**, (37 mg, 14% total) δ_H 8.85 (1H, br, NH), 7.96-7.91 (2H, m), 7.74 (1H, dd, 3J 8.8, 4J 0.7), 7.63-7.40 (3H, m), 7.24 (1H, m) and 6.70 (1H, dd, 3J 3.0, 4J 2.1); δ_C (CH signals only, literature data¹⁰ in parentheses) 128.7 (128.9), 125.3 (125.4), 123.7 (123.9), 122.6 (122.2), 120.7 (120.8), 120.6 (120.8), 119.2 (119.3) and 104.2 (104.3): a pure fraction of 3*H*-benz[*e*]indole **15** was also obtained, δ_H (360 MHz) 8.42 (1H, br, NH), 8.27 (1H, dm, 3J 8.1, 4J 1.4, 5J and nJ 0.7), 7.94 (1H, dm, 3J 8.1, 4J 1.4, 5J and nJ 0.7), 7.64 (1H, d, 3J 8.9), 7.59 (1H, ddd, 3J 8.1 and 6.9, 4J 1.4), 7.57 (1H, dd, 3J 8.9, 4J 0.9), 7.46 (1H, ddd, 3J 8.1 and 6.9, 4J 1.4) 7.32 (1H, ddd, 3J 2.9, 4J 2.5 and nJ 0.3) and 7.14 (1H, ddd, 3J 2.9, 4J 2.1 and nJ 0.9); δ_C (90 MHz) 132.60 (quat), 129.58 (quat), 128.93, 128.62 (quat), 126.19, 123.75, 123.41 (2C), 123.23 (quat), 122.59, 113.16 and 102.37; m/z 167 (M^+ , 100%), 155 (12), 140 (23), 139 (32), 105 (14), 86 (35), 84 (68) and 77 (14).

FVP of methyl 3-(1-phenylpyrrol-2-yl) propenoate (4). Methyl 3-(1-phenylpyrrol-2-yl) propenoate **4** (0.514 g, 2.26 mmol) was pyrolysed as above using the following parameters: T_f 850 °C, T_i 55-60 °C, P 0.01 Torr. The pyrolysate was absorbed onto silica and separated using dry flash chromatography (1% ethyl acetate in *n*-hexane) to give 5-phenylpyrrolizin-3-one **16** (0.169 g, 38%) bp 98 °C (0.4 Torr) [lit.,¹ 97-98 °C (0.2 Torr)]; δ_H 7.81-7.85 (2H, m), 7.25-7.49 (3H, m), 7.10 (1H, d, J 5.8), 6.18 (1H, dd, J 3.6 and 0.7), 6.07 (1H, d, J 3.3) and 5.70 (1H, dd, J 5.8 and 0.4); δ_C 166.21 (quat), 138.69 (quat), 138.49 (quat), 137.53, 129.63 (quat), 128.62, 128.14 (2CH), 126.75 (2CH), 121.76, 114.64 and 112.17; m/z 195 (M^+ , 100%), 167 (33), 139

(13) and 115 (11). Note that the chemical shift of the signal at δ_{H} 5.70 is wrongly reported as 5.09 in ref. 1; the spectra are otherwise compatible with those previously listed.¹

Acknowledgements

We are grateful to the Engineering and Physical Sciences Research Council for the award of the 360 MHz NMR spectrometer.

References

1. Campbell, S. E.; Comer, M. C.; Derbyshire, P. A.; Despinoy, X. L. M.; McNab, H.; Morrison, R.; Sommerville, C. C.; Thornley, C. *J. Chem. Soc., Perkin Trans. I* **1997**, 2195.
2. McNab, H.; Parsons, S.; Stevenson, E. *J. Chem. Soc., Perkin Trans. I* **1999**, 2047.
3. Skinner, W. A.; Schelstraete, M. G. M.; Baker, B. R. *J. Org. Chem.* **1961**, *26*, 1554.
4. Garofalo, A.; Ragno, G.; Campiani, G.; Brizzi, A.; Nacci, V. *Tetrahedron* **2000**, *56*, 9351.
5. Heffernan, M. L.; Irvine, G. M. *Aust. J. Chem.* **1976**, *29*, 847.
6. Jones, A. J.; Hanisch, P.; Heffernan, M. L.; Irvine, G. M. *Aust. J. Chem.* **1980**, *33*, 499.
7. Tominaga, Y.; Shiroshita, Y.; Kurokawa, T.; Gotou, H.; Matsuda Y.; Hosomi, A. *J. Heterocycl. Chem.* **1989**, *26*, 477.
8. Clark, B. A. J.; Despinoy, X. L. M.; McNab, H.; Sommerville, C. C.; Stevenson, E. *J. Chem. Soc., Perkin Trans. I* **1999**, 2049.
9. Stevenson, E. Ph.D. thesis, The University of Edinburgh, 1998.
10. Baudin, J.-B.; Julia, S. A.; Ruel, O. *Tetrahedron* **1987**, *43*, 881.
11. Sakamoto, T.; Kondo, Y.; Yamanaka, H. *Heterocycles* **1986**, *24*, 1845.
12. Hickson, C. L.; McNab, H. *J. Chem. Res. (S)* **1989**, 176.
13. McMillan, F. M.; McNab, H.; Milligan, A. A.; Tipping, I. D.; Tyas, R. G. unpublished work.
14. Claes, L.; François, J.-P.; Deleuze, M. S. *J. Am. Chem. Soc.* **2003**, *125*, 7129.
15. Jones, D. W.; Marmon, R. J. *J. Chem. Soc., Perkin Trans. I* **1993**, 681.
16. Kadaba, P. K. *Synthesis* **1971**, 316.
17. Defoin, A.; Geffroy, G.; Le Nouen, D.; Spileers, D.; Streith, J. *Helv. Chim. Acta* **1989**, *72*, 1199.