

Supplementary Material

Novel 1,1-diphenylhydrazine derivatives of benzofurazan and their dimerization

Elena N. Hristea,^a Marioara Bem,^a Teodor Silviu Balaban,^b Andreas Eichhöfer,^c
Miron T. Caproiu,^d Constantin Draghici,^d Gabriela Ionita,^a Tanta Spataru,^a
Cristian Enache,^e Maria Maganu,^d Adrian Beteringhe,^{a,f} Mihaela Hillebrand,^g
Titus Constantinescu,^a and Alexandru T. Balaban^{*h}

^a *Institute of Physical Chemistry “Ilie Murgulescu” of the Roumanian Academy, Splaiul
Independentei 202, 060021 Bucharest, Roumania*

^b *Université Paul Cézanne Aix Marseille III, Institut de Sciences Moléculaires de Marseille,
UMR 6263, Chirosciences, Marseille, France*

^c *Karlsruhe Institute of Technology, Institute of Nanotechnology, Postfach 3640, D-76021,
Karlsruhe, Germany*

^d *Institute of Organic Chemistry “C. D. Nenitzescu” of the Roumanian Academy, Splaiul
Independentei 202B, Bucharest 15-256, Roumania*

^e *Central Laboratory for Phytosanitary Quarantine, Bulevardul Voluntari 11, 077190 Voluntari,
Roumania*

^f *Lumina – University of South-East Europe, Colentina Road 64B, Bucharest – 021187,
Roumania*

^g *University of Bucharest, Department of Physical Chemistry, Bulevardul Elisabeta 4–12,
Bucharest, Roumania*

^h *Texas A&M University at Galveston, Department of Marine Sciences, Galveston, TX 77553-
1675, USA*

E-mail: balabana@tamug.edu

Content: X-Ray analysis for compounds 4 (see S1) and 13 (see S2)




Figure S1. Crystal packing diagrams of **4** showing strong π - π interactions between the furazan moiety and an N-phenyl group.




Figure S2. Packing diagrams of **13** in the crystal.