

Supplementary Material

Substituted quinolinones. Part 23. Synthesis of 6-ethyl-4,5-dioxo-5,6-dihydro-4H-pyrano[3,2-c]quinoline-3-carboxaldehyde and its chemical behavior towards hydroxylamine hydrochloride

Magdy A. Ibrahim,* Hany M. Hassanin, Mohamed Abass, and Shima'a Badran

*Department of Chemistry, Faculty of Education, Ain Shams University, Roxy, Heliopolis 11757,
Cairo, Egypt*

E-mail: magdy_ahmed1977@yahoo.com

Table of Contents

1. ^1H NMR spectrum of compound 4	S2
2. ^{13}C NMR spectrum of compound 4	S3
3. Mass spectrum of compound 4	S4
4. ^1H NMR spectrum of compound 6	S5
5. ^1H NMR spectrum of compound 10	S6
6. Mass spectrum of compound 10	S7
7. ^1H NMR spectrum of compound 13	S8
8. Mass spectrum of compound 13	S9




Figure S1. ^1H NMR spectrum of compound 4.




Figure S2. ^{13}C NMR spectrum of compound 4.




Figure S3. Mass spectrum of compound 4.




Figure S4. ^1H NMR spectrum of compound 6.




Figure S5. ^1H NMR spectrum of compound 10.




Figure S6. Mass spectrum of compound **10**.




Figure S7. ^1H NMR spectrum of compound **13**.




Figure S8. Mass spectrum of compound 13.