

Novel ynamide structural analogues and their synthetic transformations

Ting Lu* and Richard P. Hsung*

*Division of Pharmaceutical Sciences, School of Pharmacy, University of Wisconsin, Madison
Wisconsin 53705, United States*

E-mail: luting80@yahoo.com, rhsung@wisc.edu

DOI: <http://dx.doi.org/10.3998/ark.5550190.p008.512>

Abstract

This Highlight accounts for a recent phenomenon in which a series of novel ynamide structural analogues have emerged and caught the attention of the synthetic community. Preparations and reactions of these de novo ynamide variants are delineated here to demonstrate their accessibility as well as their reactivity. This Highlight should help reveal that these unique *N*-containing alkynes can become highly versatile building blocks in organic syntheses.


Keywords: Ynimides, yne-hydrazides, amidinyl-ynamides, yne-sulfoximines, yne-imines, diaminoacetylenes

Table of Contents

1. Introduction: Ynamide Analogues
2. Discussions: Syntheses and Reactions
 - 2.1 Diaminoacetylenes
 - 2.2 Ynimides
 - 2.3 Yne-imines
 - 2.4 Yne-hydrazides
 - 2.5 Amidinyl-ynamides
 - 2.6 Yne-sulfoximines
3. Conclusions

1. Introduction: Ynamide Analogues

Ynamides **1** (Scheme 1), the electron-deficient alternatives of the important but labile ynamines,¹⁻¹⁴ represent one of the most significant and versatile *N*-containing building blocks in organic synthesis.¹⁵⁻¹⁷ Particularly, in the past 15 years, interests in this versatile building block have raised dramatically.¹⁸⁻²¹ This boom is closely related to the fact that rapid development of efficient synthesis methods²²⁻²⁶ has rendered ynamides highly accessible. Very recently, preparations of structural analogues of ynamides have emerged as novel *N*-containing alkyne building blocks.²⁷⁻³⁶ These preparations mainly are based on modified methods for ynamide synthesis²²⁻²⁶ as described in Scheme 1. With the belief that these novel analogues will also become versatile synthons that will be tremendously useful to organic synthesis, especially in the arena of *N*-heterocycle constructions, we highlight herein syntheses and reactions of *de novo* ynamide analogues: diaminoacetylenes **2**, ynimides **3**, yne-imines **4**, yne-hydrazides **5**, amidinyl-ynamides **6**, and yne-sulfoximines **7** (Scheme 1).




Scheme 1

2. Discussions: Syntheses and Reactions

2.1 Diaminoacetylenes


Tamm²⁷ described a novel approach to access diaminoacetylenes **10** by Fristch-Buttenberg-Wiechell (FBW) rearrangement³⁷⁻³⁹ of 1,1-dihalo-2,2-ethenediamines **8** via the LiBr-bound carbenoids **9** (Scheme 2). This preparation of diaminoacetylenes **10** led to their X-ray structures for the first time, thereby revealing a virtually perpendicular orientations for the two NC₃ planes (See *P1* and *P2* in box at right in Scheme 2).²⁷



Scheme 2

With Brønsted acid or Lewis acid serving as catalysts, diaminoacetylene **10-a** would undergo an intramolecular hydroarylation to afford indole **11**. It is noteworthy that with polar solvent such as THF, **8-a** is not subjected to FBW rearrangement. Instead, indole **13** was formed via an intramolecular 1,5 C-H insertion of the vinylidene intermediate **12**.²⁷


The successful synthesis of 1,2-dipiperidinoacetylene **10-b** enabled studies of coordination chemistry on diaminoacetylenes through preparations of a series of structurally interesting metal complexes such as monometallic complex **14**, homobimetallic complex **15**, and heterobimetallic complex **16** (Scheme 3).⁴⁰



Scheme 3


2.2 Ynimides

Sueda²⁸ reported an efficient synthesis of ynimides **19** via a copper catalyzed amidation of alkynyl(trialkyl)bismuthonium salts **17**. An application of these *de novo* *N*-ethynyl phthanlimides was demonstrated through a copper catalyzed [3 + 2] cycloaddition followed by hydrazinolysis, leading to 4-amino-1,2,3-triazole **22**. This transformation represents as an alternative preparation of **22** to the existing [3 + 2] cycloaddition using the highly labile ethynamine.




Scheme 4

In addition, with the aid of an Au(I) catalyst, which is a preferred π electrophilic Lewis acid, ynimides **19** reacted with methanol to afford β -ketoimides **23**. On the other hand, with Ag(I) catalyst, oxazoles **24** were formed through an initial coordination of Ag(I) with the carbonyl oxygen (see **27**). Similarly, **19-e** could react with aniline, under Au(I)/Ag(I) or Ag(I) only catalysis, to afford phthalimide **29** and oxazolylbenzamide **30**, respectively (Scheme 5).²⁹



Scheme 5




Scheme 6

Muñiz³⁰ later demonstrated a metal-free amination of aryl substituted terminal alkynes **31** using hypervalent iodine reagent **32** to synthesize ynimides **33**. When treating the aniline derived acetylene **36** with **32**, benzo-1,3-oxazine **37** was obtained likely through the ynimide intermediate **38** (Scheme 6).

2.3 Yne-imines

Evano³¹ documented syntheses of yne-iminies **41** through copper catalyzed oxidative cross-couplings of terminal alkynes **39** with diaryl imines **40**. By using Lindlar's catalyst, these *de novo* yne-imines **41** could be reduced to afford *Z*-azadienes **42**. On the other hand, the addition of MeLi induced dimerization of **41** led to the highly substituted azines **43** (Scheme 7).




Scheme 7


2.4 Yne-hydrazides

Batey³² reported a concise synthesis of yne-hydrazides **46** through a straightforward addition of lithiated acetylenes **44** to diazodicarboxylates **45** (Scheme 8). These novel yne-hydrazides were shown as useful building blocks in a series of interesting transformations. In addition to a Larock-type indole formation (Scheme 8),⁴¹ Batey showcased syntheses of structurally unique heterocycles such as **48**, **50**, and **52** through a sequence of [3 + 2] cycloaddition followed by

condensation under acidic conditions (Scheme 9). It is noteworthy of the complete regioselectivity switch in the [3 + 2] cycloaddition step between using $\text{Cp}^*\text{RuCl}(\text{COD})$ and $\text{Cu}(\text{OAc})_2$ (see **47** versus **49**),⁴² albeit the former used an internal yne-hydrazide, while the latter employed a terminal yne-hydrazide.




Scheme 8



Scheme 9

Recently, Batey³³ utilized yne-hydrazide **46-f** to synthesize *Z*-ene-hydrazide **54**, which


represent an analogue of biologically active natural product hydrazidomycin A⁴³ (Scheme 10). The key Lindlar's hydrogenation approach is similar to that of Hsung's Z-enamides syntheses from ynamides.⁴⁴



Scheme 10

2.5 Amidinyl-ynamides

Neuville³⁴ reported syntheses of 1,2,4-trisubstituted imidazoles **58** from terminal alkynes **55** and amidines **56**. Amidinyl-ynamides **57** are believed to be the intermediates resulting from a copper catalyzed oxidative amination of **55** with **56** (Scheme 11).




Scheme 11


2.6 Yne-sulfoximines

Bolm³⁵ described the first preparations of yne-sulfoximines **61** through a copper catalyzed oxidative cross-coupling of sulfoximines **59** with terminal alkynes **60**. These yne-sulfoximines

61 could be transformed to *N*-acyl sulfoximines **62** via simple hydrolysis upon silica gel column chromatography (Scheme 12).



Scheme 12



Scheme 13

Recognizing the excellent potential of yne-sulfoximines **61** as new synthetic building blocks, Bolm's group also explored their usage in thermal [2 + 2] cycloaddition reactions.³⁶ They found that yne-sulfoximines **61** could react with ketenes **65** to form the sulfoximine derived cyclobutenones **63** via a [2 + 2] cycloaddition pathway. When using diphenyl ketene **65-e**, allenyl amide **66** was also isolated in addition to the expected cyclobutenone **63-e**. The formation of **66** is likely a result of pericyclic ring opening of the oxetene intermediate **67**,⁴⁵⁻⁵⁴ which could be envisioned through a hetero-[2 + 2] pathway (Scheme 13).

3. Conclusions

This Highlight presents a recent phenomenon in developing syntheses and reactions of novel structural analogues of ynamides. While their syntheses and reactivities are similar to those of ynamides in many aspects, their unique motifs have rendered them special merits in terms of reactivities. These *N*-containing alkynes should become new versatile building blocks in organic synthesis.

Acknowledgements

Authors thank NIH [GM066055] for funding.

References

1. Bode, J. *Ann.* **1892**, 267, 268.
2. Klages, F.; Drerup, E. *Ann.* **1941**, 547, 65.
3. Zaugg, H. E.; Swett, L. R.; Stone, G. R. *J. Org. Chem.* **1958**, 23, 1389.
<http://dx.doi.org/10.1021/jo01103a617>
4. Wolf, V.; Kowitz, F. *Ann.* **1960**, 638, 33.
5. Viehe, H. G. *Angew Chem. Int. Ed.* **1963**, 2, 477.
<http://dx.doi.org/10.1002/anie.196304771>
6. Viehe, H. G. *Angew Chem. Int. Ed.* **1967**, 6, 767.
<http://dx.doi.org/10.1002/anie.196707671>
7. Viehe, H. G. *Chemistry of Acetylenes*. Chapter 12. Marcel Dekker: New York, 1969; pp 861-912.
8. Ficini, J. *Tetrahedron* **1976**, 32, 448.
[http://dx.doi.org/10.1016/0040-4020\(76\)85200-3](http://dx.doi.org/10.1016/0040-4020(76)85200-3)
9. Pitacco, G.; Valentin, E., in *Chemistry of Functional Groups*. Chapter 15, 1979; pp623-714.
10. Collard-Motte, J.; Janousek, Z. *Topics in Current Chem.* **1986**, 130, 89.
http://dx.doi.org/10.1007/3-540-15810-3_3

11. Himbert, G. in *Methoden Der Organischen Chemie (Houben-Weyl)*, Kropf, H.; Schaumann, E., Eds. Georg Thieme Verlag: Stuttgart, 1993; pp3267.
[Schaumann, E., Eds. Georg Thieme Verlag: Stuttgart, 1993, pp3267.](#)
12. Zifcsak, C. A.; Mulder, J. A.; Rameshkumar, C.; Wei, L.-L.; Hsung, R. P. *Tetrahedron* **2001**, *57*, 7575.
[http://dx.doi.org/10.1016/S0040-4020\(01\)00681-0](http://dx.doi.org/10.1016/S0040-4020(01)00681-0)
13. Mulder, J. A.; Kurtz, K. C. M.; Hsung, R. P. *Synlett* **2003**, 1379.
14. Katritzky, A. R.; Jiang, R.; Singh, S. K. *Heterocycles* **2004**, *63*, 1455.
<http://dx.doi.org/10.3987/REV-04-575>
15. DeKorver, K. A.; Li, H.; Lohse, A. G.; Hayashi, R.; Lu, Z.; Zhang, Y.; Hsung, R. P. *Chem. Rev.* **2010**, *110*, 5064.
<http://dx.doi.org/10.1021/cr100003s>
PMid:20429503 PMCID:PMC2927719
16. Evano, G.; Coste, A.; Jouvin, K. *Angew. Chem., Int. Ed.* **2010**, *49*, 2840.
<http://dx.doi.org/10.1002/anie.200905817>
PMid:20354981
17. Wang, X-N.; Yeom, H-S.; Fang, L-C.; He, S.; Ma, Z-X.; Kedrowski, B. L.; Hsung, R. P. *Acc. Chem. Res.* **2013**, DOI: 10.1021/ar400193g.
<http://dx.doi.org/10.1021/ar400193g>
18. Ackermann, L.; Potukuchi, H. K. *Org. Biomol. Chem.* **2010**, *8*, 4503.
<http://dx.doi.org/10.1039/c0ob00212g>
PMid:20733972
19. Domínguez, G.; Perez-Castells, J. *Chem. Soc. Rev.* **2011**, *40*, 3430.
20. Weding, N.; Hapke, M. *Chem. Soc. Rev.* **2011**, *40*, 4525.
<http://dx.doi.org/10.1039/c0cs00189a>
PMid:21594299
21. Madelaine, C.; Valerio, V.; Maulide, N. *Chem.: Asian J.* **2011**, *6*, 2224.
<http://dx.doi.org/10.1002/asia.201100108>
PMid:21721132
22. Mulder, J. A.; Kurtz, K. C. M.; Hsung, R. P. *Synlett* **2003**, 1379.
23. Dehli, J. R.; Legros, J.; Bolm, C. *Chem. Commun.* **2005**, *43*, 973.
<http://dx.doi.org/10.1039/b415954c>
PMid:15719090
24. Tracey, M. R.; Hsung, R. P.; Antoline, J. A.; Kurtz, K. C. M.; Shen, L.; Slafer, B. W.; Zhang, Y. In *Science of Synthesis*, Houben-Weyl Methods of Molecular Transformations; Weinreb, S. M., Ed.; Georg Thieme Verlag KG: Stuttgart: Germany, 2005; Chapter 21.4.
25. Evano, G.; Blanchard, N.; Mathieu Toumi, M. *Chem. Rev.* **2008**, *108*, 3054.
<http://dx.doi.org/10.1021/cr8002505>
PMid:18698737
26. Evano, G.; Jouvin, K.; Coste, A. *Synthesis* **2013**, 17.

27. Petrov, A. R.; Daniliuc, C. G.; Jones, P. G.; Tamm, M. *Chem.: Eur. J.* **2010**, *16*, 11804.
<http://dx.doi.org/10.1002/chem.201002211>
PMid:20839193
28. Sueda, T.; Oshima, A.; Teno, N. *Org. Lett.* **2011**, *13*, 3996.
<http://dx.doi.org/10.1021/ol2014973>
PMid:21710984
29. Sueda, T.; Kawada, A.; Urashi, Y.; Teno, N. *Org. Lett.* **2013**, *15*, 1560.
<http://dx.doi.org/10.1021/ol400338x>
PMid:23496249
30. Souto, J. A.; Becker, P.; Iglesias, A.; Muñiz, K. *J. Am. Chem. Soc.* **2012**, *134*, 15505.
<http://dx.doi.org/10.1021/ja306211q>
PMid:22909000
31. Laouiti, A.; Rammah, M. M.; Rammah, M. B.; Marrot, J.; Couty, F.; Evano, G. *Org. Lett.* **2012**, *14*, 6.
<http://dx.doi.org/10.1021/ol2032152>
PMid:22171540
32. Beveridge, R. E.; Batey, R. A. *Org. Lett.* **2012**, *14*, 540.
<http://dx.doi.org/10.1021/ol2031608>
PMid:22216967
33. Beveridge, R. E.; Batey, R. A. *Org. Lett.* **2013**, *15*, 3086.
<http://dx.doi.org/10.1021/ol401275f>
PMid:23734814
34. Li, J.; Neuville, L. *Org. Lett.* **2013**, *15*, 1752.
<http://dx.doi.org/10.1021/ol400560m>
PMid:23528104
35. Wang, L.; Huang, H.; Priebbenow, D. L.; Pan, F.-F.; Bolm, C. *Angew. Chem., Int. Ed.* **2013**, *52*, 3478.
<http://dx.doi.org/10.1002/anie.201209975>
PMid:23404753
36. Pirwerdjan, R.; Priebbenow, D. L.; Becker, P.; Lamers, P.; Bolm, C. *Org. Lett.* **2013**, *15*, 5397.
<http://dx.doi.org/10.1021/ol4026028>
PMid:24151931
37. P. Fritsch, *Justus Liebigs Ann. Chem.* **1894**, 279, 319.
<http://dx.doi.org/10.1002/jlac.18942790310>
38. W. P. Bottenberg, *Justus Liebigs Ann. Chem.* **1894**, 279, 324.
<http://dx.doi.org/10.1002/jlac.18942790311>
39. H. Wiechell, *Justus Liebigs Ann. Chem.* **1894**, 279, 337.
<http://dx.doi.org/10.1002/jlac.18942790312>

40. Petrov, A. R.; Bannenberg, T.; Daniliuc, C. G.; Jones, P. G.; Tamm, M. *Dalton Transactions*. **2011**, 40, 10503.
<http://dx.doi.org/10.1039/c1dt10606f>
PMid:21842057
41. Larock, R. C.; Yum, E. K. *J. Am. Chem. Soc.* **1991**, 113, 6689.
<http://dx.doi.org/10.1021/ja00017a059>
42. Boren, B. C.; Naravan, S.; Rasmussen, L. K.; Zhang, L.; Zhao, H.; Lin, Z.; Jia, G.; Fokin, V. *J. Am. Chem. Soc.* **2008**, 130, 8923.
<http://dx.doi.org/10.1021/ja0749993>
PMid:18570425
43. Ueberschaar, N.; Le Sage Tchize Ndejoung, B.; Ding, L.; Maier, A.; Fiebig, H.-H.; Hertweck, C. *Bioorg. Med. Chem. Lett.* **2011**, 21, 5839.
<http://dx.doi.org/10.1016/j.bmcl.2011.07.108>
PMid:21868221
44. Zhang, X.; Zhang, Y.; Huang, J.; Hsung, R. P.; Kurtz, K. C. M.; Oppenheimer, J.; Petersen, M. E.; Sagamanova, I. K.; Shen, L.; Tracey, M. R. *J. Org. Chem.* **2006**, 71, 4170.
<http://dx.doi.org/10.1021/jo060230h>
PMid:16709057
45. Shindo, M. *Synthesis* **2003**, 2275.
<http://dx.doi.org/10.1055/s-2003-42417>
46. Cheng, D.; Ling, F.; Li, Z.; Yao, W.; Ma, C. *Org. Lett.* **2012**, 14, 3146.
<http://dx.doi.org/10.1021/ol3012277>
PMid:22671022
47. Aikawa, K.; Hioki, Y.; Shimizu, N.; Mikami, K. *J. Am. Chem. Soc.* **2011**, 133, 20092.
<http://dx.doi.org/10.1021/ja2085299>
PMid:22070285
48. Saito, A.; Kasai, J.; Odaira, Y.; Fukaya, H.; Hanzawa, Y. *J. Org. Chem.* **2009**, 74, 5644.
<http://dx.doi.org/10.1021/jo900857c>
PMid:19496542
49. Saito, A.; Umakoshi, M.; Yagyu, N.; Hanzawa, Y. *Org. Lett.* **2008**, 10, 1783.
<http://dx.doi.org/10.1021/ol800539a>
PMid:18396889
50. You, L.; Al-Rashid, Z. F.; Figueroa, R.; Ghosh, S. K.; Li, G.; Lu, T.; Hsung, R. P. *Synlett* **2007**, 1656.
51. Kurtz, K. C. M.; Hsung, R. P.; Zhang, Y. *Organic Lett.* **2006**, 8, 231.
<http://dx.doi.org/10.1021/ol052487s>
PMid:16408882
52. Mori, S.; Shindo, M. *Org. Lett.* **2004**, 6, 3945.
<http://dx.doi.org/10.1021/ol048499f>
PMid:15496070

53. Shindo, M.; Sato, Y.; Yoshikawa, T.; Koretsune, R.; Shishido, K. *J. Org. Chem.* **2004**, *69*, 3912.

<http://dx.doi.org/10.1021/jo0497813>

PMid:15153025

54. Shindo, M.; Matsumoto, K.; Mori, S.; Shishido, K. *J. Am. Chem. Soc.* **2002**, *124*, 6840.

<http://dx.doi.org/10.1021/ja026275r>

Authors' Biographies



Ting Lu obtained her B.S. in Chemistry from Nankai University in China in 2002. She then went to National University of Singapore to obtain her M.S. degree in Materials Science in 2005. After that, she joined Professor Richard Hsung's research group at the University of Minnesota at Twin Cities and then the University of Wisconsin at Madison, where she studied the methodology development in cyclopropanations of enamides and allenamides, and got her Ph.D. degree in Organic Chemistry in 2009. Now she is working on new catalytic systems for the production of biorenewable chemicals from biomass at the Institute of Bioengineering and Nanotechnology, Agency for Science, Technology and Research, Singapore.



Richard P. Hsung obtained his B.S. in Chemistry and Mathematics from Calvin College in

Grand Rapids, MI. He then attended The University of Chicago and received his M.S. and Ph.D. degrees in Organic Chemistry, respectively, under the supervision of Professors Jeff Winkler and Bill Wulff. After pursuing a postdoctoral stay with Professor Larry Sita in Chicago and NIH-postdoctoral work with Professor Gilbert Stork at Columbia University, he moved to the University of Minnesota at Twin Cities as an Assistant Professor in 1997 and was promoted to Associate Professor in 2002. He was promoted to Professor and moved to University of Wisconsin at Madison in 2006. He was a recipient of the Camille Dreyfus Teacher-Scholar Award and the National Science Foundation Career Award. He has coauthored over 200 publications, delivered over 200 invited lectures, and supervised over 150 students and postdoctoral fellows with research interests in developing cycloaddition and annulation approaches to natural product syntheses and stereoselective methods using allenamides, ynamides, enamides, and cyclic acetals.