





# Supplementary Material



## Toward accessible 3*H*-Pyrroles



**Boris A. Trofimov,\* Marina Yu. Dvorko, Dmitrii A. Shabalin, Elena Yu. Schmidt**


*Laboratory of Unsaturated Heteroatomic Compounds*  
*A.E. Favorsky Irkutsk Institute of Chemistry, Siberian Branch, Russian Academy of Sciences*  
*1 Favorsky Str., 664033 Irkutsk, Russia*  
*E-mail: [boris\\_trofimov@irioch.irk.ru](mailto:boris_trofimov@irioch.irk.ru)*

**$^1\text{H}$  and  $^{13}\text{C}$  NMR spectra**


<sup>1</sup>H NMR spectrum of **2b** (CDCl<sub>3</sub>)<sup>13</sup>C NMR spectrum of **2b** (CDCl<sub>3</sub>)





<sup>1</sup>H NMR spectrum of **2d** (CDCl<sub>3</sub>)<sup>13</sup>C NMR spectrum of **2d** (CDCl<sub>3</sub>)




<sup>1</sup>H NMR spectrum of 2e (CDCl<sub>3</sub>)




<sup>13</sup>C NMR spectrum of 2e (CDCl<sub>3</sub>)

<sup>1</sup>H NMR spectrum of **2f** (CDCl<sub>3</sub>)<sup>13</sup>C NMR spectrum of **2f** (CDCl<sub>3</sub>)



<sup>1</sup>H NMR spectrum of **4e** (mixture with **2e**, CDCl<sub>3</sub>)



<sup>13</sup>Cjmod NMR spectrum of **4e** (mixture with **2e**, CDCl<sub>3</sub>)



