

## Supplementary Material

# An easy synthesis of diversely functionalized 2H-chromenes and amido amines by an enol-Ugi reaction

Ana G. Neo,\* Teresa G. Castellano, Carlos F. Marcos\*





*Laboratory of Bioorganic Chemistry & Membrane Biophysics, School of Veterinary Sciences,  
University of Extremadura, 10071 Cáceres, Spain*

*E-mail: [cfernand@unex.es](mailto:cfernand@unex.es), [aneo@unex.es](mailto:aneo@unex.es)*




### Table of Contents




<sup>1</sup> H-NMR of compound <b>3</b>	S2
<sup>13</sup> C-NMR of compound <b>3</b>	S2
<sup>1</sup> H-NMR of compound <b>6a</b>	S3
<sup>13</sup> C-NMR of compound <b>6a</b>	S3
<sup>1</sup> H-NMR of compound <b>6b</b>	S4
<sup>13</sup> C-NMR of compound <b>6b</b>	S4
<sup>1</sup> H-NMR of compound <b>6c</b>	S5
<sup>13</sup> C-NMR of compound <b>6c</b>	S5
<sup>1</sup> H-NMR of compound <b>6d</b>	S6
<sup>13</sup> C-NMR of compound <b>6d</b>	S6
<sup>1</sup> H-NMR of compound <b>6e</b>	S7
<sup>13</sup> C-NMR of compound <b>6e</b>	S7
<sup>1</sup> H-NMR of compound <b>6f</b>	S8
<sup>13</sup> C-NMR of compound <b>6f</b>	S8
<sup>1</sup> H-NMR of compound <b>6g</b>	S9
<sup>13</sup> C-NMR of compound <b>6g</b>	S9
<sup>1</sup> H-NMR of compound <b>8a</b>	S10
<sup>13</sup> C-NMR of compound <b>8a</b>	S10
<sup>1</sup> H-NMR of compound <b>8b</b>	S11
<sup>13</sup> C-NMR of compound <b>8b</b>	S11
<sup>1</sup> H-NMR of compound <b>8c</b>	S12
<sup>13</sup> C-NMR of compound <b>8c</b>	S12
<sup>1</sup> H-NMR of compound <b>7a</b>	S13
<sup>13</sup> C-NMR of compound <b>7a</b>	S13
<sup>1</sup> H-NMR of compound <b>7b</b>	S14
<sup>13</sup> C-NMR of compound <b>7b</b>	S14
<sup>1</sup> H-NMR of compound <b>7c</b>	S15
<sup>13</sup> C-NMR of compound <b>7c</b>	S15




## Methyl 2-(4-hydroxy-2-oxo-2H-chromen-3-yl)-2-oxoacetate (3)






## Methyl 2-(4-((2-(cyclohexylamino)-2-oxo-1-phenylethyl)(phenyl)amino)-2-oxo-2H-chromen-3-yl)-2-oxoacetate (6a)







Methyl 2-(4-((2-*tert*-butylamino)-2-oxo-1-phenylethyl)(phenyl)amino)-2-oxo-2*H*-chromen-3-yl)-2-oxoacetate (6b)




Methyl 2-(4-((2-(cyclohexylamino)-2-oxo-1-(*p*-tolyl)ethyl)(phenyl)amino)-2-oxo-2H-chromen-3-yl)-2-oxoacetate (6c)




Methyl 2-(4-((2-(benzylamino)-2-oxo-1-(*p*-tolyl)ethyl)(phenyl)amino)-2-oxo-2*H*-chromen-3-yl)-2-oxoacetate (6d)



## Methyl 2-(4-(benzyl(2-(cyclohexylamino)-2-oxo-1-phenylethyl)amino)-2-oxo-1-phenylethyl)-2-oxo-2H-chromen-3-yl)-2-oxoacetate (6e)







Methyl 2-(4-(benzyl(2-(*tert*-butylamino)-2-oxo-1-phenylethyl)amino)-2-oxo-2*H*-chromen-3-yl)-2-oxoacetate (6f)





**Methyl 2-(4-((benzo[d][1,3]dioxol-5-ylmethyl)(2-(cyclohexylamino)-2-oxo-1-phenylethyl)amino)-2-oxo-2H-chromen-3-yl)-2-oxoacetate (6g)**

**Dimethyl 4-((2-(cyclohexylamino)-2-oxo-1-phenylethyl)(phenyl)amino)-2-hydroxy-2H-chromene-2,3-dicarboxylate (8a)**

**Dimethyl 4-((2-(*tert*-butylamino)-2-oxo-1-phenylethyl)(phenyl)amino)-2-hydroxy-2*H*-chromene-2,3-dicarboxylate (8b)**

**Dimethyl 4-((2-(cyclohexylamino)-2-oxo-1-(*p*-tolyl)ethyl)(phenyl)amino)-2-hydroxy-2*H*-chromene-2,3-dicarboxylate (8c)**

***N*-cyclohexyl-2-phenyl-2-(phenylamino)acetamide (7a)**

***N*-(*tert*-butyl)-2-phenyl-2-(phenylamino)acetamide (7b)**

***N*-cyclohexyl-2-(phenylamino)-2-(*p*-tolyl)acetamide (7c)**