

Supplementary Material

Selective oxidation of sulfides using recyclable pseudocyclic benziodoxole triflate

Akira Yoshimura*, and Viktor V. Zhdankin*



*Department of Chemistry and Biochemistry, University of Minnesota Duluth,
Duluth, Minnesota, 55812, USA*

Emails: ayoshimu@d.umn.edu, vzhdanki@d.umn.edu



Table of Contents

- | | | |
|----|-------------------------|----|
| 1. | NMR spectra of products | S2 |
|----|-------------------------|----|


^1H NMR (500 MHz, CD_3CN)




^1H NMR (500 MHz, CHCl_3)




¹H NMR (500 MHz, CHCl₃)




4b











¹H NMR (500 MHz, CHCl₃)





4c





¹H NMR (500 MHz, CHCl₃)**4d**¹H NMR (500 MHz, CHCl₃)**4e**

¹H NMR (500 MHz, CHCl₃)**4f**¹H NMR (500 MHz, CHCl₃)**4g**



¹H NMR (500 MHz, CHCl₃)





¹H NMR (500 MHz, CHCl₃)





^1H NMR (500 MHz, CHCl_3)





^1H NMR (500 MHz, CHCl_3)







^1H NMR (500 MHz, CHCl_3)





^1H NMR (500 MHz, CHCl_3)





¹H NMR (500 MHz, CHCl₃)¹H NMR (500 MHz, CHCl₃)

^1H NMR (500 MHz, CHCl_3)



^1H NMR (500 MHz, CHCl_3)



^1H NMR (500 MHz, CHCl_3)

