

Tf₂O-Mediated mild synthesis of 6*H*-chromeno[4,3-*b*]quinolines

Li-Ning Chen,[†] Zhao-Ke Jin,[†] Jian-Liang Ye,^{*} and Pei-Qiang Huang^{*}

Fujian Provincial Key Laboratory of Chemical Biology, College of Chemistry and Chemical Engineering, Xiamen University, Xiamen 361005, P. R. China

[†]These authors contributed equally

E-mail: pqhuang@xmu.edu.cn ; yejl@xmu.edu.cn

Dedicated to Prof. Léon Ghosez on the occasion of his 90th birthday

Received mm-dd-yyyy


Accepted Manuscript mm-dd-yyyy

Published on line mm-dd-yyyy

Dates to be inserted by editorial office

Abstract

A facile and versatile synthesis of 6*H*-chromeno[4,3-*b*]quinolines has been achieved through triflic anhydride/2-fluoropyridine (Tf₂O/2-F-Pyr)-promoting domino intramolecular cycloaddition reactions of salicylic-acid-derived *N*-phenyl-*ortho*-propynyloxy benzamides under mild, and metal-free conditions. This protocol is efficient, scalable, and well compatible with a broad scope of substrates bearing halogens, trifluoromethyl, and electron-donating groups.



- mild conditions
 - metal-free
 - scalable
 - readily accessible reagents and starting materials
- 20 examples
up to 97% yield

Keywords: 6*H*-Chromeno[4,3-*b*]quinoline, amide activation, triflic anhydride, intramolecular cycloaddition, tandem reactions.

Introduction

6*H*-Chromeno[4,3-*b*]quinolines represent pharmacological-privileged chromene- and quinoline- fused scaffolds, exhibiting various biological activities.¹⁻⁴ Consequently, they are acknowledged as promising druggable molecules. For example, compounds **1** are selective estrogen-receptor beta (ER β) ligands with high binding IC₅₀ values of 3.3 nM (**1a**), 3.6 nM (**1b**), and 6.1 nM (**1c**), respectively.³ Functional testing carried out on electron-withdrawing NO₂-substituted compound **2** showed good cytotoxicity against murine melanoma cell line B16F10 and female breast-cancer cell line MCF7 with IC₅₀ values of 14.8 and 21.32 μ M, respectively.² Similarly, the tri-methoxy-substituted compound **3** also displayed apoptotic effects on colon cancer cell line HT29 (IC₅₀ = 2.61 μ M).¹ Moreover, recent research indicated that 6*H*-chromeno[4,3-*b*]quinolines could be used as fluorescent probes for selective detection of diverse chemical and biological components,⁵⁻⁸ such as thiols,⁸ pH,^{6,7} lysosomes,⁶ and mitochondria.⁵ For example, Ren and Liu demonstrated that chromenoquinoline **4**, exhibiting low-cytotoxicity, good photostability, and near-infrared emission, could efficiently stain mitochondria and zebrafish for bioimaging.⁵ Lin and coworkers revealed that compound **5** is capable of ratiometric fluorescence, monitoring pH variations from 7.4 to 5.5 in live cells.⁷ Hence, the synthesis of 6*H*-chromeno[4,3-*b*]quinolines has received continuous interest in the fields of organic and pharmaceutical chemistry for a long time.⁹⁻¹⁷





Figure 1. Some bioactive (upper) and luminescent (bottom) 6*H*-Chromeno[4,3-*b*]quinolines


Among the developed methods, one-step construction of pyran and pyridine rings via intramolecular tandem cyclization reactions starting from readily available salicylic acid derivatives, also known as formal aza-Diels-Alder reactions or [4+2] cycloaddition reactions, stand out as an efficient and the most frequently employed strategy. In 1983, Miocque and coworkers synthesized 6*H*-chromeno[4,3-*b*]quinoline by subjecting *ortho*-propynyloxy benzamide to reflux conditions in toluene with POCl₃ for 48 hours (Scheme 1a).¹⁵ This reaction involved an intramolecular cascade cycloaddition between the *in-situ* formed active nitrilium and a proximal alkyne. Alternatively, Ibrahim reported that the nitrilium intermediate was produced by treating *ortho*-propynyloxy benzamide with PCl₅ under reflux conditions in benzene to yield imidoyl chloride, followed by SbCl₅.¹⁴ Also shown in Scheme 1a, the greener Lewis acid-catalyzed imine-alkyne intramolecular tandem

cyclization reaction of *ortho*-propynyloxy benzimines,¹⁶ typically generated *in-situ* from condensation of *O*-propargylated salicylaldehydes and anilines,^{5-7,9,10} represents a more broadly applicable protocol. However, due to relatively low nucleophilicity of the alkyne, the reaction often requires high-temperature conditions or microwave assistance.

a) Reported methods



b) This work



Scheme 1. The representative synthetic strategy of 6*H*-chromeno[4,3-*b*]quinolines

On the other hand, triflic anhydride (Tf₂O), an easily available mild electrophilic reagent, has been utilized to facilitate the conversions of diverse functional groups.¹⁸ Pioneered by Ghosez,^{19,20} direct transformations of amides have been extensively investigated and established as a robust method for selective synthesis.²¹⁻⁴⁸ Of particular interest is the utilization of an intermolecular alkyne-Tf₂O-activated amide cycloaddition for constructing ring structures. Ghosez's initial research demonstrated the smooth formation of cyclobutenones through Tf₂O/colidine-promoted [2+2]-cycloadditions of alkynes with tertiary amides under reflux conditions in CHCl₃.¹⁹ Following this pioneering work, Yao treated an acetylenic aniline amide with Tf₂O/Ph₃PO to generate an imidate intermediate, initiating a cascade intramolecular annulation with an alkyne, to produce indolizinoquinolinones.⁴⁵ Huang and Liang, respectively, employed Tf₂O/2-fluoropyridine (2-F-Pyr) and Tf₂O/2,6-dichloropyridine (2,6-diCl-Pyr) as an activation system to promote the coupling reactions of secondary *N*-aryl secondary amides with alkynes, leading to the concise synthesis of quinolines.^{46,47} It is worthy to note that the reactive nitrilium intermediate generated from Tf₂O/2-F-Pyr was identified by 2D NMR techniques.⁴⁶


In light of the above-mentioned achievements, and in connection with our interest in developing a practical method for the direct transformation of amides,²⁵⁻³² we envisioned to develop a mild, efficient and versatile approach toward 6*H*-chromeno[4,3-*b*]quinoline through a Tf₂O-mediated amide-activation-triggered intramolecular-cascade alkyne-nitrilium cyclization reaction of salicylic-acid-derived *N*-phenyl *ortho*-propynyloxy benzamides (Scheme 1b above). Herein, we present the outcomes of our study.

Results and Discussion

We commenced our investigation by using *N*-phenyl-*ortho*-propynyloxy benzamide **6a** as a model compound to determine the optimal reaction conditions (Table 1). We first examined the conditions that we established

previously for secondary amide activation.⁴⁸ The amide **6a** was treated with Tf₂O (1.1 equiv.) and 2-F-Pyr. (1.2 equiv.) in CH₂Cl₂ (0.1 M) at 0 °C for 15 min, then stirred at room temperature until the complete consumption of the substrate (monitored with TLC, ~8 h). To our delight, the desired product **7a** was afforded smoothly in 82% yield. Encouraged by this result, a screening of bases was conducted, however, the surveyed bases, including 2-Cl-Pyr., 2-Br-Pyr., 2,6-lutidine, 2,4,6-tri-*tert*-butylpyrimidine (TTBP), and pyridine (Pyr), all gave inferior results (entries 2-6). It is worth noting that **7a** could be generated in 50% yield under base-free conditions (entry 7). When running the reaction in 1,2-dichloroethane (DCE), **7a** was produced in the same yield (82%, entry 8). Changing the solvent to toluene resulted in a sharply reduced yield of 38% (entry 9). A slightly higher yield of 89% was obtained in the solvent CH₃CN (entry 10). Thus, CH₃CN was identified as the solvent of choice for subsequent optimization. Pleasingly, when increasing the concentration from 0.1 to 0.2 M (CH₃CN), **7a** was provided in an excellent isolated yield of 97% (entry 11). Further increase of the concentration to 0.5 M resulted in a loss of yield (90%, entry 12). Thus, the optimized conditions of the intramolecular tandem-cycloaddition reaction of *N*-phenyl *ortho*-propynyloxy benzamide **6a** were defined as follows: the amide **6a** was treated with Tf₂O (1.1 equiv.) and 2-F-Pyr. (1.2 equiv.) in CH₃CN (0.2 M) at 0 °C for 15 min, then stirred at room temperature for 8 h.


Table 1. Optimization of the Reaction Conditions^a



Entry	Base	Solvent	C(M)	Yield (%) ^b
1	2-F-Pyr	DCM	0.1	82%
2	2-Cl-Pyr	DCM	0.1	65%
3	2-Br-Pyr	DCM	0.1	45%
4	2,6-Lutidine	DCM	0.1	19%
5	TTBP ^c	DCM	0.1	32%
6	Pyr	DCM	0.1	22%
7	None	DCM	0.1	50%
8	2-F-Pyr	DCE	0.1	82%
9	2-F-Pyr	Toluene	0.1	38%
10	2-F-Pyr	CH ₃ CN	0.1	89%
11	2-F-Pyr	CH₃CN	0.2	97%^d
12	2-F-Pyr	CH ₃ CN	0.5	90%

^aReaction conditions: To the mixture of amide **6a** (0.5 mmol), the base under investigation (0.60 mmol) in corresponding solvent Tf₂O (0.55 mmol) was added dropwise at 0 °C. After stirring at 0 °C for 15 minutes, the reaction was allowed to warm to room temperature and stirred until the complete consumption of the substrate (monitored with TLC, ~8 h). ^bYields determined by ¹H NMR using 1,3,5-trimethoxybenzene as an internal standard. ^cTTBP=2,4,6-tri-*tert*-butylpyrimidine, 0.30 mmol. ^dIsolated yield.

With optimized conditions in hand, the scope of amides was explored and the results are summarized in Table 2. For *N*-4-halogen-phenyl (F, Cl, Br, and I) benzamide **6b-e**, the corresponding 6*H*-chromeno[4,3-*b*]quinolines **7b-e** were afforded in good yields of 70%, 82%, 71%, and 87%, respectively. In the case of 3,4-dichloro-substituted **6f**, however, the product **7f** was provided in a modest yield of 27%. It is suggested that electron-withdrawing substituents at the *N*-phenyl moiety, therefore, have negative effects on the yield of the reaction, which was further demonstrated by the fact that the reaction of 4-CF₃-substituted **6g** produced the desired product **7g** in only 10% yield. The structure of **7g** was confirmed by single crystal X-ray diffraction analysis.

Table2. Scope of Amides **6**


^aReaction conditions: To the mixture of amide **6** (0.5 mmol), 2-F-Pyr (0.60 mmol) in CH₃CN (2.5 mL) Tf₂O (0.55 mmol) was added dropwise at 0 °C. After stirring for 15 minutes at 0 °C, the reaction was allowed to warm to room temperature and stirred until the complete consumption of the substrate (monitored with TLC, 2~8 h).

^b80 °C, overnight. ^cPerformed on the 6.0 mmol scale, 1.50 g of **7u** was obtained.

When raising the reaction temperature to 80 °C, and prolonging the reaction time to overnight, the yield of **7g** was slightly increased to 27%. Furthermore, no cycloaddition products **7h** and **7i** were observed in the

reactions of the stronger electron-withdrawing methoxycarbonyl (CO₂Et) and nitro (NO₂) substituted benzamides **6h** and **6i** under standard conditions or at 80 °C overnight conditions. Comparatively, the reactions of 4-methyl and 4-*t*-butyl substituted **6j** and **6k** proceeded smoothly to afford the corresponding products **7j** and **7k** in good yields of 76% and 75%, respectively. In the case of the electron-rich 4-methoxy substituted **6l**, the corresponding chromenoquinoline **7l** was produced in an excellent yield of 94%. Unexpectedly, the cycloaddition of 3,4-dimethoxy substituted **6m** resulted in the corresponding product **7m** only in a moderate yield of 57% maybe owing to steric-hindrance. In addition, the reaction was well compatible with the sterically hindered *N*-2-methyl-phenyl benzamide **6n** to yield product **7n** in a slightly lower yield of 62%. For *N*-naphthyl benzamide **6o**, the reaction provided the cycloadduct **7o** in 37% yield. Next, the *N*-phenyl substituted benzamides were examined. Similarly, the reactions of *N*-phenyl 4-halogen (F, Cl, Br) substituted benzamides **6p-6r** gave the corresponding cycloadducts **7p-7r** in good yields of 65%, 79%, 82%, respectively. Delightfully, for 4-CF₃-substituted substrate **6s**, the product **7s** was obtained in a moderate yield of 50% under the standard conditions. Expectedly, the benzamides (**6t** and **6u**) bearing an electron-donating methyl or methoxy group at the *para*-position provided corresponding cycloadducts **7t** and **7u** in excellent yields of 87% and 95%, respectively. The structure of **7g** has been determined by single crystal X-ray diffraction analysis. Additionally, the 6.0 mmol scale reaction of **6u** proceeded smoothly to give chromenoquinoline **7u** (1.50 g) in same yield of 95%.

Based on these results and previous NMR studies,⁴⁶ a plausible mechanism for the Tf₂O/2-F-Pyr-promoting domino intramolecular cycloaddition reaction of *N*-phenyl-*ortho*-propynyloxy benzamide is depicted in Scheme 2. Upon the amide **6** treatment with Tf₂O, the imidate **Int-1** and nitrilium **Int-2** were formed. And the presence of 2-F-Pyr facilitates the complete conversion of **Int-1** into **Int-2**. The highly electrophilic nitrilium **Int-2** couples with alkyne group to give an alkenyl cation intermediate **Int-3**,^{46,49} which is prone to be captured by the proximal *N*-phenyl group to produce, via a Friedel-Crafts cyclization, 6*H*-chromeno[4,3-*b*]quinoline **7** (Path A). Alternatively, **Int-2** has the potential to undergo a [4+2] cycloaddition process to form **Int-4**, ultimately leading to the formation of chromenoquinoline **7** (Path B).



Scheme 2. Plausible mechanism

Conclusions

In summary, we have disclosed a facile and efficient approach towards bio-important and fluorescent 6*H*-chromeno[4,3-*b*]quinolines, via Tf₂O-mediated tandem intramolecular cycloaddition reactions of the easily accessible *N*-phenyl *ortho*-propynyloxy benzamides. This method features good yields, good substrate scope, mild and metal-free reaction conditions, and scalability, thereby providing a viable alternative for the

preparation of functionalized chromenoquinolines from amides without the use of toxic/sensitive reagents such as POCl₃ and PCl₅.

Experimental Section

General. Melting points were determined by a Switzerland Büchi M-560 automatic melting point apparatus. Infrared spectra were measured with a Nicolet Avatar 330 FT-IR spectrometer using film KBr pellet techniques. ¹H NMR and ¹³C NMR spectra were recorded on 400 MHz spectrometer with CDCl₃ as solvent. Chemical shifts (δ) are reported in ppm and respectively referenced to either the internal standard Me₄Si (TMS) or solvent signals (Me₄Si at 0 ppm for ¹H NMR, and CDCl₃ at 77.0 ppm for ¹³C NMR). HRMS spectra were recorded on a Bruker Dalton Esquire 3000 plus mass spectrometer by the ESI method. Unless otherwise stated, reactions were performed in oven-dried glassware under a nitrogen atmosphere using standard Schlenk techniques. Flash-column chromatography was performed with silica gel (300-400 mesh), eluting with EtOAc/petroleum ether. Triflic anhydride (Tf₂O) was distilled over phosphorous pentoxide and was stored under 0 °C for no more than a week before use. All other commercially available compounds were used as received.

General procedure for the synthesis of 6H-chromeno[4,3-b]quinolines. To a flame-dried Schlenk tube were added sequentially an amide (0.5 mmol, 1.0 equiv), 2-F-Pyr (0.60 mmol, 1.2 equiv) and MeCN (2.5 mL) at room temperature, then trifluoromethanesulfonic anhydride (Tf₂O, 0.55 mmol, 1.1 equiv) was added dropwise at 0 °C. After stirring for 15 minutes at 0 °C, the mixture was allowed to warm to room temperature and stirred until the complete consumption of the substrate (monitored with TLC, 2~8 h). The reaction was then quenched with a saturated aqueous NaHCO₃ solution (1.0 mL). The reaction mixture was extracted with CH₂Cl₂ (5 mL × 3). The combined organic layers were washed with brine (5 mL), dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure. The residue was purified by flash column chromatography (elution EtOAc/petroleum ether) on silica gel to afford the desired product.

6H-Chromeno[4,3-b]quinoline (7a). Following the general procedure, the reaction of **6a** (125.5 mg, 0.5 mmol) gave, after FC on silica gel (eluent: EtOAc/petroleum ether = 1:20), compound **7a** as a white floppy solid (113mg, yield: 97%). Mp: 125-127 °C (lit.¹⁶ Mp 64-66 °C ; lit.⁵⁰ Mp 126 °C); ¹H NMR (400 MHz, CDCl₃) δ 8.49 (dd, *J* 7.7, 1.7 Hz, 1H), 8.13 (d, *J* 8.4 Hz, 1H), 7.87 (s, 1H), 7.77 (d, *J* 8.2 Hz, 1H), 7.73–7.64 (m, 1H), 7.50 (t, *J* 7.5 Hz, 1H), 7.42–7.33 (m, 1H), 7.17 (t, *J* 7.4 Hz, 1H), 7.02 (d, *J* 8.2 Hz, 1H), 5.36 (s, 2H) ppm; ¹³C NMR (100 MHz, CDCl₃) δ 157.65, 149.27, 148.59, 132.11, 131.17, 129.77, 129.66, 127.82, 127.66, 126.52, 125.79, 125.48, 123.50, 122.79, 117.57, 68.65 ppm; HRMS (ESI) *m/z*: calcd for C₁₆H₁₂NO⁺ [*M* + *H*]⁺: 234.0913, found: 234.0913.

9-Fluoro-6H-chromeno[4,3-b]quinoline (7b). Following the general procedure, the reaction of **6b** (134.5 mg, 0.5 mmol) gave, after FC on silica gel (eluent: EtOAc/petroleum ether = 1:20), compound **7b** as a white floppy solid (88 mg, yield: 70%). Mp: 118-121 °C (lit.⁵⁰ Mp 99-101 °C); ¹H NMR (400 MHz, CDCl₃) δ 8.44 (dd, *J* 7.8, 1.9 Hz, 1H), 8.11 (dd, *J* 9.2, 5.4 Hz, 1H), 7.79 (s, 1H), 7.45 (td, *J* 8.8, 2.7 Hz, 1H), 7.41–7.33 (m, 2H), 7.16 (t, *J* 7.5 Hz, 1H), 7.02 (d, *J* 8.2 Hz, 1H), 5.34 (s, 2H) ppm; ¹³C NMR (100 MHz, CDCl₃) δ 160.58 (d, *J*_{C-F} = 248.0 Hz), 157.51, 148.72, 145.63, 132.15 (d, *J*_{C-F} = 4.5 Hz), 132.00, 130.47 (d, *J*_{C-F} = 5.3 Hz), 128.34 (d, *J*_{C-F} = 9.9 Hz), 126.31, 125.60, 123.24, 122.84, 119.87 (d, *J*_{C-F} = 25.6 Hz), 117.59, 110.79 (d, *J*_{C-F} = 21.9 Hz), 68.51 ppm; HRMS (ESI) *m/z*: calcd for C₁₆H₁₁FNO⁺ [*M* + *H*]⁺: 252.0819, found: 252.0819.

9-Chloro-6H-chromeno[4,3-b]quinoline (7c). Following the general procedure, the reaction of **6c** (142.5 mg, 0.5 mmol) gave, after FC on silica gel (eluent: EtOAc/petroleum ether = 1:20), compound **7c** as a pale-yellow

crystalline solid (109 mg, yield: 82%). ^1H NMR (400 MHz, CDCl_3) δ 8.42 (dd, J 7.8, 1.7 Hz, 1H), 8.02 (d, J 9.0 Hz, 1H), 7.72–7.65 (m, 2H), 7.59 (dd, J 9.0, 2.3 Hz, 1H), 7.41–7.33 (m, 1H), 7.15 (t, J 7.4 Hz, 1H), 7.00 (d, J 8.1 Hz, 1H), 5.30 (s, 2H) ppm; ^{13}C NMR (100 MHz, CDCl_3) δ 157.58, 149.47, 146.87, 132.33, 132.07, 131.15, 130.58, 130.09, 128.29, 126.33, 126.29, 125.71, 123.08, 122.82, 117.59, 68.41 ppm; HRMS (ESI) m/z : calcd for $\text{C}_{16}\text{H}_{11}\text{ClNO}^+ [\text{M} + \text{H}]^+$: 268.0524, found: 268.0522.

9-Bromo-6H-chromeno[4,3-*b*]quinoline (7d). Following the general procedure, the reaction of **6d** (164.5 mg, 0.5 mmol) gave, after FC on silica gel (eluent: EtOAc/petroleum ether = 1:15), compound **7d** as a yellow floppy solid (110 mg, yield: 71%). Mp: 142–144 °C (lit.¹⁶ Mp 137–139 °C; lit.⁵⁰ Mp 138 °C); ^1H NMR (400 MHz, CDCl_3) δ 8.42 (dd, J 7.8, 1.7 Hz, 1H), 7.95 (d, J 8.9 Hz, 1H), 7.86 (d, J 2.2 Hz, 1H), 7.74–7.66 (m, 2H), 7.39–7.35 (m, 1H), 7.15 (td, J 7.5, 1.2 Hz, 1H), 7.00 (dd, J 8.2, 1.1 Hz, 1H), 5.30 (s, 2H) ppm; ^{13}C NMR (100 MHz, CDCl_3) δ 157.62, 149.60, 147.07, 133.13, 132.39, 131.26, 130.01, 129.63, 128.82, 126.32, 125.76, 123.06, 122.83, 120.24, 117.61, 68.41 ppm; HRMS (ESI) m/z : calcd for $\text{C}_{16}\text{H}_{11}\text{BrNO}^+ [\text{M} + \text{H}]^+$: 312.0019, found: 312.0019.

9-Iodo-6H-chromeno[4,3-*b*]quinoline (7e). Following the general procedure, the reaction of **6e** (188.5 mg, 0.5 mmol) gave, after FC on silica gel (eluent: EtOAc/petroleum ether = 1:15), compound **7e** as a pale-yellow crystalline solid (156 mg, yield: 87%). Mp: 138–141 °C; ^1H NMR (400 MHz, CDCl_3) δ 8.40 (dd, J 7.8, 1.5 Hz, 1H), 8.08 (t, J 1.5 Hz, 1H), 7.87 (dt, J 8.9, 1.6 Hz, 1H), 7.79 (d, J 8.9 Hz, 1H), 7.64 (s, 1H), 7.40–7.33 (m, 1H), 7.14 (t, J 7.5 Hz, 1H), 6.99 (d, J 8.2 Hz, 1H), 5.28 (s, 2H) ppm; ^{13}C NMR (100 MHz, CDCl_3) δ 157.62, 149.69, 147.38, 138.39, 136.28, 132.43, 131.18, 129.78, 129.37, 126.13, 125.79, 123.00, 122.82, 117.60, 91.83, 68.39 ppm; HRMS (ESI) m/z : calcd for $\text{C}_{16}\text{H}_{11}\text{INO}^+ [\text{M} + \text{H}]^+$: 359.9880, found: 359.9878.

9,10-Dichloro-6H-chromeno[4,3-*b*]quinoline (7f). Following the general procedure, the reaction of **6f** (159.5 mg, 0.5 mmol) gave, after FC on silica gel (eluent: EtOAc/petroleum ether = 1:50), compound **7f** as a white solid (41 mg, yield: 27%). Mp: 166–168 °C; IR (film) ν_{max} : 2917, 1607, 1580, 1471, 1239, 1150, 1037, 915, 748 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 8.41 (dd, J 7.8, 1.8 Hz, 1H), 8.22 (s, 1H), 7.84 (s, 1H), 7.73 (s, 1H), 7.42–7.37 (m, 1H), 7.16 (t, J 7.6 Hz, 1H), 7.01 (d, J 8.2 Hz, 1H), 5.32 (s, 2H) ppm; ^{13}C NMR (100 MHz, CDCl_3) δ 157.74, 150.46, 147.16, 134.11, 132.72, 130.81, 130.66, 129.84, 128.08, 126.78, 126.54, 125.86, 122.92, 122.81, 117.68, 68.37 ppm; HRMS (ESI) m/z : calcd for $\text{C}_{16}\text{H}_{10}\text{Cl}_2\text{NO}^+ [\text{M} + \text{H}]^+$: 302.0134, found: 302.0134.

9-(Trifluoromethyl)-6H-chromeno[4,3-*b*]quinoline (7g). Following the general procedure, the reaction of **6g** (159.5 mg, 0.5 mmol) gave, after FC on silica gel (eluent: EtOAc/petroleum ether = 1:20), compound **7g** as a pale-yellow solid (41 mg, yield: 27%). Mp: 147–149 °C (lit.¹⁶ Mp 122–124 °C); ^1H NMR (400 MHz, CDCl_3) δ 8.43 (d, J 7.8 Hz, 1H), 8.16 (d, J 8.8 Hz, 1H), 8.00 (s, 1H), 7.82–7.80 (m, 2H), 7.38 (t, J 7.6 Hz, 1H), 7.15 (t, J 7.5 Hz, 1H), 7.00 (d, J 8.2 Hz, 1H), 5.30 (s, 2H) ppm; ^{13}C NMR (100 MHz, CDCl_3) δ 157.85, 151.24, 149.47, 132.83, 131.68, 130.66, 128.06 (q, $J_{\text{C-F}}$ = 32.0 Hz), 126.63, 125.61, 125.99, 125.50 (q, $J_{\text{C-F}}$ = 5.0 Hz), 125.37, 125.33, 124.26 (q, $J_{\text{C-F}}$ = 271.0 Hz) 122.87, 117.67, 68.32 ppm; HRMS (ESI) m/z : calcd for $\text{C}_{17}\text{H}_{11}\text{F}_3\text{NO}^+ [\text{M} + \text{H}]^+$: 302.0787, found: 302.0787.

9-Methyl-6H-chromeno[4,3-*b*]quinoline (7j). Following the general procedure, the reaction of **6j** (132.5 mg, 0.5 mmol) gave, after FC on silica gel (eluent: EtOAc/petroleum ether = 1:15), compound **7j** as a brown-yellow floppy solid (94 mg, yield: 76%). Mp: 125–127 °C; ^1H NMR (400 MHz, CDCl_3) δ 8.45 (dd, J 7.8, 1.6 Hz, 1H), 8.00 (d, J 8.5 Hz, 1H), 7.68 (s, 1H), 7.54–7.41 (m, 2H), 7.39–7.29 (m, 1H), 7.18–7.11 (m, 1H), 7.00 (d, J 8.1 Hz, 1H), 5.28 (s, 2H), 2.49 (s, 3H) ppm; ^{13}C NMR (100 MHz, CDCl_3) δ 157.43, 148.32, 147.09, 136.36, 131.97, 131.78, 130.43, 129.26, 127.77, 126.54, 125.57, 125.35, 123.57, 122.68, 117.46, 68.62, 21.75 ppm; HRMS (ESI) m/z : calcd for $\text{C}_{17}\text{H}_{14}\text{NO}^+ [\text{M} + \text{H}]^+$: 248.1070, found: 248.1070.

9-(*tert*-Butyl)-6H-chromeno[4,3-*b*]quinoline (7k). Following the general procedure, the reaction of **6k** (153.5 mg, 0.5 mmol) gave, after FC on silica gel (eluent: EtOAc/petroleum ether = 1:15), compound **7k** as a brown-yellow solid (108 mg, yield: 75%). Mp: 143–146 °C; IR (film) ν_{max} : 3048, 2978, 2897, 2867, 1584, 1465, 1236,

917, 838, 756 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 8.47 (dd, J 7.8, 1.7 Hz, 1H), 8.05 (d, J 8.9 Hz, 1H), 7.82–7.72 (m, 2H), 7.66 (d, J 2.2 Hz, 1H), 7.40–7.30 (m, 1H), 7.15 (t, J 7.4 Hz, 1H), 7.01 (d, J 8.1 Hz, 1H), 5.32 (s, 2H), 1.42 (s, 9H) ppm; ^{13}C NMR (100 MHz, CDCl_3) δ 157.46, 149.38, 148.58, 147.12, 131.78, 131.10, 129.15, 128.66, 127.50, 125.65, 125.31, 123.68, 122.70, 122.67, 117.48, 68.68, 35.11, 31.40 ppm; HRMS (ESI) m/z : calcd for $\text{C}_{20}\text{H}_{20}\text{NO}^+$ $[\text{M} + \text{H}]^+$: 290.1539, found: 290.1539.

9-Methoxy-6H-chromeno[4,3-b]quinoline (7l). Following the general procedure, the reaction of **6l** (140.5 mg, 0.5 mmol) gave, after FC on silica gel (eluent: EtOAc/petroleum ether = 1:4), compound **7l** as a brown-yellow solid (124 mg, yield: 94%). Mp: 140–143 °C (lit.⁵⁰ Mp 125 °C); ^1H NMR (400 MHz, CDCl_3) δ 8.43 (dd, J 7.8, 1.7 Hz, 1H), 8.02 (d, J 9.2 Hz, 1H), 7.72 (s, 1H), 7.38–7.30 (m, 2H), 7.19–7.11 (m, 1H), 7.04–6.96 (m, 2H), 5.31 (s, 2H), 3.92 (s, 3H) ppm; ^{13}C NMR (100 MHz, CDCl_3) δ 157.92, 157.21, 146.94, 144.59, 131.53, 131.06, 129.92, 128.79, 125.72, 125.35, 123.65, 122.73, 122.29, 117.44, 105.40, 68.65, 55.74 ppm; HRMS (ESI) m/z : calcd for $\text{C}_{17}\text{H}_{14}\text{NO}_2^+$ $[\text{M} + \text{H}]^+$: 264.1019, found: 264.1019.

9,10-Dimethoxy-6H-chromeno[4,3-b]quinoline (7m). Following the general procedure, the reaction of **6m** (155.5 mg, 0.5 mmol) gave, after FC on silica gel (eluent: EtOAc/petroleum ether = 1:15), compound **7m** as a white solid (84 mg, yield: 57%). ^1H NMR (400 MHz, CDCl_3) δ 8.40 (dd, J 7.8, 1.7 Hz, 1H), 7.71 (s, 1H), 7.47 (s, 1H), 7.37–7.30 (m, 1H), 7.15 (t, J 7.5 Hz, 1H), 7.01 (d, J 6.6 Hz, 2H), 5.32 (s, 2H), 4.06 (s, 3H), 4.02 (s, 3H) ppm; ^{13}C NMR (100 MHz, CDCl_3) δ 157.17, 152.68, 149.94, 147.14, 145.56, 131.44, 129.56, 125.18, 123.80, 123.58, 123.32, 122.68, 117.44, 108.38, 105.25, 68.66, 56.39, 56.27 ppm; HRMS (ESI) m/z : calcd for $\text{C}_{18}\text{H}_{16}\text{NO}_3^+$ $[\text{M} + \text{H}]^+$: 294.1125, found: 294.1124.

11-Methyl-6H-chromeno[4,3-b]quinoline (7n). Following the general procedure, the reaction of **6n** (132.5 mg, 0.5 mmol) gave, after FC on silica gel (eluent: EtOAc/petroleum ether = 1:20), compound **7n** as a yellow floppy solid (76 mg, yield: 62%). Mp: 92–95 °C (lit.¹⁶ Mp 72–74 °C); ^1H NMR (400 MHz, CDCl_3) δ 8.54 (dd, J 7.8, 1.8 Hz, 1H), 7.81 (s, 1H), 7.59 (d, J 8.1 Hz, 1H), 7.53 (d, J 7.0 Hz, 1H), 7.39–7.34 (m, 2H), 7.16 (t, J 7.4 Hz, 1H), 7.01 (d, J 8.1 Hz, 1H), 5.36 (s, 2H), 2.88 (s, 3H) ppm; ^{13}C NMR (100 MHz, CDCl_3) δ 157.53, 147.82, 147.47, 137.70, 131.83, 131.27, 129.81, 127.73, 126.25, 125.74, 125.58, 124.92, 123.91, 122.65, 117.48, 68.62, 18.11 ppm; HRMS (ESI) m/z : calcd for $\text{C}_{17}\text{H}_{14}\text{NO}^+$ $[\text{M} + \text{H}]^+$: 248.1070, found: 248.1070.

6H-Benzo[h]chromeno[4,3-b]quinoline (7o). Following the general procedure, the reaction of **6o** (150.5 mg, 0.5 mmol) gave, after FC on silica gel (eluent: EtOAc/petroleum ether = 1:15), compound **7o** as a white floppy solid (52 mg, yield: 37%). ^1H NMR (400 MHz, CDCl_3) δ 8.62 (s, 1H), 8.55 (d, J 8.0 Hz, 1H), 8.49 (d, J 8.0 Hz, 1H), 8.05–7.90 (m, 3H), 7.69–7.60 (m, 2H), 7.41–7.34 (m, 1H), 7.18 (t, J 7.5 Hz, 1H), 7.03 (d, J 8.1 Hz, 1H), 5.46 (s, 2H) ppm; ^{13}C NMR (100 MHz, CDCl_3) δ 157.34, 148.66, 148.46, 131.95, 131.83, 131.04, 129.78, 129.00, 128.50, 127.37, 127.28, 126.38, 125.48, 125.06, 124.80, 123.42, 122.78, 122.68, 117.46, 68.83 ppm; HRMS (ESI) m/z : calcd for $\text{C}_{20}\text{H}_{14}\text{NO}^+$ $[\text{M} + \text{H}]^+$: 284.1070, found: 284.1070.

3-Fluoro-6H-chromeno[4,3-b]quinoline (7p). Following the general procedure, the reaction of **6p** (134.5 mg, 0.5 mmol) gave, after FC on silica gel (eluent: EtOAc/petroleum ether = 1:70), compound **7p** as a pale-yellow floppy solid (82 mg, yield: 65%). Mp: 115–118 °C; ^1H NMR (400 MHz, CDCl_3) δ 8.50–8.41 (m, 1H), 8.09 (d, J 8.5 Hz, 1H), 7.83 (s, 1H), 7.75 (d, J 8.1 Hz, 1H), 7.69 (t, J 7.7 Hz, 1H), 7.48 (t, J 7.5 Hz, 1H), 6.87 (td, J 8.5, 2.5 Hz, 1H), 6.72 (dd, J 9.8, 2.5 Hz, 1H), 5.35 (s, 2H) ppm; ^{13}C NMR (101 MHz, CDCl_3) δ 165.23 (d, $J_{\text{C-F}}$ = 250.3 Hz), 158.85 (d, $J_{\text{C-F}}$ = 12.4 Hz), 148.54, 148.51, 131.24, 129.88, 129.52, 127.69, 127.67, 127.49 (d, $J_{\text{C-F}}$ = 10.4 Hz), 126.50, 124.52, 119.86 (d, $J_{\text{C-F}}$ = 2.7 Hz), 110.30 (d, $J_{\text{C-F}}$ = 22.2 Hz), 104.87 (d, $J_{\text{C-F}}$ = 24.7 Hz), 68.96 ppm; MS (ESI) m/z : calcd for $\text{C}_{16}\text{H}_{11}\text{FNO}^+$ $[\text{M} + \text{H}]^+$: 252.0819, found: 252.0819.

3-Chloro-6H-chromeno[4,3-b]quinoline (7q). Following the general procedure, the reaction of **6q** (142.5 mg, 0.5 mmol) gave, after FC on silica gel (eluent: EtOAc/petroleum ether = 1:70), compound **7q** as a white floppy solid (105 mg, yield: 79%). Mp: 138–139 °C; ^1H NMR (400 MHz, CDCl_3) δ 8.40 (d, J 8.4 Hz, 1H), 8.10 (d, J 8.5 Hz,

1H), 7.83 (s, 1H), 7.75 (d, *J* 8.2 Hz, 1H), 7.71–7.67 (m, 1H), 7.51–7.47 (m, 1H), 7.12 (dd, *J* 8.4, 2.0 Hz, 1H), 7.02 (d, *J* 2.0 Hz, 1H), 5.34 (s, 2H) ppm; ¹³C NMR (100 MHz, CDCl₃) δ 158.00, 148.52, 148.31, 137.34, 131.27, 129.94, 129.59, 127.82, 127.68, 126.83, 126.69, 124.79, 123.16, 122.05, 117.85, 68.83 ppm; HRMS (ESI) *m/z*: calcd for C₁₆H₁₁ClNO⁺ [M + H]⁺: 268.0524, found: 268.0524.

3-Bromo-6H-chromeno[4,3-*b*]quinoline (7r). Following the general procedure, the reaction of **6r** (164.5 mg, 0.5 mmol) gave, after FC on silica gel (eluent: EtOAc/petroleum ether = 1:70), compound **7r** as a white floppy solid (127 mg, yield: 82%). Mp: 150–152 °C; ¹H NMR (400 MHz, CDCl₃) δ 8.30 (d, *J* 8.4 Hz, 1H), 8.07 (d, *J* 8.4 Hz, 1H), 7.76 (s, 1H), 7.73–7.64 (m, 2H), 7.47 (t, *J* 7.3 Hz, 1H), 7.25 (dd, *J* 8.3, 2.0 Hz, 1H), 7.16 (d, *J* 1.9 Hz, 1H), 5.29 (s, 2H) ppm; ¹³C NMR (100 MHz, CDCl₃) δ 157.96, 148.46, 148.26, 131.21, 129.90, 129.55, 127.79, 127.65, 126.96, 126.67, 125.96, 125.41, 124.75, 122.41, 120.75, 68.76 ppm; HRMS (ESI) *m/z*: calcd for C₁₆H₁₁BrNO⁺ [M + H]⁺: 312.0019, found: 312.0018.

3-(Trifluoromethyl)-6H-chromeno[4,3-*b*]quinoline (7s). Following the general procedure, the reaction of **6s** (159.5 mg, 0.5 mmol) gave, after FC on silica gel (eluent: EtOAc/petroleum ether = 1:80), compound **7s** as a pale-yellow solid (75 mg, yield: 50%). Mp: 138–140 °C; IR (film) *ν*_{max}: 3077, 2924, 1582, 1496, 1450, 1327, 1164, 1118, 880, 763 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 8.55 (d, *J* 8.1 Hz, 1H), 8.11 (d, *J* 8.5 Hz, 1H), 7.81 (s, 1H), 7.76–7.67 (m, 2H), 7.52–7.48 (m, 1H), 7.36 (dd, *J* 8.2, 1.7 Hz, 1H), 7.25 (s, 1H), 5.35 (s, 2H) ppm; ¹³C NMR (100 MHz, CDCl₃) δ 157.34, 148.51, 147.65, 133.44 (q, *J* C-F = 33.0 Hz), 131.41, 130.05, 129.78, 128.04, 127.69, 127.09, 126.37, 125.27, 125.06, 123.92 (q, *J* C-F = 271.0 Hz), 119.11 (q, *J* C-F = 3.8 Hz), 114.94 (q, *J* C-F = 4.0 Hz), 68.67 ppm; HRMS (ESI) *m/z*: calcd for C₁₇H₁₁F₃NO⁺ [M + H]⁺: 302.0787, found: 302.0787.

3-Methyl-6H-chromeno[4,3-*b*]quinoline (7t). Following the general procedure, the reaction of **6t** (132.5 mg, 0.5 mmol) gave, after FC on silica gel (eluent: EtOAc/petroleum ether = 1:80), compound **7t** as a pale-yellow floppy solid (107 mg, yield: 87%). Mp: 101–103 °C; ¹H NMR (400 MHz, CDCl₃) δ 8.31 (d, *J* 7.9 Hz, 1H), 8.05 (d, *J* 8.7 Hz, 1H), 7.62–7.56 (m, 3H), 7.38–7.33 (m, 1H), 6.92 (d, *J* 7.9 Hz, 1H), 6.77 (s, 1H), 5.17 (s, 2H), 2.32 (s, 3H) ppm; ¹³C NMR (101 MHz, CDCl₃) δ 157.42, 149.20, 148.36, 142.66, 130.77, 129.45, 129.33, 127.50, 127.48, 126.01, 125.47, 125.07, 123.66, 120.65, 117.71, 68.42, 21.73 ppm; HRMS (ESI) *m/z*: calcd for C₁₇H₁₄NO⁺ [M + H]⁺: 248.1070, found: 248.1069.

3-Methoxy-6H-chromeno[4,3-*b*]quinoline (7u). Following the general procedure, the reaction of **6u** (140.5 mg, 0.5 mmol) gave, after FC on silica gel (eluent: EtOAc/petroleum ether = 1:70), compound **7u** as a pale-yellow solid (125 mg, yield: 95%). Mp: 119–122 °C; ¹H NMR (400 MHz, CDCl₃) δ 8.36 (d, *J* 8.7 Hz, 1H), 8.05 (d, *J* 8.4 Hz, 1H), 7.68 (s, 1H), 7.66–7.59 (m, 2H), 7.42–7.37 (m, 1H), 6.70 (dd, *J* 8.7, 2.5 Hz, 1H), 6.50 (d, *J* 2.5 Hz, 1H), 5.24 (s, 2H), 3.80 (s, 3H) ppm; ¹³C NMR (100 MHz, CDCl₃) δ 163.06, 158.95, 149.24, 148.49, 130.81, 129.54, 129.23, 127.56, 127.34, 126.92, 125.86, 124.54, 116.41, 109.81, 101.87, 68.78, 55.59 ppm; HRMS (ESI) *m/z*: calcd for C₁₇H₁₄NO₂⁺ [M + H]⁺: 264.1019, found: 264.1019.

Acknowledgements

The authors are grateful to the NSF of China (21931010) for financial support.

Supplementary Material

Experimental procedures for the synthesis of the amides, and copies of ^1H and ^{13}C NMR spectra of compounds **6d-6h**, **6p-6u**, **7a-7g**, **7j-7u**, and X-ray crystallographic data of compounds **7g**, **7u** (CCDC 2335879, 2335878) are presented in the Supplementary Material associated with this paper.

References

1. Sultana, R.; Tippanna, R. R. *Lett. Org. Chem.* **2021**, *18*, 265-272.
<https://doi.org/10.2174/1570178617666200122095829>
2. Kumar, A. S.; Kumar, R. A.; Satyanarayana, V.; Reddy, E. P.; Reddy, B. J. M.; Kumar, D. N.; Khurana, A.; Chandraiah, G.; Yadav, J. S. *Nat. Prod. Commun.* **2017**, *12*, 1934578X1701200732.
<https://doi.org/10.1177/1934578x1701200732>
3. Vu, A. T.; Campbell, A. N.; Harris, H. A.; Unwalla, R. J.; Manas, E. S.; Mewshaw, R. E. *Bioorg. Med. Chem. Lett.* **2007**, *17*, 4053-4056.
<https://doi.org/10.1016/j.bmcl.2007.04.068>
4. Chaudhary, C. L.; Ko, S.; Lee, C.; Kim, Y.; Jung, C.; Hyun, S.; Kwon, Y.; Kang, J.-S.; Jung, J.-K.; Lee, H. *Pharmaceuticals* **2022**, *15*, 399.
<https://doi.org/10.3390/ph15040399>
5. Niu, P.; Tian, R.; Liu, Z.; Ran, J.; Liu, J.; Li, Y.; Ren, X.; Liu, X.; Wei, L. *Dyes Pigm.* **2022**, *205*, 110485.
<https://doi.org/10.1016/j.dyepig.2022.110485>
6. Liu, X.; Su, Y.; Tian, H.; Yang, L.; Zhang, H.; Song, X.; Foley, J. W. *Anal. Chem.* **2017**, *89*, 7038-7045.
<https://doi.org/10.1021/acs.analchem.7b00754>
7. Huang, W.; Lin, W.; Guan, X. *Tetrahedron Lett.* **2014**, *55*, 116-119.
<https://doi.org/10.1016/j.tetlet.2013.10.130>
8. Kand, D.; Kalle, A. M.; Varma, S. J.; Talukdar, P. *Chem. Commun.* **2012**, *48*, 2722-2724.
<https://doi.org/10.1039/C2CC16593G>
9. Rajput, D.; Tsering, D.; Karuppasamy, M.; Kapoor, K. K.; Nagarajan, S.; Maheswari, C. U.; Bhuvanesh, N.; Sridharan, V. *J. Org. Chem.* **2023**, *88*, 8643-8657.
<https://doi.org/10.1021/acs.joc.3c00552>
10. Kishore, D. R.; Mounika, K.; Goel, K.; Naveen, J.; Satyanarayana, G. *Synthesis* **2023**, *55*, 808-820.
<https://doi.org/10.1055/s-0041-1738429>
11. Penaranda Gomez, A.; Puerto Galvis, C. E.; Macias, M. A.; Ochoa-Puentes, C.; Kouznetsov, V. V. *Synthesis* **2022**, *54*, 1857-1869.
<https://doi.org/10.1055/a-1638-5030>
12. Weng, Y.; Chen, H.; Li, N.; Yang, L.; Ackermann, L. *Adv. Synth. Catal.* **2021**, *363*, 2773-2777.
<https://doi.org/10.1002/adsc.202100146>
13. Dong, W.; Yuan, Y.; Gao, X.; Hu, B.; Xie, X.; Zhang, Z. *ChemCatChem* **2018**, *10*, 2878-2886.
<https://doi.org/10.1002/cctc.201800192>
14. Ibrahim, Y. A.; Moustafa, A. H. *J. Chem. Res. Synop.* **1999**, 254-255.
<https://doi.org/10.1039/A809649J>
15. Rougeot, E.; Moskowitz, H.; Miocque, M. *Tetrahedron Lett.* **1983**, *24*, 2379-2382.
[https://doi.org/10.1016/S0040-4039\(00\)81930-9](https://doi.org/10.1016/S0040-4039(00)81930-9)
16. Yu, X.; Wang, J.; Xu, Z.; Yamamoto, Y.; Bao, M. *Org. Lett.* **2016**, *18*, 2491-2494.
<https://doi.org/10.1021/acs.orglett.6b01065>
17. Aradi, K.; Bombicz, P.; Novak, Z. *J. Org. Chem.* **2016**, *81*, 920-931.

- <https://doi.org/10.1021/acs.joc.5b02490>
18. Huang, H.; Kang, J. Y. *Synthesis* **2022**, 54, 1157-1202.
<https://doi.org/10.1055/a-1679-8205>
19. Falmagne, J.-B.; Escudero, J.; Taleb-Sahraoui, S.; Ghosez, L. *Angew. Chem. Int. Ed.* **1981**, 20, 879-880.
<https://doi.org/10.1002/anie.198108791>
20. Ghosez, L. *Tetrahedron* **2019**, 75, 130345.
<https://doi.org/10.1016/j.tet.2019.05.024>
21. Pace, V.; Holzer, W.; Olofsson, B. *Adv. Synth. Catal.* **2014**, 356, 3697-3736.
<https://doi.org/10.1002/adsc.201400630>
22. Kaiser, D.; Bauer, A.; Lemmerer, M.; Maulide, N. *Chem. Soc. Rev.* **2018**, 47, 7899-7925.
<https://doi.org/10.1039/C8CS00335A>
23. Matheau-Raven, D.; Gabriel, P.; Leitch, J. A.; Almeahmadi, Y. A.; Yamazaki, K.; Dixon, D. J. *ACS Catal.* **2020**, 10, 8880-8897.
<https://doi.org/10.1021/acscatal.0c02377>
24. Czerwiński, P. J.; Furman, B. *Front. Chem.* **2021**, 9, 655849.
<https://doi.org/10.3389/fchem.2021.655849>
25. Chen, H.; Chen, D.-H.; Huang, P.-Q. *Cell Rep. Phys. Sci.* **2023**, 4, 101574.
<https://doi.org/10.1016/j.xcrp.2023.101574>
26. He, Q.; Li, J.; Yu, S.-J.; Wu, D.-P.; Ye, J.-L.; Huang, P.-Q. *Acta Chim. Sinica* **2023**, 81, 1265-1270.
<https://doi.org/10.6023/A23050226>
27. Shao, D.-Y.; Huang, P.-Q. *Arkivoc* **2023**, ii, 202312048.
<https://doi.org/10.24820/ark.5550190.p012.048>
28. Han, F.; Lu, G.-S.; Wu, D.-P.; Huang, P.-Q. *Sci. China Chem.* **2023**, 66, 1094-1100.
<https://doi.org/10.1007/s11426-022-1501-y>
29. Xu, F.-F.; Chen, J.-Q.; Shao, D.-Y.; Huang, P.-Q. *Nat. Commun.* **2023**, 14, 6251.
<https://doi.org/10.1038/s41467-023-41846-x>
30. Su, X.-Y.; Huang, P.-Q. *Synthesis* **2023**, 55, 877-891.
<https://doi.org/10.1055/a-1957-4343>
31. Chen, T.-T.; Geng, H.; Weng, Z.-Y.; Huang, P.-Q. *ACS Earth Space Chem.* **2023**, 7, 243-251.
<https://doi.org/10.1021/acsearthspacechem.2c00315>
32. Yu, S.-J.; Li, J.; Ye, J.-L.; Huang, P.-Q. *Org. Chem. Front.* **2023**, 10, 1994-2001.
<https://doi.org/10.1039/D3QO00092C>
33. Bechara, W. S.; Pelletier, G.; Charette, A. B. *Nat. Chem.* **2012**, 4, 228.
<https://doi.org/10.1038/nchem.1268>
34. Movassaghi, M.; Hill, M. D. *J. Am. Chem. Soc.* **2006**, 128, 14254-14255.
<https://doi.org/10.1021/ja066405m>
35. Sugiyama, Y.; Yamada, K.; Kaneko, D.; Kusagawa, Y.; Okamura, T.; Sato, T. *Angew. Chem. Int. Ed.* **2024**, 63, e202317290.
<https://doi.org/10.1002/anie.202317290>
36. Liyama, S; Mizutani, K and Sato, T. *Chem. Lett.* **2023**, 52, 682-684.
<https://doi.org/10.1246/cl.230245>
37. Iwamoto, S.; Tokuyama, A.; Hiraoka, S.; Takei, K.; Matsuzaka, K.; Matsumoto, T.; Chida, N.; Sato, T. *Bull. Chem. Soc. Jpn.* **2023**, 96, 529-537.
<https://doi.org/10.1246/bcsj.20230088>

38. Soda, Y.; Sugiyama, Y.; Sato, S.; Shibuya, K.; Saegusa, J.; Matagawa, T.; Kawano, S.; Yoritake, M.; Fukaya, K.; Urabe, D.; Oishi, T.; Mori, K.; Simizu, S.; Chida, N.; Sato, T. *Synthesis* **2023**, 55, 617-636.
<https://doi.org/10.1055/a-1941-8680>
39. Feng, M.; Fernandes, A. J.; Meyrelles, R.; Maulide, N. *Chem* **2023**, 9, 1538-1548.
<https://doi.org/10.1016/j.chempr.2023.03.002>
40. Heindl, S.; Riomet, M.; Matyasovsky, J.; Lemmerer, M.; Malzer, N.; Maulide, N. *Angew. Chem. Int. Ed.* **2021**, 60, 19123-19127.
<https://doi.org/10.1002/anie.202104023>
41. Miller, A. A. M.; Biallas, P.; Shennan, B. D. A.; Dixon, D. J. *Angew. Chem. Int. Ed.* **2024**, 63, e202314308.
<https://doi.org/10.1002/anie.202314308>
42. Biallas, P.; Yamazaki, K.; Dixon, D. J. *Org. Lett.* **2022**, 24, 2002-2007.
<https://doi.org/10.1021/acs.orglett.2c00438>
43. Zhao, C.; Ge, Z.-L.; Hu, J.-H.; Tian, H.-J.; Wang, X.-M. *Cell Rep. Phys. Sci.* **2023**, 4, 101474.
<https://doi.org/10.1016/j.xcrp.2023.101474>
44. Zhao, F.; Jiang, F.; Wang, X.-M. *Sci. China Chem.* **2022**, 65, 2231-2237.
<https://doi.org/10.1007/s11426-022-1331-y>
45. Zhou, H.-B.; Liu, G.-S.; Yao, Z.-J. *Org. Lett.* **2007**, 9, 2003-2006.
<https://doi.org/10.1021/ol0706307>
46. Ye, J.-L.; Zhu, Y.-N.; Geng, H.; Huang, P.-Q. *Sci. China Chem.* **2018**, 61, 687-694.
<https://doi.org/10.1007/s11426-017-9160-1>
47. Li, L.-H.; Niu, Z.-J.; Liang, Y.-M. *Chem.-Eur. J.* **2017**, 23, 15300-15304.
<https://doi.org/10.1002/chem.201703832>
48. Yu, S.-J.; Ye, J.-L.; Hong, Y.-C.; Huang, P.-Q. *J. Org. Chem.* **2021**, 86, 16926-16939.
<https://doi.org/10.1021/acs.joc.1c02098>
49. Lee, C. J.; Swain, M.; Kwon, O. *Org. Lett.* **2018**, 20, 5474-5477.
<https://doi.org/10.1021/acs.orglett.8b02398>
50. Nagarajan, R.; Ramesh, S.; Gaddam, V. *Synlett*, **2010**, 757-760.
<http://doi.org/10.1055/s-0029-1219364>

This paper is an open access article distributed under the terms of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>)