

Novel pyrrolocoumarin derivatives

Valery F. Traven, Vasily V. Suslov, and Evgeny N. Gordeev

*Department of Organic Chemistry, D. Mendeleev University of Chemical Technology of Russia,
125047 Moscow, Russia*

E-mail: traven@main-gw.muctr.edu.ru

(received 20 May 00; accepted 21 Sep 00; published on the web 29 Sep 00)

DOI: <http://dx.doi.org/10.3998/ark.5550190.0001.411>

Abstract

Ethyl esters of 2-[(4',7'-dimethyl-2'-oxo-2'H-1'-benzopyran-6'-yl)azo]-2-alkyl-3-oxobutanoic acids have been synthesized from diazotized 6-amino-4,7-dimethyl-2H-1-benzopyran-2-one and ethyl esters of 2-alkyl-3-oxobutanoic acids in the presence of sodium acetate. The esters have then been transformed into indoles (obviously via intermediate hydrazones) by the Fischer reaction in the presence of acids. This is a way to novel class of indole derivatives - ethyl esters of 3,7-dihydro-7-oxopyrano[3,2-*e*]indol-2-carboxylic acids with different substituents. All the new compounds have been characterised by ¹H-NMR and mass spectra, and elemental analyses.

Keywords: Pyrrolocoumarins, diazotization Fisher indole synthesis

Introduction

2H-1-benzopyran-2-one (coumarin) derivatives belong to one of the most widespread classes of natural compounds (1). Some of them, for example, furocoumarin derivatives - psoralens and angelicins have been used as medicines (2). Pyrrolocoumarins showed also photobiological activity and antiproliferative effect (3).


Some of the pyrrolocoumarins have previously been synthesized by the reaction between aminocoumarins and benzoin (4). Cyclization of the corresponding hydrazones by the Fischer reaction has also been used (3,5). However, these procedures have only been applied for synthesis of alkyl- and aryl-substituted pyrrolocoumarins.

In this article we report a new way to ethoxycarbonyl substituted derivatives of pyrrolocoumarins. This procedure is based on the Japp-Klingemann synthesis of hydrazones (6, 7) and provides a convenient way to pyrrolocoumarins with carboethoxy function in pyrrole ring. A number of derivatives of 3,7-dihydro-4,9-dimethyl-7-oxopyrano[3,2-*e*]indol-2-carboxylic acid ethyl esters has been synthesized.

Results and Discussion


4,7-Dimethyl-6-aminocoumarin **1** has been prepared by reduction of 4,7-dimethyl-6-nitrocoumarin (**8**) with powder of cast iron in a mixture of dioxan-water and pH = 4–5. This procedure is more convenient and provides a better yield of **1** than the syntheses reported previously (**4**).

Aminocoumarin **1** has been diazotized by a standard procedure. The formed diazo-compound has then been coupled with α -substituted acetoacetic esters in the presence of sodium acetate (scheme 1).



Scheme 1


Compounds **2a,b** have been isolated and purified as solid products. However, product **2c** has been isolated as an oil and transformed then without purification into solid hydrazone **3** by hydrochloric acid treatment at room temperature (scheme 2).




Scheme 2

Solvent, catalyst and temperature of reaction are of a great importance for indole derivatives synthesis of by the Fischer reaction (7, 9).

Heating of hydrazone **3c** at 80 °C in HCl-ethanol gave pyrrolocoumarin **4c**. The ester **2b** has smoothly been transformed to pyrrolocoumarin **4b** in the same conditions without isolation of the intermediate hydrazone. The ester **2a** has also been transformed directly to pyrrolocoumarin **4a**, but solution of p-toluenesulphonic acid monohydrate in glacial acetic acid was used as a cyclization reagent (scheme 3).



Scheme 3

**¹H-NMR spectra**


All ¹H-NMR spectra were taken on a Bruker WR-200SV spectrometer at 200 Mhz in CDCl₃ solution using tetramethylsilane as internal standard. Chemical shifts are given in ppm.

The chemical shifts for the protons of Me-group at C-4 of compounds **2a-c** appeared at 2.40–2.44 ppm. The chemical shifts for the protons of Me-group at C-9 of compounds **4a-b** were found at 2.69 ppm for **4a** and at 2.75 ppm for **4b**. In the case of **4c**, however, the chemical shifts for the protons of Me-group at C-9 was found at a relatively higher field at 1.00 ppm. This shift towards a higher field could be caused by an anisotropic shielding effect of phenyl group located at C-1 of **4c**.

Mass spectra

All mass spectra were scanned on a SSQ-710 (Finnigan MAT) spectrometer at the energy of ionising electrons equal to 70 eV.

Molecular ions of compounds **3c** and **4a-c** have high intensity peaks (near 75–85 %). Azo compounds **2a** and **2b** have not molecular ions peaks. Their molecular ions decompose under electron impact in several directions (scheme 4).



Scheme 4

The decomposition of molecular ions of compounds **4a-c** goes either with loss of CH₃CO, then CO or with loss of COOEt.

Experimental Section

6-Amino-4,7-dimethyl-2H-1-benzopyran-2-one (6-amino-4,7-dimethylcoumarin) 1. A mixture of 35 g powder of cast iron, 100 mL of water and 10 mL of glacial acetic acid was refluxed for 15 minutes. A boiling solution of 6-nitro-4,7-dimethylcoumarin (42 g, 0.2 mole) in 500 mL of 1,4-dioxane was then added into the mixture. It has then been refluxed for 7 h. The resulting reaction mixture was neutralized with 46 g solid of sodium carbonate and filtered from a solid precipitate. The precipitate was washed with boiling 1,4-dioxane. The filtrate was then poured into 300 mL of cold water. A solid was filtered off, washed with water, dried and recrystallized from 1,4-dioxane. Yield 31.0 g (~82%), mp=203–205 (lit. (4) 203–205 (1,4-dioxane)).

Synthesis of Azo Compounds 2a-c (General Procedure)

A solution of 6-amino-4,7-dimethylcoumarin **1** (9.45 g, 0.05 mole) in a mixture of 25 mL of hydrochloric acid and 50 mL of glacial acetic acid was diazotized at 0 °C with sodium nitrite solution (4 g, (0.055 mole) dissolved in 10 mL of water). After keeping at this temperature for 3 h the resulting solution of diazonium salt was filtered and added into mixture of 75 mL of glacial acetic acid, 0.051 mole of 2-alkyl-3-oxobutanoic acid ethyl ester and 50 g, (0.37 mole) of sodium acetate (AcONa 3H₂O) at 0 to 5 °C and pH =5.5. The mixture was left for 10 h. After that an equal volume of water was added into the mixture. The crude product was collected, washed with ethanol, then water and recrystallized from ethanol (treatment of compound **2c** see below).

2-[(4',7'-Dimethyl-2'-oxo-2'H-1'-benzopyran-6'-yl)azo]-2-methyl-3-oxobutanoic acid ethyl ester 2a. Yield 15.0 g. (~87%), mp=84–85 °C. ¹H-NMR (CDCl₃, J/Hz) δ 1.29 (t, 3H, Me, J_{Me,-CH₂-}=7.5, COOEt); 1.71 (s, 3H, 2-Me); 2.34 (s, 3H, 7'-Me); 2.44 (d, 3H, 4'-Me, J_{4'-Me,3'}=1.6); 2.64 (s, 3H, MeCO); 4.29 (q, 2H, J-CH₂-,Me=7.5, COOEt); 6.28 (d, 1H, 3'-H, J_{3',4'-Me}=1.6); 7.25 (s, 1H, 8'-H); 7.64 (s, 1H, 5'-H). MS: m/z(%): 302 (M⁺ -CH₂=C=O, 9), 271 (M⁺ -COOEt, 25), 201 (Coum-N=N⁺, 65), 173 (Coum⁺, 100). Anal. calcd. for C₁₈H₂₀N₂O₅: C 62.78; H 5.85; N 8.13. Found C 63.00; H 5.81; N 8.15.

2-[(4',7'-Dimethyl-2'-oxo-2'H-1'-benzopyran-6'-yl)azo]-2-ethyl-3-oxobutanoic acid ethyl ester 2b. Yield 15.4 g. (~86%), mp=88–90 °C. ¹H-NMR (CDCl₃, J/Hz) δ 1.04 (t, 3H, Me, J_{Me,-CH₂-}=7.5, 2-Et); 1.29 (t, 3H, Me, J_{Me,-CH₂-}=7.5, COOEt); 2.29 (q, 2H, J-CH₂-,Me =7.5, 2-Et); 2.31 (s, 3H, 7'-Me); 2.44 (d, 3H, 4'-Me, J_{4'-Me,3'}=1.8); 2.64 (s, 3H, MeCO); 4.29 (q, 2H, -CH₂-, J-CH₂-,Me=7.5, COOEt); 6.28 (d, 1H, 3'-H, J_{3',4'-Me}=1.8); 7.26 (s, 1H, 8'-H); 7.63 (s, 1H, 5'-H). MS: m/z(%): 330 (M⁺ -CH₂=CH₂, 7), 316 (M⁺ -CH₂=C=O, 6), 288 (M⁺ -CH₂=CH₂ -CH₂=C=O, 12), 285 (M⁺ -COOEt, 5), 201 (Coum-N=N⁺, 76), 173 (Coum⁺, 100). Anal. calcd. for C₁₉H₂₂N₂O₅: C 63.68; H 6.19; N 7.82. Found: C 63.83; H 6.31; N 7.80.

a-[(E)-2-(4',7'-Dimethyl-2'-oxo-2H-1'-benzopyran-6'-yl)hydrazono]-benzenepropanoic acid ethyl ester 3c. Azo compound **2c** as an oil was treated with cold diluted hydrochloric acid and product was obtained as crystals. The hydrazone **3c** was filtered off, washed with ethanol, then water. The product **3c** was recrystallized from ethanol. Yield 13g. (~70%), mp=160–162 °C. ¹H-NMR (CDCl₃, J/Hz) δ 1.29 (t, 3H, Me, J_{Me,-CH₂-}=7.3, COOEt); 2.35 (d, 3H, 7'-Me, J_{7'-Me,8'}=0.3); 2.40 (d, 3H, 4'-Me, J_{4'-Me,3'}=1.8); 3.89 (s, 2H, -CH₂-Ph); 4.27 (q, 2H, -CH₂-, J-CH₂-,Me=7.3, COOEt); 6.23 (d, 1H, 3'-H, J_{3',4'-Me}=1.8); 7.10 (d, 1H, 8'-H, J_{8',7'-Me}=0.3); 7.30 (m, 5H, Ph-); 7.59 (s, 1H, 5'-H); 12.24 (s, 1H, NH). MS: m/z(%): 378 (M⁺, 75), 188 (Coum-N⁺, 100), 173 (Coum⁺, 16). Anal. calcd. for C₂₂H₂₂N₂O₄: C 69.83; H 5.86; N 7.40. Found: C 69.62; H 5.92; N 7.34.

3,7-Dihydro-4,9-dimethyl-7-oxopyrano[3,2-e]indole-2-carboxylic acid ethyl ester 4a. A mixture of azo compound **2a** 1g (0.0035 mole), 10 mL of glacial acetic acid and 1g (0.0052 mole) of TosOH·H₂O has been refluxed for 12 h. After cooling the reaction mixture was poured into 100 g. of crushed ice. The product was filtered off, washed with water and recrystallized

from 1,4-dioxan (activated charcoal). Yield 0.3 g. (~36%), mp >300° C. ¹H-NMR (CDCl₃, J/Hz) δ 1.45 (t, 3H, Me, J_{Me,-CH₂-}=7.5, COOEt); 2.61 (d, 3H, 4-Me, J_{4-Me,5} =1.0); 2.75 (d, 3H, 9-Me, J_{9-Me,8}=1.3); 4.46 (q, 2H, -CH₂-, J-CH₂-,Me= 7.5, COOEt); 6.28 (d,1H, 8-H, J_{8,9-Me}=1.3); 7.17 (d,1H, 5-H, J_{5,4-Me}=1.0); 7.56 (d, 1H, 1-H, J_{1H,NH} = 2.14); 9.08 (c,1H, NH). MS: m/z(%): 285 (M⁺, 81), 257 (M⁺-CH₂=CH₂, 11), 240(M⁺-Et-O, 14), 212(M⁺-COOEt, 19). Anal. calcd. for C₁₆H₁₅NO₄: C 67.36; H 5.30; N 4.91. Found: C 67.29; H 5.25; N 4.89.

3,7-Dihydro-1,4,9-trimethyl-7-oxopyrano[3,2-*e*]indole-2-carboxylic acid ethyl ester 4b. Azo compound **2b** (12.4 g, 0.035 mol) was added into a mixture of 30 mL ethanol and 15 mL SOCl₂. After refluxing for 12 h the reaction mixture was cooled. The precipitate was filtered off, washed with ethanol, then water and recrystallized from ethanol. Yield 8.8 g. (~85.0%), mp=178–180 °C. ¹H-NMR (CDCl₃, J/Hz) δ 1.45 (t, 3H, Me, J_{Me,-CH₂-}=7.5, COOEt); 2.57 (d, 3H, 4-Me, J_{4-Me,5} =0.8); 2.69 (d, 3H, 9-Me, J_{9-Me,8}=1.1); 2.78 (s, 3H, 1-Me); 4.46 (q, 2H, -CH₂-, J-CH₂-,Me =7.5, COOEt); 6.25 (d, 1H, 8-H, J_{8,9-Me}=1.1); 7.12 (d, 1H, 5-H, J_{5,4-Me}=0.8); 9.0 (s,1H, NH). MS: m/z(%): 299 (M⁺, 80), 254(M⁺-Et-O, 67), 226(M⁺-COOEt, 21). Anal. calcd. for C₁₇H₁₇NO₄: C 68.22; H 5.72; N 4.68. Found: C 67.98; H 5.68; N 4.73.

3,7-dihydro-4,9-dimethyl-7-oxo-1-phenylpyrano[3,2-*e*]indole-2-carboxylic acid ethyl ester 4c. Hydrazone **3c** (10.5 g,0.028 mol) was added to a mixture of 40 mL ethanol and 15 mL SOCl₂. After refluxing for 12 h the reaction mixture was cooled. The precipitate was filtered off, washed with ethanol, then water. The compound **4c** was recrystallized from ethanol. Yield 5 g. (~49.7%), mp=223–225 °C. ¹H-NMR (CDCl₃, J/Hz) δ 1.10 (t, 3H, Me, J_{Me,-CH₂-}=7.5, COOEt); 1.56 (d, 3H, 9-Me, J_{9-Me,8} =1.0); 2.61 (d, 3H, 4-Me, J_{4-Me,5} =0.8); 4.20 (q, 2H, -CH₂-, J-CH₂-,Me =7.5, COOEt); 6.02 (d, 1H, 8-H, J_{8,9-Me}=1.0); 7.20 (d, 1H, 5-H, J_{5,4-Me}=0.8); 7.37 (m, 5H, Ph); 9.28 (s,1H, NH). MS: m/z(%): 361 (M⁺, 85), 316(M⁺-Et-O, 27), 288(M⁺-COOEt, 4). Anal. calcd. for C₂₂H₁₉NO₄: C 73.12; H 5.30; N 3.88. Found: C 73.11; H 5.28; N 3.77.

References

1. Dean F.M. *Naturally Occurring Oxygen Ring Compounds*, Butterworth: London, 1963.
2. Dall'Acqua F.; Vedaldi D.; Caffieri S.; Guiotto A.; Rodighiero P.; Baccichetti F.; Carlassare F.; Bordin F. *J. Med. Chem.* **1981**, *24*, 178.
3. Guiotto A.; Chilin A.; Manzini P.; Dall'Acqua F.; Bordin F.; Rodighiero P. *Farmaco* **1995**, *50*, 479.
4. Meyer H. *Acta Chem. Scand., Ser. B*, **1975**, *29*, 133.
5. Khan M.; Morley B. *J. Heterocyclic Chem.* **1979**, *16*, 997.
6. Japp F.R.; Klingemann F.; *Ber.*, **1887**, *20*, 2942, 3284, 3398.
7. Sundberg R.J.; *Indoles, Academic Press*, **1996**.
8. Clayton A. *J. Chem. Soc.* **1910**, *97*, 1398.
9. Robinson R. *The Fischer Indole Synthesis*, Wiley, Chichester, **1982**.

