

Microwave-assisted asymmetric Diels-Alder reaction using chiral auxiliaries derived from biomass

Ariel M. Sarotti, Rolando A. Spanevello, and Alejandra G. Suárez*

Instituto de Química Rosario, Facultad de Ciencias Bioquímicas y Farmacéuticas – Universidad Nacional de Rosario-CONICET, Suipacha 531, S2002LRK Rosario, Argentina

E-mail: suarez@iquir-conicet.gov.ar

Dedicated to Professors Manuel González Sierra, Rita M. H. de Rossi, Julio Podestá and Oscar Giordano for their outstanding contribution to organic chemistry in Argentina

DOI: <http://dx.doi.org/10.3998/ark.5550190.0012.706>

Abstract

Microwave-assisted asymmetric Diels-Alder reactions using levoglucosenone-derived chiral auxiliaries have been investigated. The cycloaddition reaction between the corresponding chiral acrylate and cyclopentadiene showed important rate enhancements under microwave irradiation compared to conventional heating conditions, whereas the yields and diastereoselectivities were not significantly affected.


Keywords: Microwaves, Diels-Alder, asymmetric synthesis, levoglucosenone, acrylates

Introduction

The Diels-Alder reaction has proven to be an exceptionally powerful method for carbon-carbon bond forming reactions in the synthesis of natural products and novel structures of both fundamental and applied interest. Indeed, in this process the simultaneous creation of two new C-C bonds is coupled with the possibility of regio- and stereo-control of the newly formed six-membered ring.¹

As part of our program on the pyrolytic conversion of biomass into useful chemicals, we have been concerned with the synthetic application of levoglucosenone (1,6-anhydro-3,4-dideoxy- β -D-glycero-hex-3-enopyranos-2-ulose) **1** as a chiral building block for the development of new tools for asymmetric synthesis. Conventional pyrolysis of cellulose-containing materials such as waste paper is typically used to generate **1**,^{2,3} but microwave (MW) irradiation of microcrystalline cellulose is also effective.⁴ Recently, we have reported the design and synthesis of alcohols **2-6** derived from levoglucosenone (Scheme 1) and their use as chiral inductors in asymmetric Diels-Alder reactions of the corresponding acrylate with

cyclopentadiene yielding the products with up to 98% diastereomeric excess in the presence of Lewis acids.⁵



Scheme 1


In general non-catalyzed cycloaddition reactions do not afford good yields and selectivities of products. However, in our case the thermal Diels-Alder reactions produced the desired cycloadducts in very good yields (86-96%) with an inversion of the *endo* selectivity compared to the catalyzed reactions.⁵ The best diastereo- (*endo/exo*, ~72:28) and π -facial selectivities (up to 76% d.e.) were achieved at room temperature with the expected increase in reaction times (48-56 h). As the non-catalyzed Diels-Alder reactions needed long reaction times, we considered the use of a MW assisted procedure to improve the outcome of these reactions.⁶ This approach would save energy by shortening the reaction time and also by making the overall process more efficient and sustainable. We present herein the use of MW assisted Diels-Alder reaction of chiral acrylates derived from levoglucosenone with cyclopentadiene. The interest in this research is focused not only on the possibility to reduce reaction times, but also to evaluate the selectivity of the asymmetric process under MW irradiation which is of great synthetic and industrial significance.

Results and Discussion

In order to test the effectiveness of the chiral inductors **2-6** in asymmetric Diels-Alder reaction under MW irradiation we prepared the acrylates **7-11** according to the published procedures.⁵ The acylation step was easily performed in good yields by reaction of acryloyl chloride with the corresponding alcohol in the presence of triethylamine at 0 °C. Cycloaddition reactions of acrylates **7-11** with cyclopentadiene were carried out under MW irradiation, yielding the four isomeric products depicted in Scheme 2.

To make a systematic study of the different factors involved in these cycloaddition reactions, we first tested the role of the solvent as it can be a crucial factor in the outcome of the reaction. The selection of solvents was made considering a wide range of MW energy absorption capacity.

We used as a model reaction the cycloaddition of acrylate **7** with cyclopentadiene using a 300 W irradiation power. The use of hexane or THF did not produce the desired products since extensive polymerization of the cyclopentadiene took place under these conditions. However, after only 5 min of MW irradiation at 300 W, total conversion of the acrylate into the expected cycloadducts was observed using toluene (PhMe), ethanol (EtOH) or 1,1,1-trichloroethane (TCE) as solvents. Following these encouraging results, we next evaluated the cycloaddition reaction of several chiral acrylates using the same MW irradiation conditions (Table 1).



Scheme 2

Table 1. MW assisted Diels-Alder reaction between chiral acrylates **7-11** and cyclopentadiene. General conditions: 300 W, 5 min

Entry	Chiral Acrylate	Solvent	Yield (%)	Ratio <i>endo/exo</i>	Ratio <i>endo S/R</i>
1	7	PhMe	98	66 : 34	74 : 26
2	7	EtOH	97	72 : 28	69 : 31
3	7	TCE	93	66 : 34	71 : 29
4	8	PhMe	99	67 : 33	73 : 27
5	8	EtOH	79	71 : 29	73 : 27
6	8	TCE	94	71 : 29	72 : 28
7	9	PhMe	98	70 : 30	80 : 20
8	9	EtOH	94	73 : 27	76 : 24
9	9	TCE	90	66 : 34	76 : 24
10	10	PhMe	98	70 : 30	55 : 45
11	10	EtOH	98	75 : 25	51 : 49
12	10	TCE	98	69 : 31	53 : 47
13	11	PhMe	89	71 : 29	31 : 69
14	11	EtOH	91	70 : 30	32 : 68
15	11	TCE	90	64 : 36	35 : 65

The experimental data shown in Table 1 reveal that excellent yields were obtained in all cases. The stereochemical outcome of the cycloadditions showed little dependence on the solvent employed, although best *endo/exo* ratios were observed in EtOH whereas PhMe afforded the highest π -facial selectivities. Accordingly, further investigation was made using EtOH and PhMe as solvents.

Given the impressive reduction in reaction time when using MW irradiation (from 48-56 h to 5 min), a more comprehensive study directed toward finding optimal reaction conditions seemed to be appropriate. At this stage, there was a presumed relationship between the reaction temperature with the *endo/exo* and *endo R/S* ratios. We next investigated the effect of the irradiation time in the reaction of acrylate **7** and cyclopentadiene, with a power-controlled program at 300 W (Table 2).

Table 2. MW assisted Diels-Alder reactions between chiral acrylate **7** and cyclopentadiene. General conditions: PhMe, 300 W

Entry	Time (sec)	T (°C) ^a	Conv.	Yield (%)	Ratio <i>endo/exo</i>	Ratio <i>endo S/R</i>
1	75	68	44	42	70 : 30	80 : 20
2	150	103	87	81	67 : 33	75 : 25
3	225	128	95	91	67 : 33	74 : 26
4	300	140	100	98	66 : 34	74 : 26

^aMaximum temperature reached at the indicated time.

These experimental results indicate that there is a slight increase in selectivity at lower temperatures. For this reason, we considered diminishing the MW irradiation power, the reduction in reaction temperature obtained thereby, could increase the selectivity ratios. Table 3 shows the results obtained in a temperature-controlled program at 70 °C, which is the lowest temperature achieved employing PhMe or EtOH as solvents (MW irradiation power was approximately 5 W for PhMe and 1 W for EtOH). At this temperature, a longer irradiation time (25 min) was needed for total consumption of the starting material.

Table 3. MW assisted Diels-Alder reactions between chiral acrylates **7-11** and cyclopentadiene. General conditions: 70 °C, 25 min.

Entry	Chiral Acrylate	Solvent	Yield (%)	Ratio <i>endo/exo</i>	Ratio <i>endo S/R</i>
1	7	PhMe	92	69 : 31	78 : 22
2	7	EtOH	99	75 : 25	71 : 29
3	8	PhMe	97	71 : 29	80 : 20
4	8	EtOH	96	77 : 23	76 : 24
5	9	PhMe	91	69 : 31	83 : 17
6	9	EtOH	97	78 : 22	81 : 19
7	10	PhMe	92	73 : 27	57 : 43
8	10	EtOH	89	79 : 21	53 : 47
9	11	PhMe	86	78 : 22	30 : 70
10	11	EtOH	95	75 : 25	31 : 69

As shown in Table 3, the reduction in reaction temperature gave a slight improvement in selectivity whereas the overall yields were excellent in all cases. As a general trend, EtOH gave better *endo/exo* ratios and PhMe was the best in terms of facial selectivity.

From the experimental results collected in Tables 1 and 3, it can be observed that the power of the MW irradiation had an important effect on the reaction rate. Comparing the results obtained under thermal conditions *vs* MW irradiation it was evident that the latter produced an impressive increment in the reaction rates without significant changes in yields and selectivities. Finally, it is noteworthy to mention that none of the cycloaddition reactions reported in this study showed a retro Diels-Alder path, as was demonstrated to be the case in other MW accelerated Diels-Alder reactions.⁷

Conclusions

We have shown the impact of MW activation in asymmetric Diels-Alder reactions of chiral acrylates derived from levoglucosenone and cyclopentadiene. After extensive investigation, we found the optimal reaction conditions (PhMe or EtOH, 70 °C, 25 min) in terms of yield and selectivity. The desired products were isolated in excellent yields, and most importantly, the reaction time was dramatically shortened without loss of selectivity.

Experimental Section

General. The MW heating was performed in a CEM-Discover® System using a 10 mL pressure-rated reaction vials. HPLC analyses were performed with a chromatograph Varian ProStar

equipped with UV-V detector ProStar 320 at 270 nm, using a Beckman C-18, 25 cm column. Acetonitrile and water HPLC grade were used as eluents in a mixture 80:20, respectively. Flow rate was 1 mL/min. Nuclear magnetic resonance spectra were recorded on a Bruker Avance-300 DPX spectrometer with tetramethylsilane as internal standard and deuteriochloroform as solvent. The reactions were monitored by thin layer chromatography carried out on 0.25 mm E. Merck silica gel plates (60F₂₅₄) that were developed using UV light and *p*-anisaldehyde-sulfuric acid-acetic acid with subsequent heating. Flash column chromatography was performed using Merck silica gel 60H, by gradient elution created by mixtures of hexanes and increasing amounts of ethyl acetate. For complete characterization data of products **12a-d**, **13a-d** and **14a-d** see reference 5(a), for products **15a-d** and **16a-d**, see reference 5(c).

General procedure for the cycloaddition reaction of acrylates (7-11) and cyclopentadiene under MW irradiation

The corresponding acrylate (0.1 mmol) and cyclopentadiene (1 mmol) were placed in a 10 mL vial, dissolved in 4 mL of the suitable solvent, and heated for the time and MW irradiation power indicated in Tables 1-3. The solvent was then evaporated under reduced pressure and the residue was purified by flash chromatography to separate the excess of diene; the adducts were collected together to obtain an accurate yield. *Endo/exo* and *R/S* ratios were determined by HPLC, ¹H and ¹³C NMR.

Acknowledgements

This research was supported by grants from Universidad Nacional de Rosario, Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Agencia Nacional de Promoción Científica y Tecnológica (ANPCyT) and Secretaría de Estado de Ciencia, Tecnología e Innovación de la Provincia de Santa Fe.

References

1. Fringuelli, F.; Taticchi, A. *The Diels-Alder Reaction. Selected Practical Methods*; John Wiley & Sons: New York, 2002. (b) Fleming, I. *Pericyclic Reactions*; Oxford University Press Inc.: New York, 1999. (c) Nicolaou, K. C.; Snyder, S. A.; Montagnon, T.; Vassilikogiannakis, G. *Angew. Chem., Int. Ed.* **2002**, *41*, 1668.
2. Witczak, Z. J. *Levoglucosone and Levoglucosans: Chemistry and Applications*; ATL: Mount Prospect, 1994. (b) Witczak, Z. J.; Tatsuta, K.; *Carbohydrate Synthons in Natural Products Chemistry. Synthesis, Functionalization, and Applications*; ACS Symposium Series; American Chemical Society: Washington, D.C., 2003.
3. Swenton, J. S.; Freskos, J. N.; Dalidowicz, P.; Kerns, M. L. *J. Org. Chem.* **1996**, *61*, 459.

4. Sarotti, A. M.; Spanevello, R. A.; Suárez, A. G. *Green Chem.* **2007**, *9*, 1137.
5. Sarotti, A. M.; Spanevello, R. A.; Suárez, A. G. *Tetrahedron* **2009**, *65*, 3502. (b) Sarotti, A. M.; Fernández, I.; Spanevello, R. A.; Sierra, M. A.; Suárez, A. G. *Org. Lett.* **2008**, *10*, 3389. (c) Sarotti, A. M.; Spanevello, R. A.; Duhayon, C.; Tuchagues, J.-P.; Suárez, A. G. *Tetrahedron* **2007**, *63*, 241. (d) Sarotti, A. M.; Spanevello, R. A.; Suárez, A. G. *Org. Lett.* **2006**, *8*, 1487.
6. Hayes, B. L. *Microwave Synthesis. Chemistry at the Speed of Light*; CEM Publishing: Matthews, NC, 2002; Chapter 4. (b) Martínez Palau, R. *Química en Microondas*; CEM Publishing: Matthews, NC, 2006; Chapter 3. (c) Kappe, C. O. *Angew. Chem., Int. Ed.* **2004**, *43*, 6250. (d) Lidström, P.; Tierney, J.; Wathey, B.; Westman, J. *Tetrahedron* **2001**, *57*, 9225. (e) de la Hoz, A.; Díaz-Ortiz, A.; Moreno, A.; Langa, F. *Eur. J. Org. Chem.* **2000**, 3659.
7. Sarotti A. M.; Joullié, M. M.; Spanevello, R. A.; Suárez, A. G. *Org. Lett.* **2006**, *8*, 5561.