

Supplementary Material

Design and synthesis of *N*-benzimidazol-2-yl-*N'*-sulfonyl acetamidines

Nadezhda A. Rupakova,^a Vasiliy A. Bakulev,^a Uwe Knippschild,^b Balbina García-Reyes,^b Oleg S. Eltsov,^a Grigoriy P. Slesarev,^a Nikolai Beliaev,^a Pavel A. Slepukhin,^c Lydia Witt,^d Christian Peifer,^d and Tetyana V. Beryozkina^{a*}

^aUral Federal University named after the first President of Russia B. N. Yeltsin, 19 Mira st., Yekaterinburg 620002, Russia

^bKlinik für Allgemein- und Viszeralchirurgie Zentrum für Chirurgie Universitätsklinikum Ulm, 23 Albert-Einstein-Allee, 89081 Ulm, Germany

^cI. Ya. Postovsky Institute of Organic Synthesis, Ural Branch of Russian Academy of Sciences, 20 S. Kovalevskaya st., Yekaterinburg 620990, Russia


^dInstitute of Pharmacy, Christian-Albrechts-University of Kiel, Gutenbergstraße 76, D- 24118 Kiel, Germany


Email: tetber@mail.ru


Table of Contents


NMR spectra of 1d	3
NMR spectra of 1f	7
NMR spectra of 1h	9
NMR spectra of 2a	11
NMR spectra of 2b	13
NMR spectra of 2c	15
NMR spectra of 2d	18
NMR spectra of 2e	20
NMR spectra of 2f	22
NMR spectra of 2g	24
NMR spectra of 4a	26


NMR spectra of 4b	28
NMR spectra of 4c	30
NMR spectra of 4d	32
NMR spectra of 4e	34
NMR spectra of 4f	36
NMR spectra of 4g	38
NMR spectra of 4h	41
NMR spectra of 4i	44
NMR spectra of 4j	47
NMR spectra of 4k	49
NMR spectra of 4l	52
NMR spectra of 4m	55
NMR spectra of 4n	57
NMR spectra of 4o	59
Inhibitory effects of selected compounds on the kinase activity of CK1 isoforms	64
Effect on cell viability of selected synthesized compounds	65











































—7.94
—7.77
—7.75
—7.51
—7.49


—2.73
—2.51















































AT-USA, Inc
































































































Inhibitory effects of selected compounds on the kinase activity of CK1 isoforms

Selected compounds were initially screened for their activity against different CK1 isoforms against a concentration of 10 μ M ATP (Figure SI1). Most compounds had a weak effect on CK1 δ (Figure SI1C), with **2c** being the most potent inhibitor of this series. CK1 ϵ was inhibited only moderately by compound **4l** (Figure SI1D). All compounds did not inhibit CK1 α and CK1 γ 3 at all (Figure SI1A and SI1B). We next calculated IC₅₀ values for CK1 δ and CK1 ϵ regarding compounds **2c** and **4l** by assaying different concentrations in a range of 0.005 μ M to 30 μ M (Figure SI2). Hit compound **2c** showed an IC₅₀ value of approximately 7 μ M for CK1 δ and **4l** an IC₅₀ of 4.86 μ M for CK1 ϵ . Although these values are significantly weaker than IC₅₀ values of known highly potent CK1-specific inhibitors, our results suggest that our scaffolds might be improved for further development of potent CK1 isoform kinase inhibitors.




Figure SI1. Inhibitory effects of selected compounds on the kinase activity of CK1 isoforms.




Figure SI2. IC₅₀ curves of relevant compounds against CK1 δ and CK1 ϵ . Curve A: **2c**; curve B: **4l**.

Selected compounds were further evaluated with regard to their efficacy in cell culture by MTT viability assay using the glioblastoma cell line DK-MG (Figure SI3). The cells were exposed to 10 μ M of each compound with a periodic standard dosing scheme of 48 h. Compared to vehicle treatment the cell viability was not significantly diminished by the end of an exposure period of 7 days. The lack of biological activity of these compounds in this particular cellular system probably indicates suboptimal physicochemical properties.

Glioblastoma cell line DK-MG cultures were exposed to 10 μ M of selected synthesized compounds and analyzed for their effects on cell viability using MTT assay. Results are shown as normalized bar graphs using DMSO as a control for 100% cell viability (dotted line), error bars represent the standard deviation (SD) (DMSO: dimethyl sulfoxide) (Figure SI3)




Figure SI3. Effect on cell viability of selected synthesized compounds.