

## Computational study of hydrido boronium dications and comparison with the isoelectronic carbon analogs

Golam Rasul, George A. Olah, and G. K. Surya Prakash\*

Loker Hydrocarbon Research Institute and Department of Chemistry, University of Southern California,  
University Park, Los Angeles, CA 90089-1661

E-mail: [gprakash@usc.edu](mailto:gprakash@usc.edu)

Dedicated to Prof. Kenneth Laali on the occasion of his 65<sup>th</sup> Birthday and life-long contributions to chemistry


Received 03-14-2017

Accepted 06-19-2017

Published on line 07-30-2017

### Abstract

Calculations at the CCSD(T)/cc-pVTZ level show that higher coordinate, hydrido boronium dications are viable species despite their charge-charge repulsion. Structure and bonding of the studied boronium dications are discussed and are compared with their isoelectronic carbon analogs. Although the studied boronium dications are thermodynamically unstable towards deprotonation, their computational characterization supports their transient existence.



$\text{BH}_7^{2+} (\text{C}_{3v})$

**Keywords:** Hydrido boronium dications, higher coordinate, two-electron three-center bond, isoelectronic carbon analog

## Introduction

Main group hydrido cations and dications have been a subject of numerous experimental and theoretical investigations.<sup>1-6</sup> A while ago we have reported<sup>7</sup> the calculated structures of parent hexa- and heptacoordinate boronium ions,  $\text{BH}_6^+$  and  $\text{BH}_7^{2+}$ , respectively, with two and three two-electron three-center (2e-3c) bonds. The structures were found to be isostructural with their isoelectronic carbon analogs,  $\text{CH}_6^{2+}$ ,<sup>8,9</sup> and  $\text{CH}_7^{3+}$ ,<sup>10</sup> respectively. We have also previously investigated the structures and energetics of  $\text{BH}_n^{2+}$  ( $n=1-8$ )<sup>11</sup> and  $\text{B}_2\text{H}_{2n}^{2+}$  ( $n=1-4$ )<sup>12</sup> dications at the MP2/aug-cc-pVTZ and QCISD(T)/6-311G\*\* levels, respectively. In continuation of our study of higher coordinate onium ions, we have carried out a study on the structure and bonding of higher coordinate hydrido boronium dications. For comparison, all the structures were calculated at the same CCSD(T)/cc-pVTZ level.

## Results and Discussion


Gaussian 09 program<sup>13</sup> was employed for geometry optimizations and frequency calculations. The geometries were first optimized at the MP2/cc-pVTZ level. Vibrational frequencies at the MP2/cc-pVTZ//MP2/cc-pVTZ level were used to characterize stationary points as minima (NIMAG (number of imaginary frequency) = 0 or transition state NIMAG = 1) and to compute zero point vibrational energies (ZPE), which were scaled by a factor of 0.96.<sup>14</sup> The MP2/cc-pVTZ geometries were further optimized at the CCSD(T)/cc-pVTZ level. CCSD(T)/cc-pVTZ optimizations have been performed with the CFOUR program.<sup>15</sup> Computed energies are given in Table 1.

### $\text{BH}_n^{2+}$ ( $n=3-8$ ) systems

In line with previous calculations at the MP2/aug-cc-pVTZ level,<sup>11</sup> our present calculations at the CCSD(T)/cc-pVTZ level also shows that the  $C_{2v}$  symmetric structure **1** is a viable minimum for  $\text{BH}_3^{2+}$  (Figure 1). The structure **1** is characterized with a two-electron three-center (2e-3c) bond, a two-electron two-center (2e-2c) bond, resembling a complex between  $\text{BH}^{2+}$  and  $\text{H}_2$ . The B-H and H-H bond distances of the 2e-3c bond are computed to be 1.491 Å and 0.889 Å, respectively. Expectedly, both thermodynamically and kinetically the dication is unstable due to charge-charge repulsion. The dissociation of **1** into  $\text{BH}_2^+$  and  $\text{H}^+$  is exothermic by 100.6 kcal/mol and the process has a kinetic barrier of 8.2 kcal/mol at the CCSD(T)/cc-pVTZ//CCSD(T)/cc-pVTZ + ZPE level (Table 2).

Previously we have reported<sup>7</sup> the MP2/6-31G\*\* calculated structure of the boronium ion  $\text{BH}_4^+$  **2**. The structure of  $\text{BH}_4^+$  was found to be isostructural with the corresponding carbon analog  $\text{CH}_4^{2+}$ .<sup>16</sup> DePuy et al were able to generate  $\text{BH}_4^+$  **2** in the gas phase by the reaction of  $\text{BH}_2^+$  and  $\text{H}_2$ .<sup>17</sup> Removal of an electron from  $\text{BH}_4^+$  **2** leads to the dication  $\text{BH}_4^{2+}$  **3**. The planar  $D_{2h}$  symmetric form **3** (Figure 1) is found to be the only minimum on the potential energy surface of  $\text{BH}_4^{2+}$ . Structure **3** resembles a complex between  $\text{B}^{2+}$  with two hydrogen molecules resulting in formation of two 2e-3c bonds. Deprotonation of **3** has a substantial kinetic barrier of 24.7 kcal/mol (although exothermic by 68.7 kcal/mol, Table 1). Despite charge-charge repulsion the 2e-3c B-H bond distance of dication **3** (1.320 Å) is considerably shorter than that of monocation **2** (1.422 Å).


Protonation of  $\text{BH}_4^+$  **2** leads to the dication  $\text{BH}_5^{2+}$  **4**. We previously reported<sup>18</sup> the structure of  $\text{BH}_5^{2+}$  at the QCISD(T)/6-311G\*\* level. CCSD(T)/cc-pVTZ optimized structure of  $\text{BH}_5^{2+}$  **4** is now computed and displayed in Figure 1. The structure **4** resembles a complex between  $\text{BH}_2^+$  and two  $\text{H}_2$  molecules resulting in formation of two 2e-3c bonds with an empty p-orbital orthogonal to the plane of the molecule. The dissociation of **4** into  $\text{BH}_4^+$  **2** and  $\text{H}^+$  was calculated to be exothermic by 56.6 kcal/mol but has a high kinetic barrier of 36.5 kcal/mol (Table 2).



**Figure 1.** CCSD(T)/cc-pVTZ calculated structures of **1** – **8**.

The parent hexacoordinated boronium ion  $\text{BH}_6^+$  **5** is isoelectronic and isostructural with the corresponding carbon analog  $\text{CH}_6^{2+}$ .<sup>8,9</sup> The ion **5** was studied previously by us.<sup>2</sup> DePuy et al. were able to generate **5** in the gas phase by the reaction of  $\text{BH}_4^+$  **2** and  $\text{H}_2$ .<sup>12</sup> The  $C_{2v}$  symmetric form **5** was found to be the only minimum for  $\text{BH}_6^+$ . **5** contains two 2e-3c bonds and two 2c-2e bonds. Calculated B-H bond distance of 2e-3c interactions is found to be 1.366 Å. Removal of an electron from  $\text{BH}_6^+$  **5** leads to dication  $\text{BH}_6^{2+}$  **6**. The  $D_{3d}$  symmetric **6** is

a propeller-shaped molecule resembling a complex between  $B^{2+}$  with three hydrogen molecules resulting in formation of three 2e-3c bonds. B-H bond distance of 2e-3c interactions of the structure **6** is computed to be 1.337 Å. Deprotonation of **6** has a high kinetic barrier of 52.5 kcal/mol (Table 2).



**Figure 2.** Plot of relative energy of the structures, **2**, **4**, **5** and **7**.


Protonation of  $BH_6^+$  **5** leads to heptacoordinate boronium dication  $BH_7^{2+}$  **7**. The  $C_{3v}$  symmetric dication **7** is also propeller-shaped resembling a complex between  $BH^{2+}$  with three hydrogen molecules resulting in formation of three 2e-3c bonds. Average B-H bond distances of 2e-3c interactions of the structure **7** is found to be 1.359 Å. Deprotonation of **7** into  $BH_6^+$  **5** is exothermic by only 23.2 kcal/mol and the process has a high kinetic barrier of 55.1 kcal/mol at the CCSD(T)/cc-pVTZ//CCSD(T)/cc-pVTZ + ZPE level (Table 1). Relative energies of the structures **2**, **4**, **5**, **7** are plotted in Figure 2.

We also searched for any eight coordinate energy-minimum for dication  $BH_8^{2+}$ . The structure **8** was found to be a minimum on the potential energy surface of  $BH_8^{2+}$  (Figure 1). The  $C_{3v}$  symmetrical structure **8** can be considered as a weak complex between  $BH_6^{2+}$  **6** and  $H_2$  with a long bond distance of 1.844 Å between boron atom and  $H_2$  moiety.

### $B_2H_{2n}^{2+}$ (n=2-4) systems

In line with previous calculations at the QCISD(T)/6-311G\*\* level,<sup>12</sup> our present calculations at the CCSD(T)/cc-pVTZ level also shows that the  $C_s$  symmetric structure **9** and  $D_{2h}$  symmetric structure **10** are the viable minima for  $B_2H_4^{2+}$  (Figure 4). Structure **9** resembles a complex between  $B_2H_2^{2+}$  and  $H_2$  resulting in the formation of a 2e-3c bond. Structure **10** is characterized as a doubly hydrogen-bridged structure.

Structure **9** is 9.4 kcal/mol more stable than **10**. The dissociation of **9** into  $B_2H_3^+$  and  $H^+$  was computed to be exothermic by 6.7 kcal/mol but has a high kinetic barrier of 61.6 kcal/mol (Table 2).



**Figure 3.** CCSD(T)/cc-pVTZ calculated structures of **9** – **14**.

**Table 1.** Total energies (-au), ZPE (kcal/mol) and relative energies (RA in kcal/mol)

No.	MP2/cc-pVTZ	ZPE <sup>a</sup>	CCSD(T)/cc-pVTZ	RA <sup>b</sup>
<b>1</b>	25.39039	12.7	25.41983	
<b>2</b>	26.74205	21.9	26.78047	
<b>3</b>	25.96738	15.9	26.00121	
<b>4</b>	26.65701	25.2	26.69554	
<b>5</b>	27.94345	34.2	27.99208	
<b>6</b>	27.24920	31.5	27.29300	
<b>7</b>	27.91237	38.5	27.96199	
<b>8</b>	28.42359	38.9	28.47827	
<b>9</b>	50.90219	23.3	50.96293	0.0
<b>10</b>	50.88575	22.8	50.94717	9.4
<b>11</b>	52.13230	35.5	52.20481	0.7
<b>12</b>	52.13278	35.3	52.20553	0.0
<b>13</b>	52.12357	34.7	52.19450	6.3
<b>14</b>	53.35921	47.9	53.44053	

<sup>a</sup>ZPE at the MP2/cc-pVTZ//MP2/cc-pVTZ level scaled by a factor of 0.96;

<sup>b</sup>at the CCSD(T)/cc-pVTZ//CCSD(T)/cc-pVTZ + ZPE (MP2/cc-pVTZ//MP2/cc-pVTZ) level.

Three viable minimum structures (**11-13**) were computed for the diboronium dication,  $B_2H_6^{2+}$  (Figure 4). Triply hydrogen-bridged structure **12** was found to be the lowest minimum. This structure **12** also contains a 2e-3c bond involving a boron atom and two hydrogen atoms. One of the borons in structure **12** is six-coordinated and the other one is five-coordinated. Dication **11**, on the other hand, is doubly hydrogen-bridged structure. Additionally it also contains a 2e-3c bond involving a boron atom and two hydrogen atoms. Structure **11** is only 0.7 kcal/mol less stable than **12**. Structure **13**, unlike structures **11** and **12**, was characterized to be a non-hydrogen-bridged (involving both boron atoms) structure. Dication **13** with two 2e-3c bonds can be considered as a complex between  $B_2H_2^{2+}$  and two hydrogen molecules. Structure **13** is 6.3 kcal/mol less stable than the structure **12**. The dissociation of **12** into  $B_2H_5^+$  and  $H^+$  was calculated to be exothermic by 33.9 kcal/mol but has a kinetic barrier of 45.1 kcal/mol (Table 2).

**Table 2.** Dissociation Energy ( $\Delta E_0$ ) and Respective Kinetic Barrier at 298 K for the Selected Processes<sup>a</sup>

Process	$\Delta E_0$ (kcal/mol) <sup>b</sup>	Barrier (kcal/mol) <sup>b</sup>
$BH_3^{2+}$ ( <b>1</b> ) $\rightarrow$ $BH_2^+ + H^+$	-100.6	8.2
$BH_4^{2+}$ ( <b>3</b> ) $\rightarrow$ $BH_3^+ + H^+$	-68.7	20.2
$BH_5^{2+}$ ( <b>4</b> ) $\rightarrow$ $BH_4^+ + H^+$	-56.6	36.5
$BH_6^{2+}$ ( <b>6</b> ) $\rightarrow$ $BH_5^+ + H^+$	-25.8	52.5
$BH_7^{2+}$ ( <b>7</b> ) $\rightarrow$ $BH_6^+ + H^+$	-23.2	55.1
$B_2H_4^{2+}$ ( <b>9</b> ) $\rightarrow$ $B_2H_3^+ + H^+$	-6.7	61.6
$B_2H_6^{2+}$ ( <b>12</b> ) $\rightarrow$ $B_2H_5^+ + H^+$	-33.9	45.1

<sup>a</sup>at the CCSD(T)/cc-pVTZ//CCSD(T)/cc-pVTZ + ZPE (MP2/cc-pVTZ//MP2/cc-pVTZ level scaled by a factor of 0.96) level.

Structure **14** was computed for diprotonated diborane  $B_2H_8^{2+}$ . Dication **14** is a doubly hydrogen-bridged structure. It was also characterized to possess two 2e-3c bonds. Structure **14** (Figure 3) contains two six coordinate boron atoms. The structure formally resembles a complex between a  $B_2H_4^{2+}$  **10** with two hydrogen molecules resulting in formation of two 2e-3c bonds. Calculated central B-B bond length of **14** is computed to be 1.700 Å.

## Conclusions

Structures of higher coordinate hydrido boronium dications were calculated using CCSD(T)/cc-pVTZ level of theory and compared with their isoelectronic carbon analogs. These calculations indicate that higher coordinated doubly charged boronium dications are viable species in the gas phase despite strong charge-charge repulsion. In such small first row highly charged cations, charge-charge repulsions are clearly substantial, but the bonding interactions are strong enough to counter this. Although such hydrido boronium dications are thermodynamically unstable, their computational characterization gives support for their transient existence. Structures, stability and multicenter bonding concept of the intriguing higher coordinate boronium and carbonium (hypercarbon)<sup>19</sup> ions as well characterized compounds have now been

calculationally established. By development of new techniques, the characterization of these intriguing species is considered possible.

## Acknowledgements

Support of our work by the Loker Hydrocarbon Research Institute is gratefully acknowledged.

## References

1. Koch, W.; Schwarz, H.; *Structure / Reactivity and Thermochemistry of Ions*; Ausloos, P.; Lias, S. G., Eds.; Reidel: Dordrecht, 1987, p 413.  
[https://doi.org/10.1007/978-94-009-3787-1\\_21](https://doi.org/10.1007/978-94-009-3787-1_21)
2. Lammertsma, K.; Schleyer, P. v. R.; Schwarz, H. *Angew. Chem. Int. Ed. Engl.* **1989**, *28*, 1321.  
<https://doi.org/10.1002/anie.198913211>
3. Lammertsma, K. *Rev. Chem. Intermed.* **1988**, *9*, 141.  
<https://doi.org/10.1007/BF03155682>
4. Koch, W.; Maquin, F.; Stahl, D.; Schwarz, H. *Chimica* **1985**, *39*, 376.
5. Schleyer, P. v. R.; *Adv. Mass. Spectrom.* **1986**, *10*, 287.
6. Olah, G.A. *Angew. Chem. Int. Ed. Engl.* **1993**, *32*, 767.  
<https://doi.org/10.1002/anie.199307673>
7. Rasul, G.; Olah, G. A., *Inorg. Chem.*, **1997**, *36*, 1278.  
<https://doi.org/10.1021/ic960843k>
8. Lammertsma, K.; Olah, G. A.; Barzaghi, M.; Simonetta, M.; *J. Am. Chem. Soc.*, **1982**, *104*, 6851.  
<https://doi.org/10.1021/ja00388a095>
9. Lammertsma, K.; Barzaghi, M.; Olah, G. A.; Pople, J. A.; Schleyer, P. v. R.; Simonetta, M.; *J. Am. Chem. Soc.*, **1983**, *105*, 5258.  
<https://doi.org/10.1021/ja00354a013>
10. Olah, G. A.; Rasul, G.; *J. Am. Chem. Soc.*, **1996**, *118*, 8503.  
<https://doi.org/10.1021/ja961081v>
11. Rasul, G.; Dias, J. F.; Seidl, P. R.; Prakash, G. K. S., Olah, G. A., *J. Phys. Chem. A*, **2003**, *107*, 4731.  
<https://doi.org/10.1021/jp0226735>
12. Dias, J. F.; Rasul, G.; Seidl, P. R.; Prakash, G. K. S., Olah, G. A., *J. Phys. Chem. A*, **2003**, *107*, 7981.  
<https://doi.org/10.1021/jp0307405>
13. Gaussian 09, Revision A.02, Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Scalmani, G.; Barone, V.; Mennucci, B.; Petersson, G. A.; Nakatsuji, H.; Caricato, M.; Li, X.; Hratchian, H. P.; Izmaylov, A. F.; Bloino, J.; Zheng, G.; Sonnenberg, J. L.; Hada, M.; Ehara, M.; Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Vreven, T.; Montgomery, Jr., J. A.; Peralta, J. E.; Ogliaro, F.; Bearpark, M.; Heyd, J. J.; Brothers, E.; Kudin, K. N.; Staroverov, V. N.; Kobayashi, R.; Normand, J.; Raghavachari, K.; Rendell, A.; Burant, J. C.; Iyengar, S. S.; Tomasi, J.; Cossi, M.; Rega, N.; Millam, N. J.; Klene, M.; Knox, J. E.; Cross, J. B.; Bakken, V.; Adamo, C.;

- Jaramillo, J.; Gomperts, R.; Stratmann, R. E.; Yazyev, O.; Austin, A. J.; Cammi, R.; Pomelli, C.; Ochterski, J. W.; Martin, R. L.; Morokuma, K.; Zakrzewski, V. G.; Voth, G. A.; Salvador, P.; Dannenberg, J. J.; Dapprich, S.; Daniels, A. D.; Farkas, Ö.; Foresman, J. B.; Ortiz, J. V.; Cioslowski, J.; Fox, D. J. Gaussian, Inc., Wallingford CT, 2009.
14. Alecu, I. M.; Zheng, J.; Zhao, Y.; Truhlar, D. G. *J. Chem. Theory Comput.* **2010**, *6*, 2872.  
<https://doi.org/10.1021/ct100326h>
15. CFOUR, a quantum chemical program package, Stanton, J. F.; Gauss, J.; Harding, M. E.; Szalay, P. G. with contributions from Auer, A. A.; Bartlett, R. J.; Benedikt, U.; Berger, C.; Bernholdt, D. E.; Bomble, Y. J.; Cheng, L.; Christiansen, O.; Heckert, M.; Heun, O.; Huber, C.; Jagau, C.; Jonsson, D.; Jusélius, J.; Klein, K.; Lauderdale, W. J.; Mathews, D. A.; Metzroth, T.; O'Neill, D. P.; Price, D. R.; Prochnow, E.; Ruud, K.; Schiffmann, F.; Schwalbach, W.; Stopkowicz, S.; Tajti, A.; Vázquez, J.; Wang, F.; Watts, J. D.; and the integral packages MOLE-CULE (Almlöf, J and Taylor, P. R.), PROPS (Taylor, P. R.), ABACUS (Helgaker, T.; Aa, H. J.; Jensen H. J. A.; Jørgensen, P.; Olsen, J.), and ECP routines (Mitin, A. V.; Wüllen, C. van), **2010**; Harding, M. E.; Metzroth, T.; Gauss, J.; Auer, A. A. *J. Chem. Theory Comput.* **2008**, *4*, 64.
16. Wong, M. W.; Radom, L. *J. Am. Chem. Soc.* **1989**, *111*, 1155.  
<https://doi.org/10.1021/ja00185a070>
17. DePuy, C. H.; Gareyev, R.; Hankin, J.; Davico, G. E. *J. Am. Chem. Soc.* **1997**, *119*, 427.  
<https://doi.org/10.1021/ja963399t>
18. Olah, G. A.; Rasul, G. *J. Am. Chem. Soc.*, **1996**, *118*, 12922.  
<https://doi.org/10.1021/ja961597e>
19. Olah, G. A.; Prakash, G. K. S.; Wade, K.; Molnar, A.; Williams, R. E. *Hypercarbon Chemistry*, John Wiley & Sons: New York, 2011.  
<https://doi.org/10.1002/9781118016466>