Supplementary Material

The synthesis of fluorine-containing endothiopeptide analogs by the reaction of perfluorinated dithiocarboxylic acid amides with esters of α -amino acids and dipeptides

Nadiia V. Pikun,^a Sergiy S. Mykhaylychenko,^a and Yuriy G. Shermolovich^{*a}

^aInstitute of Organic Chemistry, National Academy of Sciences of Ukraine, 5, Murmanska, 02094, Kiev, Ukraine <u>sherm@ioch.kiev.ua</u> corresponding author

Table of Contents

1.	¹⁹ F NMR Spectra of dimethyl N,N'-(2,2-difluoropropanedithioyl)	S2
	bis(L-phenylalaninate) 8b (in CDCl ₃)	
2.	¹ H NMR Spectra of dimethyl N,N'-(2,2-difluoropropanedithioyl)	S2
	bis(L-phenylalaninate) 8b (in CDCl ₃)	
3.	¹³ C NMR Spectra of dimethyl N,N'-(2,2-difluoropropanedithioyl)	S3
	bis(L-phenylalaninate) 8b (in CDCl ₃)	
4.	¹⁹ F NMR Spectra of dimethyl N,N'-(2,2-difluoropropanedithioyl)	S3
	bis(DL- α -alanyl- β -alaninate) 9a (in CDCl ₃)	
5.	¹ H NMR Spectra of dimethyl N,N'-(2,2-difluoropropanedithioyl)	S4
	bis(DL- α -alanyl- β -alaninate) 9a (in CDCl ₃)	
6.	¹³ C NMR Spectra of dimethyl N,N'-(2,2-difluoropropanedithioyl)	S5
	bis(DL-α-alanyl-β-alaninate) 9a (in CDCl ₃)	

¹⁹F NMR Spectra of dimethyl N,N'-(2,2-difluoropropanedithioyl) bis(L-phenylalaninate) **8b** (in CDCl₃)



¹H NMR Spectra of dimethyl N,N'-(2,2-difluoropropanedithioyl) bis(L-phenylalaninate) **8b** (in CDCl₃)



¹³C NMR Spectra of dimethyl N,N'-(2,2-difluoropropanedithioyl) bis(L-phenylalaninate) **8b** (in CDCl₃)



¹⁹F NMR Spectra of dimethyl N,N'-(2,2-difluoropropanedithioyl) bis(DL-α-alanyl-βalaninate) **9a** (in CDCl₃)



¹H NMR Spectra of dimethyl N,N'-(2,2-difluoropropanedithioyl) bis(DL- α -alanyl- β -alaninate) **9a** (in CDCl₃)



¹³C NMR Spectra of dimethyl N,N'-(2,2-difluoropropanedithioyl) bis(DL- α -alanyl- β alaninate) **9a** (in CDCl₃)

