

Supplementary Material

Ring-closing metathesis as a key step to construct the 2,6-dihydropyrano[2,3-c]pyrazole ring system

Aurimas Bieliauskas,^{a,b} Sonata Krikštolaitytė,^b Wolfgang Holzer,^c and Algirdas Šačkus^{a,b}

^aInstitute of Synthetic Chemistry, Kaunas University of Technology, K. Baršausko g. 59, Kaunas LT-51423, Lithuania

^b Department of Organic Chemistry, Kaunas University of Technology, Radvilėnų pl. 19, Kaunas LT-50254, Lithuania

^c Department of Pharmaceutical Chemistry, University of Vienna, Althanstrasse 14, Vienna 1090, Austria
Email: algirdas.sackus@ktu.lt

Table of Contents

1. ¹ H NMR spectrum of compound 5a	S2
2. ¹³ C NMR spectrum of compound 5a	S3
3. DEPT-135 ¹³ C NMR spectrum of compound 5a	S4
4. ¹ H- ¹⁵ N HMBC NMR spectrum of compound 5a	S5
5. The overlaid ¹ H- ¹³ C gs-HSQC and 60 Hz 1,1- ADEQUATE NMR spectra of compound 5a	S6




Figure S2. ^1H NMR spectrum of compound 5a.




Figure S3. ^{13}C NMR spectrum of compound 5a.




Figure S4. DEPT-135 ¹³C NMR spectrum of compound 5a.




Figure S5. ^1H - ^{15}N HMBC NMR spectrum of compound 5a.




Figure S6. The overlaid ^1H - ^{13}C gs-HSQC (black) and 60 Hz 1,1- ADEQUATE (red) NMR spectra of compound 5a