

Addition of bis(trimethylsilyl)ketene acetals to activated 2-(pyridin-3-yl)-1,3-benzothiazole: Synthesis and cytotoxic activity of novel carboxylic acids and δ -bromolactone derivatives

Ricardo Ballinas-Indili,^a Ricardo Corona-Sánchez,^b Saúl R. Mereciás,^a M. Teresa Ramírez-Apan,^a Alfredo Toscano,^a Leticia Lomas-Romero,^b Roberto Guerrero-Reyes,^a and Cecilio Álvarez-Toledano*^a

^a Instituto de Química, Universidad Nacional Autónoma de México, 04510 Ciudad de México, México

^b Departamento de Química, Universidad Autónoma Metropolitana-Iztapalapa, Av. San Rafael Atlixco 186, Leyes de Reforma 1ra Secc., 09340 Ciudad de México, México

Email: cecilio@unam.mx


Received 01-07-2019

Accepted 06-03-2019

Published on line 06-21-2019

Abstract

The nucleophilic addition of the alkyl substituted bis(trimethylsilyl)ketene acetals to 2-(pyridin-3-yl)-1,3-benzothiazole in the presence of triflic anhydride has been examined. The behavior of the chosen substrate in these transformations is peculiar due to an exclusive activation of its pyridine fragment, resulting in the exclusive formation of 3-benzothiazolyl-1,4-dihydropyridine carboxylic acids, as confirmed by single crystal X-ray crystallography. These isolated acids have been efficiently turned into novel δ -bromolactones by the regioselective ring closing procedure promoted by NBS. *In vitro* cytotoxic activities of the synthesized carboxylic acids and lactones have shown moderate anticancer activity against six human tumor cell lines; the derivatives bearing a cyclobutyl group present the highest activity.



Keywords: Benzothiazole, ketene acetals, chemoselectivity, carboxylic acids, lactones, cytotoxic activity

Introduction

Nitrogen-containing heterocycles play an important role not only in the life sciences, but also in many industrial fields related to fine chemistry.¹ Among nitrogen-containing heterocycles, benzothiazoles and pyridines stand out as two classes of bioactive and industrially important organic compounds. Molecules with a benzothiazole ring are attractive targets for synthesis since they often exhibit diverse and important biological properties.² These compounds possess potential application as anticancer,³ antimicrobial,⁴ anti-convulsant,⁵ antiviral,⁶ anthelmintic,⁷ antifungal,⁸ analgesic,⁹ anti-inflammatory,¹⁰ antidiabetic,¹¹ etc. Thereby, the benzothiazole ring system is considered as a privileged structure.¹²

Likewise, pyridine derivatives have occupied a privileged position in the field of medicinal chemistry.¹³ The incorporation of the pyridine moiety into other organic structures is an important synthetic strategy in drug discovery.¹⁴ The high therapeutic properties of the pyridine-containing drugs have encouraged the organic chemists to synthesize a large number of novel pyridine compounds with important biologically active structures.^{15,16} Similarly, 1,4-dihydropyridine (1,4-DHP) nucleus is a privileged scaffold that, if appropriately substituted, exerts potent and selective action at diverse receptors, ion channels and enzymes.^{17,18} The ability of 1,4-DHP to revert multi drug resistance continues to be of great interest in medicinal chemistry.¹⁹

Among the synthetic methods to obtain 1,4-DHPs, the addition of nucleophiles to N-activated pyridines stands out from others, as it allows access to derivatives with high regio- and stereoselectivity.²⁰ In this context, our group have developed new methods based upon the inherent reactivity of bis(trimethylsilyl)ketene acetals to act as 1,3-carbon-oxygen dinucleophiles to allow access to novel carboxylic acid substituted 1,4-DHP and other dihydro(aza-aromatic) compounds using pyridine or other wide-ranging aza-aromatic substrates.²¹ These transformations occur by activation of nitrogen atoms *via* the formation of the corresponding salts with triflic anhydride or by interaction with alkyl chloroformates. The isolable carboxylic acid intermediates could be turned into novel functionalized γ - and δ -lactones through a subsequent lactonization step,²²⁻²⁴ and these often show a broad range of biological activities.²⁵


As was early reported by Ito,²⁶ the addition of silyl enol ethers to N-alkoxycarbonyl quaternary salts of thiazole or benzothiazole in the presence of triethylamine affords the corresponding 2-substituted 4-thiazolines or 2,3-dihydrobenzothiazoles in good yields. Thus, one might expect that either the pyridine or the benzothiazole (or both) components of 2-(pyridin-3-yl)-1,3-benzothiazole **4** would be involved in the reaction with bis(trimethylsilyl)ketene acetals **2** to furnish novel carboxylic acid or lactone scaffolds which may exhibit interesting biological properties.

In continuation of our recent research, we report herein the synthesis of novel carboxylic acid substituted 1,4-dihydropyridines **5** and bicyclic δ -lactones **6** bearing a benzothiazole ring from activated 2-(pyridin-3-yl)-1,3-benzothiazole **4** and different bis(trimethylsilyl)ketene acetals **2**. The anticancer properties of the compounds were evaluated *in vitro* against a panel of human tumor cell lines.

Results and Discussion

Synthesis

Firstly, we were interested in evaluating the reactivity of N-triflyl quaternary salt of benzothiazole towards the nucleophilic attack of bis(trimethylsilyl)ketene acetals. When benzothiazole **1** was treated with triflic anhydride during 30 min at -78 °C followed by the addition of bis(trimethylsilyl)ketene acetals **2a-d**, 2-substituted 2,3-dihydrobenzothiazoles **3a-d** were obtained in moderate to good yields (Table 1, entries 1-4).


Table 1. Synthesis of 2,3-dihydrobenzothiazoles **3a-e** using different bis(trimethylsilyl)ketene acetals^a

Entry	Acetal	$\text{R}^1 + \text{R}^2$	Product	Yield ^b %
1	2a	CH_3	3a	86
2	2b	$-(\text{CH}_2)_3-$	3b	62
3	2c	$-(\text{CH}_2)_4-$	3c	59
4	2d	$-(\text{CH}_2)_5-$	3d	65

^a Reaction conditions: 1.0 mmol of benzothiazole **1**, 1.1 mmol of Tf_2O , 1.2 mmol of bis(TMS)ketene acetal **2a-d** and 10 mL of CH_2Cl_2 . ^b Isolated yield after SiO_2 column chromatography.

Thus, we have demonstrated the viability of bis(trimethylsilyl)ketene acetals to act as nucleophiles in their reaction with N-activated benzothiazoles. Thereafter, we focus in the issue of competition between the benzothiazole and pyridine fragments to be activated by triflic anhydride by using 2-(pyridin-3-yl)-1,3-benzothiazole **4** as substrate. This issue was initially addressed by reacting triflic anhydride with **4** followed by addition of bis(trimethylsilyl)ketene acetal **2a**. The precursor compound **4** was efficiently prepared by SiO_2 -catalyzed condensation of 2-aminothiophenol and 3-pyridinecarboxaldehyde under microwave irradiation for 20 min in an air atmosphere. Based on our previous findings,²¹⁻²⁵ when 1.1 equiv of Tf_2O was added to a solution of **4** in dry CH_2Cl_2 at -78°C and subsequently treated with 1.2 equiv of ketene acetal **2a** for 2 hours at room temperature, the corresponding 1,4-dihydropyridine **5a** was obtained as white solid in 90% yield. Thus, selective activation of **4** took place exclusively at the pyridine fragment. Attempts to introduce an extra triflyl group at the benzothiazole moiety by increasing the amount of Tf_2O failed, even when the reaction time with triflic anhydride was increased probably due to the steric hindrance of the bulky triflyl groups.

Afterward, under conditions described earlier, the reaction was tested with other nucleophiles bearing different cycloalkyl substituents (Table 2). The reaction is successful for the four ketene acetals used and gives very good yields of the resulting carboxylic acids, after recrystallization from hexane / dichloromethane. Compounds **5a-d** were characterized by various spectroscopic techniques. According to their physical data, the observed products result from regioselective nucleophilic addition of the ketene acetal to C-4 of the pyridine fragment. In the ^1H NMR spectrum of **5a**, the evidence for the addition of ketene acetal **2a** to the pyridine component was supported by signals appearing as doublets at 6.94 (H-13) and 4.50 ppm (H-15), as well as a doublet of doublets at 5.63 ppm (H-14) corresponding to the three vicinal protons of the 1,4-DHP moiety. The ^{13}C NMR spectra confirmed the incorporation of a carboxylic acid by the signal appearing at 176.2 ppm and the introduction of a triflyl group by showing a quartet signal at 119.4 ppm. In addition, the structural arrangement of **5b** was unequivocally established by X-ray diffraction analysis (Figure 1).

Table 2. Synthesis of 1,4-dihydropyridines **5a-d** using different bis(trimethylsilyl)ketene acetals^a

Entry	Ketene acetal	Product	Yield ^b %
1	2a 	5a 	90
2	2b 	5b 	88
3	2c 	5c 	93
4	2d 	5d 	79

^a Reaction conditions: 1.0 mmol of 2-(pyridin-3-yl)-1,3-benzothiazole **4**, 1.1 mmol of Tf_2O , 1.2 mmol of bis(trimethylsilyl)ketene acetal **2a-d** and 25 mL of CH_2Cl_2 . ^b Isolated yield after purification by recrystallization from hexane/dichloromethane.





Figure 1. ORTEP view of compound **5b**. Thermal ellipsoids are at 30% probability level.

We next turned our attention to the synthesis of novel substituted δ -lactones. Since carboxylic acid bearing dihydropyridines are known to produce lactones by using typical hydroxy-, halo- or protolactonization conditions,²⁷ the carboxylic acids **5** were transformed to the novel bicyclic δ -lactones **6** under established bromolactonization conditions which involve the addition of 1.1 equiv. of NBS to a dichloromethane solution of the carboxylic acids **5a-d** and 20 mol % of TBAB to give, after 2 h at 0 °C and silica gel purification, white solids in moderate to good yields (Table 3).

Table 3. Synthesis of δ -bromolactones **6a-c**^a



Entry	Carboxylic acid	R ¹ + R ²	Product	Yield ^b %
1	5a	-CH ₃	6a	88
2	5b	-(CH ₂) ₃ -	6b	90
3	5c	-(CH ₂) ₄ -	6c	79
4	5d	-(CH ₂) ₅ -	6d	65

^a Reaction conditions: 0.5 mmol of **5a-d**, 0.55 mmol of NBS, 0.6 mmol NaHCO₃, 0.1 mmol of TBAB and 20 mL of CH₂Cl₂. ^b Isolated yield after SiO₂ column chromatography.

The observed products result from a regioselective ring-closing step as was confirmed by spectroscopic data of compounds **6a-d**. For example, high resolution mass spectrometry (ESI-TOF) confirmed the expected mass $m/z = 510.96142$ ($[M + 1]^+$) for **6a** in accordance with the molecular formula. Characteristic signals of protons H-13 and H-15 of compound **6a** were observed in ¹H NMR as doublets at 6.94 and 4.50 ppm ($J = 6.1$ Hz) respectively, while proton H-14 appears as a doublet of doublets signal at 5.63 ppm with the same

coupling constant. Lactone **6a** formation was verified in the ^{13}C NMR spectrum by the appearance of the lactone carbonyl signal at 171.5 ppm and the alkyl oxygen-base carbon (site of lactone ring closure) at 82.9 ppm, as well as in the IR spectrum where characteristic lactone carbonyl group appears at 1761 cm^{-1} .

Cytotoxic activity

The cytotoxicity of the two-different series of compounds **5a–d** and **6a–d** toward different cancer cell lines, including human glioblastoma (CNS U251), human prostatic adenocarcinoma (PC-3), human chronic myelogenous leukemia (K562), human colorectal adenocarcinoma (HCT-15), human mammary adenocarcinoma (MCF-7), and non-small cell lung cancer (SKLU) were determined by using the protein-binding dye sulforhodamine B (SRB) assay in microculture to determine cell growth.²⁸

Table 4. Inhibition of the growth (%) of human tumor cell lines for **5a–d** and **6a–d** at 50 μM in DMSO

Entry	Compound	Cell lines					
		U251	PC-3	K562	HCT-15	MCF-7	SKLU-1
1	5a	46.2	33.4	27.0	12.5	45.0	28.7
2	5b	60.3	35.4	33.7	32.7	61.0	50.0
3	5c	35.2	27.6	19.4	26.9	45.6	38.8
4	5d	22.9	23.3	6.3	10.7	21.5	29.6
5	6a	33.4	64.7	64.7	68.2	75.6	67.7
6	6b	87.9	78.6	81.7	96.1	100	100
7	6c	41.8	39.6	33.1	37.1	53.6	54.2
8	6d	33.1	33.1	39.9	21.8	35.3	31.8

The initial cytotoxicity screening data listed in Table 4 show moderate to good activities toward almost all tumor cell lines. From those data, we observe that in general lactones **6** were more active than their corresponding carboxylic acids **5**. Regarding the carboxylic acids, the most promising results were obtained from compound **5b**, which shows the highest cytotoxic activity against U251, MCF-7, and SKLU-1 tumor cell lines, whereas the least active compound was the compound **5d**. Similarly, as with carboxylic acids, the lactone **6b** with a cyclobutyl group exhibits the highest activity against the six tumor cell lines tested. It is worth noting that the activity of tested lactones decreases considerably for all tumor cell lines when the size of the cycloalkane ring is increased (Table 4, entries 6-8). Finally, we determined the IC_{50} of the most active compounds over HCT-15, MCF-7 and SKLU-1 and compared with a well-known chemotherapeutic drug (cisplatin) and Gefinitib which, like some benzothiazole derivatives,²⁹ is a potent epidermal growth factor receptor (EGFR) inhibitor (Table 5).

Table 5. IC₅₀ (μM) of human tumor cell lines for compounds **5b**, **6b**, Cisplatin, and Gefitinib in DMSO

Entry	Compound	Cell lines			
		HCT-15	MCF-7	SKLU-1	COS7
1	5b	86.8 ± 6.5	71.6 ± 4.3	74.7 ± 4.3	92.2 ± 1.3
2	6b	40.0 ± 4.7	27.4 ± 0.5	54.0 ± 3.1	47.7 ± 1.2
3	Cisplatin	21.4 ± 0.9	16.6 ± 0.4	2.2 ± 0.1	6.4 ± 1.5
4	Gefitinib	10.4 ± 1.8	9.0 ± 0.8	9.7 ± 0.9	3.2 ± 0.4

Analyzing these results, we observe that the compounds **5b** and **6b** have higher IC₅₀ values in comparison with references Cisplatin and Gefitinib, which indicates their moderate activity against the evaluated tumor cell lines. Additionally, compounds were also tested against normal monkey kidney fibroblast (COS7 cell line). In general, IC₅₀ values determined for COS7 cells are higher than those found for cancer cell lines, which means that this type of compounds exhibit certain selectivity towards cancer cell lines.

Conclusions

We have demonstrated that bis(TMS)ketene acetals react efficiently with *N*-triflil quaternary salt of benzothiazole to affords 2-substituted 2,3-dihydrobenzothiazoles. In addition, a simple method for the regioselective synthesis of 4-substitued 1,4-dihydropyridines bearing a benzothiazole moiety was developed from 2-(pyridin-3-yl)-1,3-benzothiazole and different *bis*(trimethylsilyl)ketene acetals by using triflic anhydride as activating agent. In these reactions, *N*-activation of the pyridine moiety is preferred over benzothiazole fragment. Using a regioselective bromolactonization protocol, novel bicyclic δ-bromolactones were accessed. Finally, we have demonstrated that carboxylic acids and lactones shown moderate anticancer activity against human tumor cell lines where the derivatives bearing a cyclobutyl group present the highest activity.

Experimental Section

General. All the reactions were performed under an inert atmosphere of nitrogen or argon. All reagents were obtained from commercial suppliers and used without further purification. Merck silica gel (type 60, 0.063-0.200 mm) was used for column chromatography. All compounds were characterized by IR spectra, recorded on a Perkin-Elmer 283B or 1420 spectrophotometer, by means of ATR or KBr techniques, and all data are expressed as wavenumbers (cm⁻¹). Melting points were obtained on a Melt-Temp II apparatus. NMR spectra were recorded on a Bruker Avance III at 300 MHz (1H NMR) and 75 MHz (13C NMR) in chloroform-*d*, acetone-*d*₆ or DMSO-*d*₆. Chemical shifts are given in ppm with TMS as the reference. The MS-DART data were obtained on a Jeol AccuTOF; the values of the signals are expressed as mass/charge units (*m/z*). Microwave irradiation experiments were performed using a Monowave 300 single-mode microwave reactor. The reusable 35 mL Pyrex vial is sealed with PEEK snap caps and standard PTFE coated silicone septa. Reaction cooling is performed using compressed air automatically after the heating period has elapsed. Suitable X-ray quality crystals of **5b** were grown by slow evaporation of a *n*-hexane-dichloromethane mixture at -5°C. The crystals were mounted on a glass fiber at room temperature, and then placed on a Bruker Smart Apex CCD

diffractometer, equipped with Mo K α radiation; decay was negligible in both cases. Systematic absences and intensity statistics were used in the space group determination. The structure was solved using direct methods. Anisotropic structure refinements were achieved using a full matrix, least-squares technique on all non-hydrogen atoms. All hydrogen atoms were placed in idealized positions, based on hybridization, with isotropic thermal parameters fixed at 1.2 times the value of the attached atom. Structure solutions and refinements were performed using SHELXTL V6.10.

Synthesis of bis(trimethylsilyl)ketene acetals. The syntheses of ketene acetals 2a-d were carried out using the methodology previously described elsewhere.³⁰

Synthesis of 2,3-dihydrobenzothiazoles 3a-e, general procedure. Benzothiazole 1 (1.0 mmol) in anhydrous dichloromethane (10 mL) were placed in a 100 mL round-bottom flask under nitrogen atmosphere. Trifluoromethanesulfonic anhydride (0.12 mL, 1.1 mmol) was added by syringe at -78 °C and the mixture was stirred at this temperature for 30 min. After this time, the corresponding *bis*(trimethylsilyl)ketene acetal 2a-d (1.2 mmol) was added. The mixture was warmed to room temperature and stirred for 4 h. The crude product was diluted with dichloromethane and washed with water and brine. The organic layer was dried over Na₂SO₄ and the solvent removed. Finally, the 2,3-dihydrobenzothiazoles were purified by silica gel column chromatography, eluting with a solvent mixture of hexane/ethyl acetate.

2-Methyl-2-{3-[(trifluoromethyl)sulfonyl]-2,3-dihydro-1,3-benzothiazol-2-yl}propanoic acid (3a). White crystals (305 mg, 86%), mp 147-148 °C. ¹H NMR (300 MHz, Acetone-*d*₆, ppm): δ 8.57 (s, 1H, OH), 7.55 (dd, *J* 7.8, 1.5 Hz, 1H, H-3), 7.45 (dd, *J* 7.6, 1.6 Hz, 1H, H-6), 7.32 (td, *J* 7.5, 1.5 Hz, 1H, H-2), 7.25 (td, *J* 7.7, 1.5 Hz, 1H, H-1), 6.06 (s, 1H, H-8), 1.39 (s, 3H, CH₃), 1.06 (s, 3H, CH₃). ¹³C NMR (75 MHz, Acetone-*d*₆, ppm): δ 175.0 (C=O), 136.4 (C-4), 133.0 (C-5), 128.4 (C-2), 126.1 (C-6), 122.8 (C-1), 120.3 (q, CF₃, *J* 327 Hz), 119.4 (C-3), 73.6 (C-8), 50.6 (C-10), 22.9 (CH₃), 17.0 (CH₃). IR (KBr, cm⁻¹): ν_{\max} 1704 (C=O). MS (DART, *m/z*): 373 [M+18]⁺, 356 [M+H]⁺. HRMS (ESI-TOF): calcd for C₁₂H₁₃F₃NO₄S₂ [M+H]⁺ 356.02381; found 356.02362.

1-{3-[(Trifluoromethyl)sulfonyl]-2,3-dihydro-1,3-benzothiazol-2-yl}cyclobutane-1-carboxylic acid (3b). White crystals (227 mg, 62%), mp 140-142 °C. ¹H NMR (300 MHz, Acetone-*d*₆, ppm): δ 10.18 (s, 1H, OH), 7.62 (d, 1H, *J* 7.7 Hz, H-3), 7.43 (dd, *J* 7.3, 1.9 Hz, 1H, H-6), 7.24-7.33 (m, 2H, H-2 and H-1), 6.10 (s, 1H, H-8), 2.67-2.57 (m, 1H, CH₂ cyclobutyl), 2.44-2.26 (m, 2H, CH₂ cyclobutyl), 2.02-1.78 (m, 3H, CH₂ cyclobutyl). ¹³C NMR (75 MHz, Acetone-*d*₆, ppm): δ 174.2 (C=O), 135.9 (C-4), 132.8 (C-5), 128.4 (C-2), 126.1 (C-6), 122.8 (C-1), 120.3 (q, CF₃, *J*=327 Hz), 119.2 (C-3), 70.8 (C-8), 54.5 (C-10), 27.4 (CH₂ cyclobutyl), 24.3 (CH₂ cyclobutyl), 15.3 (CH₂ cyclobutyl). IR (KBr, cm⁻¹): ν_{\max} 1705 (C=O). MS (DART, *m/z*): 385 [M+18]⁺, 367 [M]⁺. HRMS (ESI-TOF): calcd for C₁₃H₁₃F₃NO₄S₂ [M+H]⁺ 367.01598; found 367.01518.

1-{3-[(Trifluoromethyl)sulfonyl]-2,3-dihydro-1,3-benzothiazol-2-yl}cyclopentane-1-carboxylic acid (3c). White crystals (225 mg, 59%), mp 146-147 °C. ¹H NMR (300 MHz, DMSO-*d*₆, ppm): δ 13.08 (s, 1H, OH), 7.50 (dd, *J* 7.7, 1.5 Hz, 1H, H-3), 7.46 (dd, *J* 7.9, 1.3 Hz, 1H, H-6), 7.30 (td, *J* 7.6, 1.5 Hz, 1H, H-2), 7.23 (td, *J* 7.7, 1.5 Hz, 1H, H-1), 6.09 (s, 1H, H-8), 2.01-1.87 (m, 3H, CH₂ cyclopentyl), 1.66-1.32 (m, 5H, CH₂ cyclopentyl). ¹³C NMR (75 MHz, DMSO-*d*₆, ppm): δ 175.9 (C=O), 135.8 (C-4), 133.3 (C-5), 129.1 (C-2), 126.6 (C-6), 123.4 (C-1), 120.2 (q, CF₃, *J*=327 Hz), 119.7 (C-3), 72.7 (C-8), 61.9 (C-10), 33.5 (CH₂ cyclopentyl), 30.0 (CH₂ cyclopentyl), 25.3 (CH₂ cyclopentyl), 25.1 (CH₂ cyclopentyl). IR (KBr, cm⁻¹): ν_{\max} 1706 (C=O). MS (DART, *m/z*): 399 [M+18]⁺, 382 [M+H]⁺. HRMS (ESI-TOF): calcd for C₁₄H₁₅F₃NO₄S₂ [M+H]⁺ 382.03946; found 382.03880.

1-{3-[(Trifluoromethyl)sulfonyl]-2,3-dihydro-1,3-benzothiazol-2-yl}cyclohexane-1-carboxylic acid (3d). White crystals (257 mg, 65%), mp 168-170 °C. ¹H NMR (300 MHz, DMSO-*d*₆, ppm): δ 12.94 (s, 1H, OH), 7.49 (dd, *J* 7.6, 1.5 Hz, 1H, H-3), 7.43 (dd, *J* 7.8, 1.4 Hz, 1H, H-6), 7.28 (td, *J* 7.6, 1.5 Hz, 1H, H-2), 7.22 (td, *J* 7.7, 1.6 Hz, 1H, H-1), 5.85 (s, 1H, H-8) 1.88 (dd, *J* 30.8, 11.9 Hz, 2H, CH₂ cyclohexyl), 1.54 (d, *J* 10.4 Hz, 3H, CH₂

cyclohexyl), 1.33-1.01 (m, 5H, CH₂ cyclohexyl). ¹³C NMR (75 MHz, DMSO-*d*₆, ppm): δ 173.4 (C=O), 136.1 (C-4), 132.5 (C-5), 128.9 (C-2), 126.6 (C-6), 123.3 (C-1), 120.2 (q, CF₃, *J* 327 Hz), 119.3 (C-3), 75.4 (C-8), 55.1.9 (C-10), 29.3 (CH₂ cyclohexyl), 28.8 (CH₂ cyclohexyl), 25.2 (CH₂ cyclohexyl), 22.9 (CH₂ cyclohexyl), 22.8 (CH₂ cyclohexyl). IR (KBr, cm⁻¹): ν_{max} 1701 (C=O). MS (DART, *m/z*): 413 [M+18]⁺, 396 [M+H]⁺. HRMS (ESI-TOF): calcd for C₁₅H₁₇F₃NO₄S₂ [M+H]⁺ 396.05511; found 396.05383.

Synthesis of 2-(pyridin-3-yl)-1,3-benzothiazole (4). To a 35 mL microwave vial equipped with a magnetic stirrer were added 2-aminothiophenol (1 g, 7.98 mmol), 3-pyridinecarboxaldehyde (854 mg, 7.98 mmol) and SiO₂ (24 mg, 0.4 mmol) in EtOH (15 mL). The reaction mixture was irradiated at 70 °C for 20 min. After cooling to room temperature, the reaction was filtered, and the solvent was evaporated under reduced pressure. The crude product was purified by silica gel column chromatography with a solvent mixture of hexane/ethyl acetate. Yellow powder (1.25 g, 74%), mp 135-136 °C. ¹H NMR (300 MHz, CDCl₃, ppm): δ 9.31 (s, 1H, H-11), 8.73 (dd, *J* 4.8, 1.6 Hz, 1H, H-13), 8.39 (dt, *J* 8.0, 2.0 Hz, 1H, H-3), 8.12 (d, *J* 8.1 Hz, 1H, H-15), 7.94 (d, 1H, *J* 7.6 Hz H-6), 7.57-7.52 (m, 1H, H-2), 7.47-7.4 (m, 2H, H-1, H-14). ¹³C NMR (75 MHz, CDCl₃, ppm): δ 164.57 (C-8), 153.95 (C-5), 151.62 (C-13), 148.61 (C-11), 134.96 (C-4), 134.52 (C-15), 129.68 (C-10), 126.64 (C-1), 125.72 (C-14), 123.79 (C-2), 123.49 (C-6), 121.75 (C-3). The data obtained are in agreement with that previously reported.³¹

Synthesis of 1,4-dihydropyridines 5a-d, general procedure. 2-(Pyridin-3-yl)-1,3-benzothiazole 4 (212 mg, 1.0 mmol) and 35 mL of anhydrous dichloromethane were placed in a 100 mL round-bottom flask under a nitrogen atmosphere. Trifluoromethanesulfonic anhydride (0.17 mL, 1.1 mmol) was added by syringe at -78 °C and the mixture was stirred at this temperature for 30 min. After this time, the corresponding *bis*(trimethylsilyl)ketene acetal 2a-d (1.2 mmol) was added. The mixture was warmed to room temperature and stirred for 8 h. The crude product was diluted with 20 mL of dichloromethane and washed with water and brine. The organic layer was dried over Na₂SO₄ and the solvent removed. Finally, the crude product was purified by recrystallization from hexane/dichloromethane to afford the desired product.

2-{3-[1,3-Benzothiazol-2-yl]-1-[(trifluoromethyl)sulfonyl]-1,4-dihydropyridin-4-yl}-2-methylpropanoic acid (5a). White powder (389 mg, 90%), mp 198-199 °C. ¹H NMR (300 MHz, Acetone-*d*₆, ppm): δ 10.94 (s, 1H, OH), 8.04 (d, *J* 7.7 Hz, 1H, H-3), 7.97 (d, *J* 7.7 Hz, 1H, H-6), 7.58-7.42 (m, 3H, H-1, H-2, H-11), 6.94 (d, *J* 7.7 Hz, 1H, H-13), 5.63 (dd, *J* 7.9, 5.6 Hz, 1H, H-14), 4.50 (d, *J* 5.6 Hz, 1H, H-15), 1.19 (s, 3H, CH₃), 1.08 (s, 3H, CH₃). ¹³C NMR (75 MHz, CDCl₃, ppm): δ 176.2 (C=O), 166.1 (C-8), 153.2 (C-5), 134.3 (C-4), 126.8 (C-11), 126.5 (C-1), 125.8 (C-2), 123.2 (C-13), 123.0 (C-6), 121.8 (C-3), 119.4 (q, CF₃, *J* 320 Hz), 118.8 (C-10), 112.1 (C-14), 47.8 (C-16), 41.3 (C-15), 23.7 (CH₃), 19.5 (CH₃). IR (ATR, cm⁻¹): ν_{max} 1679 (C=O). MS (DART, *m/z*): 433 [M+H]⁺. HRMS (ESI-TOF): calcd for C₁₇H₁₆F₃N₂O₄S₂ [M+H]⁺ 433.05036; found 433.05072.

1-{3-[1,3-Benzothiazol-2-yl]-1-[(trifluoromethyl)sulfonyl]-1,4-dihydropyridin-4-yl}cyclobutane-1-carboxylic acid (5b). White powder (391 mg, 88%), mp 184-185 °C. ¹H NMR (300 MHz, DMSO-*d*₆, ppm): δ 12.63 (s, 1H, -OH), 8.11 (d, *J* 7.7 Hz, 1H, H-3), 8.01 (d, *J* 7.3 Hz, 1H, H-6), 7.59-7.42 (m, 3H, H-1, H-2, H-11), 6.90 (d, *J* 7.9 Hz, 1H, H-13), 5.51 (dd, *J* 8.0, 5.2 Hz, 1H, H-14), 4.28 (d, *J* 5.2 Hz, 1H, H-15), 2.11-1.93 (m, 4H, CH₂ cyclobutyl), 1.83-1.52 (m, 2H, CH₂ cyclobutyl). ¹³C NMR (75 MHz, DMSO-*d*₆, ppm): δ 176.7 (C=O), 166.3 (C-8), 153.1 (C-5), 134.2 (C-4), 127.1 (C-11), 126.2 (C-1), 125.8 (C-2), 123.2 (C-13), 122.7 (C-6 and C-3), 119.4 (q, CF₃, *J* 325 Hz), 118.8 (C-10), 112.1 (C-14), 53.7 (C-16), 39.7 (C-15), 28.7 (CH₂ cyclobutyl), 27.9 (CH₂ cyclobutyl), 16.3 (CH₂ cyclobutyl). IR (ATR, cm⁻¹): ν_{max} 1689 (C=O). MS (DART, *m/z*): 445.0 [M+H]⁺. HRMS (ESI-TOF): calcd for C₁₈H₁₆F₃N₂O₄S₂ [M+H]⁺ 445.05036; found 445.05024.

1-{1,3-Benzothiazol-2-yl}-1-[(trifluoromethyl)sulfonyl]-1,4-dihydropyridin-4-yl}cyclopentane-1-carboxylic acid (5c). White powder (426 mg, 93%), mp 193-194 °C. ¹H NMR (300 MHz, DMSO-*d*₆, ppm): δ 12.48 (s, 1H, -

OH) 8.07 (d, *J* 8.1 Hz, 1H, H-3), 7.98 (d, *J* 7.8 Hz, 1H, H-6), 7.54-7.41 (m, 3H, H-1, H-2), 7.34 (s, 1H, H-11), 6.68 (d, *J* = 7.9 Hz, 1H, H-13), 5.54 (dd, *J* 7.9, 5.4 Hz, 1H, H-14), 4.32 (d, *J* 5.3 Hz, 1H, H-15), 1.93-1.83 (m, 2H, CH₂ cyclopentyl), 1.45-1.35 (m, 6H, CH₂ cyclopentyl). ¹³C NMR (75 MHz, DMSO-*d*₆, ppm): δ 176.7 (C=O), 166.6 (C-8), 153.1 (C-5), 134.2 (C-4), 127.1 (C-11), 126.5 (C-1), 126.2 (C-2), 123.2 (C-13), 122.8 (C-6), 122.6 (C-3), 119.5 (C-10), 119.3 (q, CF₃, *J* 325 Hz), 113.3 (C-14), 60.8 (C-16), 40.3 (C-15), 33.8 (CH₂ cyclopentyl), 31.8 (CH₂ cyclopentyl), 24.1 (2 X CH₂ cyclopentyl). IR (ATR, cm⁻¹): ν_{max} 1683 (C=O). MS (DART, *m/z*): 459.0 [M+H]⁺. HRMS (ESI-TOF): calcd for C₁₉H₁₈F₃N₂O₄S₂ [M+H]⁺ 459.06601; found 459.06658.

{3-{1,3-Benzothiazol-2-yl}-1-[(trifluoromethyl)sulfonyl]-1,4-dihydropyridin-4-yl}cyclohexane-1-carboxylic acid (5d). White powder (373 mg, 79%), mp 216-217 °C. ¹H NMR (300 MHz, DMSO-*d*₆, ppm): δ 12.47 (s, 1H, OH), 8.06 (d, *J* 7.7 Hz, 1H, H-6), 7.97 (d, *J* 7.4 Hz, 1H, H-3), 7.56-7.35 (m, 3H, H-1, H-2, H-11), 6.95 (d, *J* 7.7 Hz, 1H, H-13), 5.50 (dd, *J* 7.8, 5.8 Hz, 1H, H-14), 4.12 (d, *J* 5.8 Hz, 1H, H-15), 1.51-1.38 (m, 2H, CH₂ cyclohexyl), 1.26-0.90 (m, 8H, CH₂ cyclohexyl). ¹³C NMR (75 MHz, DMSO-*d*₆, ppm): δ 175.4 (C=O), 166.6 (C-8), 153.1 (C-5), 134.3 (C-4), 126.9 (C-11), 126.7 (C-1), 126.1 (C-2), 123.5 (C-13), 123.2 (C-6), 122.6 (C-3), 119.3 (q, CF₃, *J* 325 Hz), 118.8 (C-10), 112.2 (C-14), 53.4 (C-16), 42.7 (C-15), 31.4 (CH₂ cyclohexyl), 29.5 (CH₂ cyclohexyl), 25.4 (CH₂ cyclohexyl), 23.5 (CH₂ cyclohexyl), 23.3 (CH₂ cyclohexyl). IR (ATR, cm⁻¹): ν_{max} 1675 (C=O). MS (DART, *m/z*): 473.0 [M+H]⁺. HRMS (ESI-TOF, *m/z*): calcd for C₂₀H₂₀F₃N₂O₄S₂ [M+H]⁺ 473.08166; found 473.06591.

General procedure for the synthesis of ̢-bromolactones 6a-d. A 100 mL round-bottom flask was charged with the corresponding carboxylic acid **5a-d** (0.5 mmol), sodium bicarbonate (50 mg, 0.6 mmol), *N*-bromosuccinimide (98 mg, 0.55 mmol), and tetrabutylammonium bromide (32 mg, 0.1 mmol) in dry dichloromethane (20 mL). The mixture was stirred at 0 °C and kept away from light. The reaction progress was monitored by TLC and once it reached completion, the crude mixture was washed with water (3 × 20 mL). The organic phase was dried over sodium sulfate and evaporated under reduced pressure. The obtained lactones were purified by silica gel column chromatography, eluted with a mixture of hexane/ethyl acetate.

6-(1,3-Benzothiazol-2-yl)-9-bromo-4,4-dimethyl-8-[(trifluoromethyl)sulfonyl]-2-oxa-8-azabicyclo[3.3.1]non-6-en-3-one (6a). White powder (224 mg, 88%), mp 218-219 °C. ¹H NMR (300 MHz, CDCl₃, ppm): δ 8.0 (d, *J* 8.0 Hz, 1H, H-6), 7.88 (d, *J* 7.9 Hz, 1H, H-3), 7.57-7.37 (m, 3H, H-1, H-2, H-11), 6.18 (s, 1H, H-13), 5.14 (t, *J* 3.1 Hz, 1H, H-14), 3.98 (t, *J* 2.4 Hz, 1H, H-15), 1.63 (s, 3H, -CH₃), 1.31 (s, 3H, -CH₃). ¹³C NMR (75 MHz, CDCl₃, ppm): δ 171.5 (C=O), 164.9 (C-8), 153.2 (C-5), 133.9 (C-4), 126.6 (C-11), 126.1 (C-1), 123.3 (C-2), 123.1 (C-6), 121.6 (C-3), 119.2 (q, CF₃, *J* 321 Hz), 117.28 (C-10), 82.94 (C-13), 45.65 (C-14), 43.87 (C-16), 35.76 (C-15), 28.08 (CH₃), 25.9 (CH₃). IR (ATR, cm⁻¹): ν_{max} 1761 (C=O). MS (DART, *m/z*): 513 (Br, 81) [M+H+2]⁺, 511 (Br, 79) [M+H]⁺. HRMS (ESI-TOF): calcd for C₁₇H₁₅BrF₃N₂O₄S₂ [M+H]⁺ 510.96087; found 510.96142.

8-(1,3-Benzothiazol-2-yl)-9-bromo-6-[(trifluoromethyl)sulfonyl]-4-oxa-6-azaspiro[bicyclo[3.3.1]nonane-2,1'-cyclobutan]-7-en-3-one (6b). White powder (235 mg, 90%), mp 168-169 °C. ¹H NMR (300 MHz, DMSO-*d*₆, ppm): δ 8.11 (d, *J* 7.1 Hz, 1H, H-6), 8.04 (d, *J* 7.9 Hz, 1H, H-6), 7.61-7.40 (m, 3H, H-1, H-2, H-11), 6.43 (br, 1H, H-13), 5.31 (t, *J* 3.1 Hz, 1H, H-14), 4.24 (t, *J* 2.4 Hz, 1H, H-15), 2.77-2.67 (m, 1H, CH₂ cyclobutyl), 2.29-2.18 (m, 4H, CH₂ cyclobutyl), 1.98-1.93 (m, 3H, CH₂ cyclobutyl). ¹³C NMR (75 MHz, DMSO-*d*₆, ppm): δ 171.2 (C=O), 165.3 (C-8), 153.0 (C-5), 134.2 (C-4), 127.4 (C-11), 126.6 (C-1), 123.3 (C-2), 123.1 (C-6), 122.8 (C-3), 119.4 (q, CF₃, *J* 324 Hz), 117.6 (C-10), 83.2 (C-13), 49.9 (C-14), 43.0 (C-16), 36.8 (C-15), 34.14 (CH₂ cyclobutyl), 27.3 (CH₂ cyclobutyl), 15.7 (CH₂ cyclobutyl). IR (ATR, cm⁻¹): ν 1756 (C=O). MS (DART, *m/z*): 525 (Br, 81), 523 (Br, 79) [M+H]⁺. HRMS (ESI-TOF, *m/z*): calcd for C₁₈H₁₅BrF₃N₂O₄S₂ [M+H]⁺ 522.96087; found 522.98093.

8-(1,3-Benzothiazol-2-yl)-9-bromo-6-[(trifluoromethyl)sulfonyl]-4-oxa-6-azaspiro[bicyclo[3.3.1]nonane-2,1'-cyclopentan]-7-en-3-one (6c). White powder (212 mg, 79%), mp 192-193 °C. ¹H NMR (300 MHz, CDCl₃, ppm): δ 7.98 (d, *J* 8.1 Hz, 1H, H-6), 7.86 (d, *J* 7.9 Hz, 1H, H-3), 7.52-7.36 (m, 3H, H-1, H-2, H-11), 6.15 (br, 1H, H-13), 5.05 (br, 1H, H-14), 3.95 (br, 1H, H-15), 2.18-1.99 (m, 3H, CH₂ cyclopentyl), 1.87-1.65 (m, 5H, CH₂ cyclopentyl).

^{13}C NMR (75 MHz, CDCl_3 , ppm): δ 171.9 (C=O), 165.1 (C-8), 153.2 (C-5), 134.0 (C-4), 126.6 (C-11), 126.0 (C-1), 123.6 (C-2), 123.3 (C-6), 121.6 (C-3), 119.3 (q, CF_3 , J 325 Hz), 117.1 (C-10), 82.9 (C-13), 56.5 (C-14), 43.9 (C-16), 40.7 (C-15), 36.8 (CH_2 cyclopentyl), 35.6 (CH_2 cyclopentyl), 25.4 (CH_2 cyclopentyl), 24.9 (CH_2 cyclopentyl). IR (ATR, cm^{-1}): ν_{max} 1760 (C=O). MS (DART, m/z): 539 (Br, 81) $[\text{M}+\text{H}]^+$, 537 (Br, 79) $[\text{M}+\text{H}]^+$. HRMS (ESI-TOF): calcd for $\text{C}_{19}\text{H}_{17}\text{BrF}_3\text{N}_2\text{O}_4\text{S}_2$ $[\text{M}+\text{H}]^+$ 536.97652; found 536.97534.

8-(1,3-Benzothiazol-2-yl)-9-bromo-6-[(trifluoromethyl)sulfonyl]-4-oxa-6-azaspiro[bicyclo[3.3.1]nonane-2,1'-cyclohexan]-7-en-3-one (6d). White powder (179 mg, 65%), mp 210-211 °C. ^1H NMR (300 MHz, CDCl_3 , ppm): δ 8.02 (d, 1H, H-6, J 7.6 Hz), 7.87 (d, 1H, H-3, J 7.2 Hz), 7.53-7.40 (m, 2H, H-1, H-2), 7.32 (s, 1H, H-11), 6.11 (br, 1H, H-13), 5.24 (t, 1H, H-14, J 2.4 Hz), 4.09 (t, 1H, H-15, J 2.4 Hz), 2.11-1.33 (m, 10H, CH_2 cyclohexyl). ^{13}C NMR (75 MHz, CDCl_3 , ppm): δ 170.7 (C=O), 165.1 (C-8), 153.2 (C-5), 134.0 (C-4), 126.6 (C-11), 126.0 (C-1), 123.9 (C-2), 123.3 (C-6), 121.6 (C-3), 119.2 (q, CF_3 , J 321 Hz), 117.1 (C-10), 82.8 (C-13), 49.2 (C-14), 43.4 (C-19), 35.6 (C-15), 35.4 (CH_2 cyclohexyl), 33.4 (CH_2 cyclohexyl), 24.9 (CH_2 cyclohexyl), 21.5 (CH_2 cyclohexyl), 21.3 (CH_2 cyclohexyl). IR (ATR, cm^{-1}): ν_{max} 1759 (C=O). MS (DART, m/z): 553 (Br, 81) $[\text{M}+\text{H}+2]^+$, 551 (Br, 79) $[\text{M}+\text{H}]^+$. HRMS (ESI-TOF): calcd for $\text{C}_{20}\text{H}_{19}\text{BrF}_3\text{N}_2\text{O}_4\text{S}_2$ $[\text{M}+\text{H}]^+$ 550.99217; found 550.99174.

Cell culture and assay for cytotoxicity

Cell lines HCT-15, MCF-7, K-562, U-251, PC-3, SKLU-1 and COS7 were supplied by The National Cancer Institute (NCI), U.S.A. and HIV/AIDS Services Center of Mexico City. The cytotoxicity of tumor cells with the test compounds was determined using the protein-binding dye sulforhodamine B (SRB) in microculture assay to measure cell growth. The cell lines were cultured in RPMI-1640 (Sigma Chemical Co., Ltd., St. Louis, MO, U.S.A.), supplemented with 10% fetal bovine serum which was purchased from Invitrogen Corporation, 2 mM L-glutamine, 100 IU/ml penicillin G, 100 mg/ml streptomycin sulfate, and 0.25 mg/ml amphotericin B (Gibco). They were maintained at 37 °C in a 5% CO_2 atmosphere with 95% humidity. For the assay, 5×10^4 cell/ml (K562, MCF-7), 7.5×10^4 cell/ml (U251, PC-3) and 10×10^4 cell/ml (HCT-15, MT2), and 100 ml/well of these cells suspension was seeded in 96-well microtiter plates and incubated to allow for cell attachment. After 24 h, 100 ml of each test compounds and positive substances (Cisplatin or Gefitinib) were added to each well. 48 h Later, adherent cell cultures were fixed *in situ* by adding 50 ml of cold 50% (wt/vol) trichloroacetic acid (TCA) and incubated for 60 min at 4 °C. The supernatant was discarded, and the plates were washed three times with water and air-dried. Cultures fixed with TCA were stained for 30 min with 100 ml of 0.4% SRB solution. Protein-bound dye was extracted with 10 mM unbuffered tris base and the optical densities were read on an Ultra Microplate Reader (Elx 808, BIO-TEK Instruments, Inc.), with a test wavelength of 515 nm.

Acknowledgements

This work was financially supported by DGAPA PAPIIT Project IN202917 and CONACYT Project 252020. We thank Rocio Patiño, M. Del Carmen García, Luis Velasco and Javier Perez for technical support with the IR spectroscopy and mass spectrometry.

Supplementary Material

Copies of ^1H and ^{13}C NMR spectra of the new compounds. CCDC 1898179 for **5b** contain the supplementary crystallographic data for this paper.

References

1. Pozharskii, A. F.; Soldatenkov, A.; Katritzky, A. R. *Heterocycles in Life and Society: An Introduction to Heterocyclic Chemistry, Biochemistry, and Applications* 2nd Ed. Wiley:New York, 2011.
2. Keri, R.S.; Patil, M.R.; Patil, S.A.; Budagumpi, S. *Eur. J. Med. Chem.* **2015**, *89*, 207-251.
<https://doi.org/10.1016/j.ejmech.2014.10.059>
3. Huang, S.T.; Hsei, I.J.; Chen, C. *Bioorg. Med. Chem.* **2006**, *14*, 6106-6119.
<https://doi.org/10.1016/j.bmc.2006.05.007>
4. Singh, M.; Singh, S.K.; Gangwar, M.; Nath, G.; Singh, S.K. *RSC Adv.* **2014**, *4*, 19013-19023.
<https://doi.org/10.1039/C4RA02649G>
5. Siddiqui, N.; Pandeya, S.N.; Khau, S.A.; Stables, J.; Rana, A.; Alam, M.; Md. Arshad, F.; Bhat, M.A. *Bioorg. Med. Chem.* **2007**, *17*, 255-259.
<https://doi.org/10.1016/j.bmcl.2006.09.053>
6. Akhtar, T.; Hameed, S.; Al-Masoudi, N.; Loddo, R.; Colla, P. *Acta Pharm.* **2008**, *58*, 135-149.
<https://doi.org/10.2478/v10007-008-0007-2>
7. Suresh, C.H.; Rao, J.V.; Jayaveera, K.N.; Subudhi, S.K. *Int. J. Pharma.* **2013**, *2*, 257-261.
Herrera-Cano, N.; Ballari, M.S.; López, A.G.; Santiago, A.N. *J. Agric. Food Chem.* **2015**, *63*, 3681-3686.
<https://doi.org/10.1021/acs.jafc.5b00150>
8. Siddiqui, N.; Alam, M.; Siddiqui, A.A. *Asian J. Chem.* **2004**, *16*, 1005-1008.
9. Patel, P.; Pillai, J.; Darji, N.; Patel, B. *Int. J. Drug Res. Tech.* **2012**, *2*, 170-176.
10. Pattan, S.R.; Suresh, C.; Pujar, V.D.; Reddy, V.V.K.; Rasal, V.P.; Koti, B.C. *Ind. J. Chem.* **2005**, *44B*, 2404-2408.
11. Gill, R. K.; Rawal, R.K.; Bariwal, J. *Arch. Pharm. Chem. Life Sci.* **2015**, *348*, 155-178.
<https://doi.org/10.1002/ardp.201400340>
12. Kiuru, P.; Yli-Kauhalauma, J. *Pyridine and Its Derivatives*, in Majumdar, K.; Chattopadhyay, S. K., *Heterocycles in Natural Product Synthesis*. 2011, Eds.; Wiley-VCH Verlag: Weinheim, pp 267-297.
13. Altaf, A.A.; Shahzad, A.; Gul, Z.; Rasool, N.; Badshah, A.; Lal, B.; Khan, E. *J. Drug Des. Med. Chem.* **2015**, *1*, 1-11.
14. Henry, G.D. *Tetrahedron* **2004**, *60*, 6043-6061.
<https://doi.org/10.1021/ja303002a>
15. Hill, M.D. *Chem. Eur. J.* **2010**, *16*, 12052-12062.
<https://doi.org/10.1002/chem.201001100>
16. Ioan, P.; Carosati, E.; Micucci, M.; Cruciani, G.; Broccatelli, F.; Zhorov, B.S.; Chiarini, A.; Budriesi, R. *Curr. Med. Chem.* **2011**, *18*, 4901-4922.
<https://doi.org/10.2174/092986711797535173>
17. Carosati, E.; Ioan, P.; Micucci, M.; Broccatelli, F.; Cruciani, G.; Zhorov, B.S.; Chiarini, A.; Budriesi, R. *Curr. Med. Chem.* **2012**, *19*, 4306-4323.
<https://doi.org/10.2174/092986712802884204>
18. Zarrin, A.; Mehdipour, A.R.; Miri, R. *Chem. Biol. Drug Des.* **2010**, *76*, 369-381.
<https://doi.org/10.1111/j.1747-0285.2010.01025.x>
19. Bull, J.A.; Mousseau, J.J.; Pelletier, G.; Charette, A.B. *Chem. Rev.* **2012**, *112*, 2642-2713.
<https://doi.org/10.1021/cr200251d>
20. Álvarez-Toledano, C.; Penieres-Carrillo J.G. *Eur. J. Org. Chem.* **2018**, 4040-4049.
<https://doi.org/10.1002/ejoc.201800227>

21. Garduño-Alva, A.; Xu, Y.; Gualo-Soberanes, N.; López-Cortés, J.; Rudler, H.; Parlier, A.; Ortega-Alfaro, M.C.; Álvarez-Toledano, C.; Toscano, R. A. *Eur. J. Org. Chem.* **2008**, 3714–3723.
<https://doi.org/10.1002/ejoc.200800338>
22. Rivera-Hernández, A.; Chans, G.M.; Rudler, H.; López-Cortés, J.G.; Toscano, R.A.; Álvarez-Toledano, C.; *Tetrahedron* **2014**, 70, 1861–1871.
<https://doi.org/10.1016/j.tet.2014.01.044>
23. Rivera-Hernández, A.; López-Jimeno, I. S.; Carmona-Reyes, G. A.; Toscano, R.A.; Penieres-Carrillo, J.G.; Álvarez-Toledano, C. *Tetrahedron Lett.* **2015**, 56, 4829–4832.
<https://doi.org/10.1016/j.tetlet.2015.06.069>
24. Valencia-Galicia, N. A.; Corona-Sánchez, R.; Ballinas-Indili, R.; Toscano, R.A.; Macías-Rubalcava, M. L.; Álvarez-Toledano, C. *Tetrahedron Lett.* **2017**, 58, 3168–3171.
<https://doi.org/10.1016/j.tetlet.2016.10.013>
25. Itoh, T.; Miyazaki, M.; Nagata, K.; Ohsawa, A. *Tetrahedron*, **2000**, 56, 4383–4395.
[https://doi.org/10.1016/S0040-4020\(00\)00331-8](https://doi.org/10.1016/S0040-4020(00)00331-8)
26. Rudler, H.; Denise, B.; Xu, Y.; Parlier, A.; Vaissermann, J. *Eur. J. Org. Chem.* **2005**, 3724–3744.
<https://doi.org/10.1002/ejoc.200500162>
27. Monks, A.; Scudeiro, D.; Skehan, P.; Shoemaker, R.; Paull, K.; Vistica, D.; Hose, C.; Langley, J.; Cronise, P.; Vaigro-Wolff, A.; GrayGoodrich, M.; Campbell, H.; Mayo, J.; Boyd, M. *J. Natl. Cancer Inst.* **1991**, 83, 757–766.
28. Noolvi, M. N.; Patel, H. M.; Kaur, M.; *Eur. J. Med. Chem.* **2012**, 54, 447–462.
<https://doi.org/10.1016/j.ejmech.2012.05.028>
29. Ainsworth, C.; Kuo, Y. N. *J. Organomet. Chem.* **1972**, 46, 73–84.
[https://doi.org/10.1016/S0022-328X\(00\)90476-5](https://doi.org/10.1016/S0022-328X(00)90476-5)
30. Huang, J.; Chan, J.; Chen, Y.; Borths, C. J.; Baucom, K. D.; Larsen, R. D.; Faul, M.M. *J. Am. Chem. Soc.*, 2010, 132, 3674–3675.
<https://doi.org/10.1021/ja100354j>