

Supplementary Material

Efficient synthesis of novel 1-hydroxy-2,4,5-trisubstituted imidazole derivatives via a one-pot, four-component reaction between hydroxylamine, benzonitriles, arylglyoxals and 1,3-dicarbonyl compounds

Farzaneh Alizadeh-Bami, Mina Salehzadeh, Hossein Mehrabi,* and Reza Ranjbar-Karimi

Department of Chemistry, Vali-e-Asr University of Rafsanjan, 77176 Rafsanjan, Iran

E-mail: mehraby_h@yahoo.com

Table of Contents

1. Experimental Section	S2
2. ^1H NMR and ^{13}C NMR data for compounds 5a-I	S3

Experimental Section

General. All chemicals were purchased from Aldrich and Merck with high-grade quality, and used without any purification. All melting points were obtained by Barnstead Electrothermal 9200 apparatus and are uncorrected. The reactions were monitored by TLC and all yields refer to isolated products. NMR spectra were obtained on a Varian 500 MHz spectrometer (¹H NMR at 500 MHz, ¹³C NMR at 125 MHz) in DMSO using TMS as an internal standard. Infrared spectra were recorded on a Bruker FT-IR Equinax-55 spectrophotometer in KBr with absorption in cm⁻¹. Elemental analyses were performed using a Carlo Erba EA 1108 instrument. All products were characterized by their spectral and physical data.

General procedure for the synthesis of 1-hydroxy-2,4,5-trisubstituted imidazole derivatives.

A mixture of hydroxylamine (**1**) (2.0 mmol) and benzonitrile (**2**) (1.0 mmol) was stirred in 10 mL of ethanol at reflux for 3 hr to give amidoxime. Then, 1,3-dicarbonyl compound (**3**) (1.0 mmol) and arylglyoxal (**4**) (1.0 mmol) was added, and obtain mixture was stirred at reflux for 1 hr. After completion of the reaction, determined by TLC, the solvent was removed under reduced pressure, and the resulting crude product was purified by recrystallization from ethanol to give the pure compounds **5a–I** (65–87%).

5-(5-(4-Chlorophenyl)-1-hydroxy-2-phenyl-1*H*-imidazol-4-yl)-6-hydroxy-2,2-dimethyl-4*H*-1,3-dioxin-4-one

(**5a**). mp 213–215 °C. IR ν/cm^{-1} (KBr): 3424, 2991, 1639. ¹H NMR (500 MHz, DMSO): δ 1.64 (s, 6H, 2CH₃), 2.06 (s, 1H, OH), 7.51 (d, *J* 10.0 Hz, 2H, ArH), 7.55 (s, 2H, ArH), 7.56 (d, *J* 5.0 Hz, 2H, ArH), 7.63 (s, 1H, ArH), 8.04 (m, 2H, ArH), 14.10 (s, 1H, OH) ppm. ¹³C NMR (125 MHz, DMSO): δ 26.2, 102.23, 122.3, 125.4, 126.5, 128.5, 129.1, 129.3, 129.8, 131.5, 131.5, 131.8, 135.9, 161.8, 165.3, 196.5 ppm. Anal. Calcd for C₂₁H₁₇ClN₂O₅ (412.83): C, 61.10; H, 4.15; N, 6.79. Found: C, 61.24; H, 4.17; N, 6.75%.

5-(2-Benzyl-1-hydroxy-5-(p-tolyl)-1*H*-imidazol-4-yl)-6-hydroxy-2,2-dimethyl-4*H*-1,3-dioxin-4-one (**5b**). mp 208–211 °C. IR ν/cm^{-1} (KBr): 3432, 2998, 1667; ¹H NMR (500 MHz, DMSO): δ 1.22 (s, 6H, 2CH₃), 1.58 (s, 3H, CH₃), 2.07 (s, 1H, OH), 3.76 (s, 2H, CH₂), 6.67 (d, *J* 10.0 Hz, 2H, ArH), 6.78 (d, *J* 10.0 Hz, 1H, ArH), 6.82 (s, 2H, ArH), 6.84 (d, *J* 7.9 Hz, 2H, ArH), 6.92 (d, 2H, *J* 7.9 Hz, ArH), 13.31 (s, 1H, OH) ppm. ¹³C NMR (125 MHz, DMSO): δ 20.8, 25.6, 28.9, 62.6, 101.7, 123.2, 123.7, 125.8, 126.1, 127.1, 128.3, 128.5, 128.6, 133.6, 136.1, 137.6, 165.1, 191.2 ppm. Anal. Calcd for C₂₃H₂₂N₂O₅ (406.44): C, 67.97; H, 5.46; N, 6.89. Found: C, 67.19; H, 5.50; N, 6.91%.

5-(2-(2-Chlorophenyl)-5-(4-chlorophenyl)-1-hydroxy-1*H*-imidazol-4-yl)-6-hydroxy-2,2-dimethyl-4*H*-1,3-dioxin-4-one (**5c**). mp 204–206 °C. IR ν/cm^{-1} (KBr): 3420, 2987, 1609; ¹H NMR (500 MHz, DMSO): δ 1.05 (s, 1H, OH), 1.65 (s, 6H, 2CH₃), 7.55 (d, *J* 5.0 Hz, 2H, ArH), 7.64 (d, *J* 5.0 Hz, 2H, ArH), 7.71 (t, 2H, ArH), 7.78 (d, *J* 6.9 Hz, 1H, ArH), 7.87 (d, *J* 7.7 Hz, 1H, ArH), 13.85 (s, 1H, OH) ppm. ¹³C NMR (125 MHz, DMSO): δ 26.3, 69.2, 102.0, 111.8, 115.6, 122.4, 128.1, 128.4, 128.8, 129.0, 130.5, 133.6, 134.0, 134.7, 135.1, 144.2, 164.8, 214.0 ppm. Anal. Calcd for C₂₁H₁₆Cl₂N₂O₅ (447.27): C, 56.39; H, 3.61; N, 6.26%.

5-(5-(4-Chlorophenyl)-1-hydroxy-2-(4-methoxyphenyl)-1*H*-imidazol-4-yl)-6-hydroxy-2,2-dimethyl-4*H*-1,3-dioxin-4-one (**5d**). mp 203–205 °C. IR ν/cm^{-1} (KBr): 3422, 2938, 1610. ¹H NMR (500 MHz, DMSO): δ 1.61 (s, 6H, 2CH₃), 2.08 (s, 1H, OH), 3.86 (s, 3H, CH₃), 7.20 (d, *J* 9.0 Hz, 2H, ArH), 7.54 (d, *J* 8.6 Hz, 2H, ArH), 7.69 (d, *J* 8.6 Hz, 2H, ArH), 8.04 (d, *J* 9.0 Hz, 2H, ArH), 13.51 (s, 1H, OH) ppm. ¹³C NMR (125 MHz, DMSO): δ 26.3, 56.0, 68.1, 101.7, 111.3, 115.0, 126.6, 128.4, 128.6, 128.8, 129.2, 130.4, 133.2, 136.9, 141.1, 164.9, 217.0 ppm. Anal. Calcd for C₂₂H₁₉ClN₂O₆ (442.85): C, 59.67; H, 4.32; N, 6.33. Found: C, 59.79; H, 4.33; N, 6.28%.

5-(5-(4-Bromophenyl)-1-hydroxy-2-phenyl-1*H*-imidazol-4-yl)-6-hydroxy-2,2-dimethyl-4*H*-1,3-dioxin-4-one (5e). mp 218-220 °C. IR ν/cm^{-1} (KBr): 3426, 2986, 1639. ^1H NMR (500 MHz, DMSO): δ 1.63 (s, 6H, 2CH₃), 2.08 (s, 1H, OH), 7.55 (d, *J* 5.0 Hz, 2H, ArH), 7.64 (d, *J* 5.0 Hz, 2H, ArH), 7.68 (s, 1H, ArH), 7.70 (d, *J* 10.0 Hz, 2H, ArH), 8.07 (d, *J* 10.0 Hz, 2H, ArH), 13.75 (s, 1H, OH) ppm. ^{13}C NMR (125 MHz, DMSO): δ 26.3, 78.7, 101.8, 127.4, 128.7, 128.9, 129.3, 129.5, 131.4, 131.7, 132.8, 133.9, 136.7, 140.7, 164.9, 187.4 ppm. Anal. Calcd for C₂₁H₁₇BrN₂O₅ (457.28): C, 55.16; H, 3.75; N, 6.13. Found: C, 55.24; H, 3.77; N, 6.08%.

2-(5-(4-Chlorophenyl)-1-hydroxy-2-phenyl-1*H*-imidazol-4-yl)-3-hydroxy-5,5-dimethylcyclohex-2-en-1-one (5f). mp 205-207 °C. IR ν/cm^{-1} (KBr): 3449, 2954, 1694. ^1H NMR (500 MHz, DMSO): δ 1.09 (s, 6H, 2CH₃), 2.59 (s, 1H, OH), 3.17 (s, 2H, CH₂), 3.32 (d, 2H, CH₂), 7.55 (d, *J* 5.0 Hz, 2H, ArH), 7.57 (s, 1H, ArH), 7.61 (d, *J* 10.0 Hz, 2H, ArH), 7.81 (d, *J* 5.0 Hz, 2H, ArH), 8.43 (d, *J* 10.0 Hz, 2H, ArH), 14.01 (s, 1H, OH) ppm. ^{13}C NMR (125 MHz, DMSO): δ 28.1, 32.9, 45.7, 51.4, 120.7, 129.2, 129.4, 129.6, 131.3, 132.9, 133.5, 136.0, 139.5, 164.4, 165.5, 173.5, 192.2, 196.7 ppm. Anal. Calcd for C₂₃H₂₁ClN₂O₃ (408.88): C, 67.56; H, 5.18; N, 6.58. Found: C, 67.72; H, 5.23; N, 6.56%.

3-Hydroxy-2-(1-hydroxy-2,5-diphenyl-1*H*-imidazol-4-yl)-5,5-dimethylcyclohex-2-en-1-one (5g). mp 188-190 °C. IR ν/cm^{-1} (KBr): 3362, 2952, 1690. ^1H NMR (500 MHz, DMSO): δ 1.09 (s, 6H, 2CH₃), 2.49 (s, 1H, OH), 2.59 (s, 2H, CH₂), 3.18 (s, 2H, CH₂), 7.53 (d, *J* 10.0 Hz, 2H, ArH), 7.54 (d, *J* 10.0 Hz, 2H, ArH), 7.59 (m, 1H, ArH), 7.67 (m, 1H, ArH), 7.78 (d, *J* 5.0 Hz, 2H, ArH), 8.41 (d, *J* 10.0 Hz, 2H, ArH), 14.10 (s, 1H, OH) ppm. ^{13}C NMR (125 MHz, DMSO): δ 28.1, 32.8, 45.7, 51.5, 120.7, 129.1, 129.2, 129.4, 129.5, 132.8, 134.5, 134.8, 136.1, 164.9, 165.5, 173.4, 193.2, 196.6 ppm. Anal. Calcd for C₂₃H₂₂N₂O₃ (374.44): C, 73.78; H, 5.92; N, 7.48. Found: C, 73.66; H, 5.91; N, 7.44%.

3-Hydroxy-2-(1-hydroxy-2-phenyl-5-(p-tolyl)-1*H*-imidazol-4-yl)-5,5-dimethylcyclohex-2-en-1-one (5h). mp 150-153 °C. IR ν/cm^{-1} (KBr): 3450, 2958, 1682. ^1H NMR (500 MHz, DMSO): δ 1.17 (s, 6H, 2CH₃), 2.05 (s, 1H, OH), 2.42 (s, 3H, CH₃), 2.55 (s, 2H, CH₂), 3.14 (s, 2H, CH₂), 7.26 (s, 2H, ArH), 7.50 (m, 2H, ArH), 7.52 (d, *J* 8.3 Hz, 1H, ArH), 7.72 (d, *J* 7.0 Hz, 2H, ArH), 8.56 (d, *J* 7.0 Hz, 2H, ArH), 13.81 (s, 1H, OH) ppm. ^{13}C NMR (125 MHz, DMSO): δ 21.8, 28.3, 32.7, 46.3, 52.0, 117.2, 120.4, 128.6, 129.4, 129.5, 132.1, 136.1, 138.9, 144.7, 148.5, 165.6, 172.0, 195.5, 197.5 ppm. Anal. Calcd for C₂₄H₂₄N₂O₃ (388.47): C, 74.21; H, 6.23; N, 7.21. Found: C, 74.37; H, 6.29; N, 7.18%.

2-(2-(4-Bromophenyl)-5-(4-chlorophenyl)-1-hydroxy-1*H*-imidazol-4-yl)-3-hydroxy-5,5-dimethylcyclohex-2-en-1-one (5i). mp 192-194 °C. IR ν/cm^{-1} (KBr): 3425, 2956, 1654. ^1H NMR (500 MHz, DMSO): δ 0.79 (s, 3H, CH₃), 0.81 (s, 3H, CH₃), 2.11 (s, 2H, CH₂), 2.31 (s, 2H, CH₂), 2.66 (s, 1H, OH), 7.03 (d, *J* 8.7 Hz, 2H, ArH), 7.11 (d, *J* 8.6 Hz, 2H, ArH), 7.42 (d, *J* 8.7 Hz, 2H, ArH), 7.59 (d, *J* 8.6 Hz, 2H, ArH), 13.05 (s, 1H, OH) ppm. ^{13}C NMR (125 MHz, DMSO): δ 27.9, 28.0, 30.3, 51.7, 51.9, 104.0, 109.6, 125.1, 126.0, 127.7, 128.3, 129.2, 132.2, 134.1, 136.7, 156.3, 179.0, 199.2, 199.9 ppm.

2-(2-(4-Bromophenyl)-1-hydroxy-5-(p-tolyl)-1*H*-imidazol-4-yl)-3-hydroxy-5,5-dimethylcyclohex-2-en-1-one (5j). mp 193-195 °C. IR ν/cm^{-1} (KBr): 3432, 2954, 1654. ^1H NMR (500 MHz, DMSO): δ 0.84 (s, 3H, CH₃), 0.87 (s, 3H, CH₃), 2.12 (s, 3H, CH₃), 2.15 (s, 2H, CH₂), 2.37 (s, 2H, CH₂), 2.54 (s, 1H, OH), 6.92 (d, *J* 8.0 Hz, 2H, ArH), 7.10 (d, *J* 8.4 Hz, 2H, ArH), 7.48 (d, *J* 8.0 Hz, 2H, ArH), 7.66 (d, *J* 8.4 Hz, 2H, ArH), 13.15 (s, 1H, OH) ppm. ^{13}C NMR (125 MHz, DMSO): δ 21.0, 28.0, 28.1, 30.3, 51.8, 51.9, 104.6, 109.7, 124.3, 125.4, 127.5, 128.9, 129.1, 132.2, 134.9, 138.3, 155.8, 180.1, 199.1, 200.0 ppm. Anal. Calcd for C₂₄H₂₃BrN₂O₃ (467.36): C, 61.68; H, 4.96; N, 5.99. Found: C, 61.70; H, 4.99; N, 6.04%.

2-(2-(4-Bromophenyl)-1-hydroxy-5-phenyl-1*H*-imidazol-4-yl)-3-hydroxy-5,5-dimethylcyclohex-2-en-1-one (5k). mp 197-199 °C. IR ν/cm^{-1} (KBr): 3433, 2924, 1655. ^1H NMR (500 MHz, DMSO): δ 0.86 (s, 3H, CH₃), 0.90 (s, 3H, CH₃), 2.18 (s, 2H, CH₂), 2.40 (s, 1H, OH), 2.50 (s, 2H, CH₂), 7.15 (d, *J* 5.0 Hz, 2H, ArH), 7.25 (s, 2H, ArH), 7.52

(d, J 5.0 Hz, 2H, ArH), 7.55 (s, 1H, ArH), 7.69 (d, J 7.1 Hz, 2H, ArH), 13.18 (s, 1H, OH) ppm. ^{13}C NMR (125 MHz, DMSO): δ 28.0, 28.0, 30.4, 51.8, 51.9, 104.7, 117.0, 124.4, 127.7, 128.2, 128.5, 129.2, 130.7, 132.2, 132.2, 138.7, 156.0, 180.0, 199.1 ppm.

2-(2-(4-Bromophenyl)-1-hydroxy-5-(4-nitrophenyl)-1*H*-imidazol-4-yl)-3-hydroxy-5,5-dimethyl cyclohex-2-en-1-one (5I). mp 196-198 °C. IR ν/cm^{-1} (KBr): 3424, 2957, 1660. ^1H NMR (500 MHz, DMSO): δ 1.04 (s, 3H, CH_3), 1.07 (s, 3H, CH_3), 1.18 (s, 1H, OH), 2.35 (s, 2H, CH_2), 2.58 (s, 2H, CH_2), 7.61 (m, 2H, ArH), 7.68 (d, J 8.1 Hz, 2H, ArH), 7.83 (d, J 8.1 Hz, 2H, ArH), 8.18 (d, J 8.3 Hz, 2H, ArH), 13.32 (s, 1H, OH) ppm. ^{13}C NMR (125 MHz, DMSO): δ 28.1, 28.3, 30.6, 51.9, 52.1, 104.0, 109.8, 123.7, 124.9, 125.7, 128.4, 129.3, 132.5, 145.1, 148.0, 157.3, 178.5, 199.5, 200.4 ppm. Anal. Calcd for $\text{C}_{23}\text{H}_{20}\text{BrN}_3\text{O}_5$ (498.33): C, 55.44; H, 4.05; N, 8.43. Found: C, 55.27; H, 4.01; N, 8.37%.























