

Synthesis of 2-pyridyl oxazoline esters and their organocatalytic application to the enantioselective silane reduction of ketones

Zhenyu Cao, Nadine McCleary, and Patrick O'Leary*

National University of Ireland, Galway, School of Chemistry, SMACT (Synthetic Methods: Asymmetric Catalytic Transformations)

Email: patrick.oleary@nuigalway.ie


Received 10-29-2019

Accepted 12-16-2019

Published on line 01-15-2020

Abstract

The synthesis of a number of novel chiral 2-pyridyl oxazoline esters is reported along with the use of these in the enantioselective organocatalytic reduction of ketones. The chiral alcohols were prepared in good to excellent yields (up to 99%) and were found to catalyse the reduction reactions with good conversions (up to 99%) and in moderate to good enantioselectivities (up to 65% at high conversion and 70% with lower reactivity). The results give an insight into the effects of the catalyst structure on the selectivity observed in the reaction.



Keywords: Organic synthesis, ligand design, enantioselective, reduction, organocatalytic, oxazoline

Introduction

Asymmetric reductions of carbonyl and heterocarbonyl functional groups are fundamental organic transformations. Non stereoselective reduction of ketones and imines can easily be achieved with hydrides but the asymmetric reduction is a very active area of research¹ with new methods being reported regularly using a wide variety of reactions and reagents for example chiral ligand metal complexes in hydrogenation^{2,3} and transfer hydrogenation,^{4,5} enzymatic reduction⁶ and chiral ligand metal complexes in hydrosilylation.^{7,8}

Recently, the asymmetric hydrosilylation of ketones and imines using chiral Lewis base activated trichlorosilane emerged as an effective system capable of competing with the conventional metal-catalysed protocols.⁹ In 1999 and 2001 a major breakthrough in this area was reported by Matsumura in which encouraging enantioselectivities were achieved in the reduction of ketones (up to 51% ee) and imines (up to 66% ee).^{10,11} Since then, rapid development in this field has been seen with efficient catalysts reported to catalyse the reaction, including chiral formamides,¹²⁻¹⁴ picolinamides¹⁵⁻¹⁷ and sulfinamides.¹⁸

In 2006, Malkov reported the design and synthesis of a series of catalysts based on the pyridyl oxazoline core (Figure 1).¹⁹ These catalysts afforded good enantioselectivities in the hydrosilylation of both ketones (some in the 80-90% ee range) and imines (Some >80% ee), whereas other catalysts were more often designed to target only one of the reactions.




Figure 1. Some pyridyl oxazoline ligands described previously.¹⁹

Results and Discussion

Our group has an ongoing interest in the design and application of novel oxazoline based catalytic systems.²⁰⁻²² The initial aim of this project was to synthesize the pyridine substituted oxazoline **1**. After that, a variety of functionalisation reactions would be carried out on the alcohol to give us various ligands with which we could study the effect of variations of functionality in the ligands **2-4** (Figure 2) on their catalytic performance, as organocatalysts in asymmetric catalytic reduction reactions. The ultimate aim of this project is to maximize the catalytic performance of the pyridine-substituted oxazoline ligands and to study the structure-selectivity relationship for this catalyst type in these reductions.





Figure 2. Initial target compounds of this work.

We designed a synthetic pathway to the pyridinyl-oxazoline ligand using aspects of the synthesis of similar compounds described by Balavoine.²³ However we aimed to start from L-serine **5**, which was much cheaper than the diol which would have been the starting material were we to follow the planned method. Thus, we needed to use an extra synthetic step to reduce the oxazoline ester **6** to the alcohol **1**.




Reagents and conditions: (a) thionyl chloride, MeOH, 0 °C-rt, then reflux overnight, 99% (b) Na metal, MeOH, rt, 24 h (c) CH₂Cl₂, reflux, 2.5 h, 78% (d) LiAlH₄, Et₂O, 0 °C-rt, 4 h, 31% (e) NaBH₄, MeOH, 0 °C-rt, 4 h, 72% after crystallization.

Scheme 1. Synthesis of pyridinyl-oxazoline ligand **1**.

As shown in Scheme 1, the first step involved the protection of L-serine **5** as its methyl ester **7**. Sodium methoxide (0.4 eq) was generated in situ and reacted with the 2-cyano-pyridine **8** at room temperature. The reaction was stirred until full disappearance of the starting material. The carboximidate **9** was used directly in the next reaction in which it reacted with the methyl ester **7** to form the key intermediate, the oxazoline ester **6**, in 78% yield. Multigram quantities of **6** were easily available by this route.

Reduction of the ester group on **6** was attempted with a lithium aluminium hydride protocol²⁵. Lithium aluminium hydride (1.2 eq) was reacted with the ester **6** (1-2 g) in ice-cold diethyl ether. After 4 hours the starting material had been consumed. The reaction was quenched by the Fieser work-up method and the crude product was purified by column chromatography on silica gel. Isolation of the oxazoline alcohol **1** was achieved in a disappointing 31% yield. Other work-up methods were also attempted to quench the reaction, but no significant improvement was achieved. Due to the modest yield an alternative reduction method using sodium borohydride/methanol²⁶ was later attempted. Sodium borohydride (2.0 eq) was reacted with the ester **6** in methanol. After the work-up, the product was purified by crystallization (CH₂Cl₂/EtOAc) to give the alcohol **1** as a white powder in a much more satisfactory 72% yield. This reduction method became the method of choice for our study.



Reagents and conditions: (a) NaH, MeI, 0 °C to r.t, 3 h, 76% (b) 2-picolinic acid, 4-DMAP, dicyclohexylcarbodiimide (DCC), CH₂Cl₂, 0 °C to r.t, 4 h, 53% (c) benzoyl chloride, Et₃N, CH₂Cl₂, 0 °C to r.t, 3 h, 88%

Scheme 2. Synthesis of ligands **2-4**, related known compounds **10** and **11**.

With the oxazoline **1** in hand, three variations of the sidearm were developed (Scheme 2). These modifications aimed to investigate the effect of varying the group attached to the chiral center particularly to see what, if any, changes would induce increased interactions between the ligand and the catalytic centre. The increased interactions may give catalytic systems with better selectivity. Ligand **2** with ether functionality and ligand **3** with a pyridine ester were expected to demonstrate different coordination chemistry in the reaction both to each other and the original ligand **1**. The benzoate ester **4**, a similar compound to **3**, was also synthesized to help establish the coordination effects and catalytic outcomes attributable to the pyridine nitrogen on **4**. The ligands **2-4** prepared from **1** are all novel compounds and their synthesis was by conventional synthetic methods as outlined in Scheme 2.


During the course of this study we became aware of a report by Christoffers²⁷ where oxazoline **10** was synthesised (Scheme 2) via a different, more complex, route involving TBS protection and deprotection of the alcohol. This was the opposite enantiomer of oxazoline **1**. This compound was a synthetic intermediate in the synthesis of ligand **11**, and has not been used as a ligand in any catalytic reactions.

In a preliminary report by Kočovský et al on the use of 2-pyridyl mono(oxazoline) ligands a mechanism was proposed for the hydrosilylation of aromatic ketones. In this report the team proposed *N,N*-chelation of trichlorosilane to the mono(oxazoline) ligand.¹⁹ Consequently, their mechanistic proposal involves coordination of both the oxazoline nitrogen and the pyridyl nitrogen to the trichlorosilane.

We wanted to investigate this hypothesis. Therefore, we completed the synthesis of a 4-pyridyl mono(oxazoline) ligand in which the pyridyl moiety is derived from 4-cyanopyridine as opposed to 2-cyanopyridine. Our reasoning behind the use of 4-substituted pyridine ring was to maintain similar electronic properties between each family of ligands to allow us to directly study the coordinating effect of the nitrogen on the 2- position of the pyridine moiety.


We used the same synthetic methodology as previously described to construct the ligand, with the substitution of 4-cyanopyridine. Primary alcohol **12** can be derived from the cyclisation of methyl-(L)-serine **7**

with 4-cyanopyridine, followed by reduction of the resulting methyl ester **13** to the primary alcohol. Finally, esterification with the acid chloride led to the isolation of the ligand **14** (Scheme 3).



Reagents and conditions: (a) (i) 4-cyanopyridine, Na metal, MeOH, rt, 40h then **7**, CH₂Cl₂, 2.5 h, reflux, 61% (over 2 steps) (b) NaBH₄, MeOH, 4h, 0 °C-rt, 77% (c) acid chloride, Et₃N, 18 h, 0 °C-rt, 68%

Scheme 3. Synthesis of 4-pyridyl ligand **14**.




Scheme 4. Trichlorosilane-assisted hydrosilylations.

The hydrosilylation reactions of both ketone **15** and imine **17** were performed following identical procedures with the obvious exception of the use of different starting materials. The procedure described below was used for the hydrosilylation of acetophenone **15**. As shown in Scheme 4, trichlorosilane (2.1 equivalents) was added to a solution of acetophenone **15** (1.0 equivalent) and 20 mol% of the appropriate ligand **1-4** at -20 °C in dichloromethane (2 mL). The reaction was stirred for 24 hours during which time 1-phenylethanol **16** was formed. The crude product was purified by column chromatography and the enantiomeric excess (ee) measured using GC using a chiral CYCLODEX- β column.

The hydrosilylation of an imine was carried out in a similar manner. *N*-(1-phenylethylidene)aniline **17** was used instead of acetophenone **15**, and the reaction was performed at 0 °C instead of -20 °C. The crude product **18** was purified by column chromatography and the ee of the purified product was measured using chiral HPLC.

The catalytic results obtained in these two reactions using ligands **1-4** and ligand **14** are recorded in Tables 1 and 2.

Table 1. Hydrosilylation of acetophenone


Ligand	Conversion (%)	ee (%) (<i>R</i>)
Alcohol 1	99	42
Methyl ether 2	91	41
Picolinate ester 3	20	28
Benzoate 4	99	62
4-pyridyl ligand 14	0	-

As shown in Table 1 for the hydrosilylation of acetophenone, the original alcohol ligand **1** showed outstanding activity, giving 99% conversion to the desired product. The modified ligands were also highly efficient in catalysing the reaction with the exception of ligand **3** where a significant drop in activity was observed.

The enantiomeric excesses obtained with these ligands ranged from 28-62%. The transformation of the hydroxyl side-arm into its methyl ether counterpart (**1** to **2**) did not appear to make a difference to the enantioselectivity achieved in the reduction. The highest ee (62%) was achieved using ligand **4**, and interestingly, the lowest one (28%) was produced by the related ligand **3** where the phenyl ring on the side-arm was replaced by a pyridine. The ees achieved in the ketone reduction while not as high as those reported by Malkov¹⁹ are encouraging in terms of designing future generations of these ligands to achieve high selectivity.

In the hydrosilylation using ligand **3**, it was observed that some precipitate was gradually formed during the addition of trichlorosilane. It was speculated that the hypervalent silicate compound generated from the silane and ligand **3** precipitated due to its reduced solubility at low temperature (-20 °C). If that was the case, the reaction was catalysed by the reduced amount of homogeneous catalyst or if the precipitated silicate complex was also a catalyst, the catalysis was performed with both homogeneous and heterogeneous catalysts. We believe that these changes in the behaviour of the complexes were part of the reason for the poor catalytic performance of **3**.

The complete lack of reactivity observed when ligand **14** the 4-pyridyl ligand was used seems to add credence to the mechanism proposed by Kočovský. It is apparent that moving the pyridyl nitrogen effectively kills the catalysis so the 2-pyridyl group is key.

Table 2. Hydrosilylation of of *N*-(1-phenylethylidene)aniline

Ligand	Conversion (%)	ee (%)
Alcohol 1	99	28(<i>R</i>)
	99 ($-10\text{ }^\circ\text{C}$)	33(<i>R</i>)
Methyl ether 2	99	10(<i>S</i>)
Picolinate ester 3	99	14(<i>R</i>)
Benzoate 4	99	4(<i>R</i>)

In the hydrosilylation of *N*-(1-phenylethylidene)aniline, the activities of the ligands were excellent, with 99% conversions obtained in all cases (Table 2). The enantioselectivities achieved in this reaction were not as good as in the hydrosilylation of acetophenone. The highest ee (28%) was obtained with the alcohol ligand **1** favouring the (*R*) enantiomer. Interestingly, changing the side-arm from the OH group to an OMe group resulted in a significant change of selectivity. Reaction using ligand **2** gave product with a 10% ee favouring the (*S*) enantiomer, a 38% swing in ee compared to that obtained with ligand **1**. Thereby we postulate that the OH sidearm of ligand **1** is playing a role mechanistically. As this was not seen in the hydrosilylation of acetophenone, it is likely that the proton of the OH side-arm was interacting, probably via hydrogen-bonding, with the imine substrate. Replacing the OH group of **1** with the OMe group would prevent this interaction and this would account for the significant change of selectivity observed.

Having determined how synthetically simple the preparation of benzoate ester **4** had turned out to be and having seen that we could achieve enantioselective catalysis in the reduction reactions, we decided to prepare another six related derivatives. Using different acyl chlorides but the same method we prepared **19-24** in yields ranging from 75-87% (Figure 3).


**Figure 3.** Second group of 2-pyridyl ligands prepared for this study.

Table 3. Hydrosilylation of acetophenone


Entry	Ligand	Conversion (%)	ee (%) ^a
1	19	78	44
2	20	3	48
3	21	>99	1
4	22	>99	46
5	23	65	0.6
6	24	99	47

^a determined by chiral GC (Cyclodex β column, 120 °C, 20 min)

The hydrosilylation reaction was used to test the catalytic performance of the chiral mono(oxazoline) ligands **19-24**. The results for the reaction shown in Scheme 5, using 20 mol% of appropriate ligand are shown in Table 3.

Excellent conversions were observed for ligands **21**, **22** and **24**. Modest enantioselectivities were obtained for ligands **19**, **20**, **22**, **24**. The ligands with the naphthyl ester **21** and the *p*-methoxyphenyl ester **23** show the poorest enantioselectivities overall, such that almost no facial selectivity is shown (Entries 3 and 5). It appears that alkyl, electron-deficient aryl or mildly electron-rich aryl groups are tolerated well but the electron-rich methoxy-aryl is less reactive and less selective and the naphthoate is not well tolerated when it comes to selectivity. Benzoate esters appear superior to ethers in terms of selectivity.

To rationalise the results Kočovský and co-workers obtained, they postulated the following mechanistic transformation (Scheme 5)


**Scheme 5.** Kočovský's proposed mechanism for the hydrosilylation using 2-pyridyl oxazoline ligands.

N,N-Chelation of Cl_3SiH of by the 2-pyridyl mono(oxazoline) leads to the formation of the activated hydrosilylating species, seen in Scheme 5 above. Trichlorosilane does not reduce the carbonyl species without the presence of a Lewis base, supporting the formation of the hexacoordinated species seen above. Another molecule of Cl_3SiH is likely to activate the ketone by coordination to the oxygen atom. It is unlikely that the carbonyl will displace a chloride ion to coordinate to the activated silicon centre, as the corresponding four membered transition state would not be energetically favourable. This ketone- Cl_3SiH complex can then approach the catalyst- Cl_3SiH from the less hindered face, as dictated by the orientation of the chiral centre. Kočovský *et al.* also suggested that this interaction will also be stabilised by arene-arene interactions between the phenyl ring of the substrate and the pyridyl ring of the chiral catalyst (Scheme 5).

We postulate that a possible competitive arene-arene interaction between the incoming acetophenone and the naphthyl moiety may be playing a role in the lack of selectivity seen for ligand **21**.

As established above, *p*-methoxyphenyl ester substituted ligand **23** was the second ligand to show disappointing selectivity. This effect could be attributed to the strong electron-donating effect exhibited by the *para*-methoxy moiety. Through resonance, it is possible to move the lone pair of electrons, originating from the oxygen of the methoxy group, as far as the ester oxygen. This may suggest that where the ligand is an ester this ester is somehow involved in the reaction mechanism and altering the electron density on the oxygen affects the selectivity.

Following on from this we wanted to establish the versatility of the reaction as regards to substrate scope. Chiral alcohols are important building blocks in organic synthesis and are commonly used synthons in the pharmaceutical industry, such as in the total synthesis of Duloxetine.²⁸ Therefore, it is desirable that our ligands are able to catalyse the transformation of various ketones. Scheme 6 shows the transformation of 1-(4-methyl)ethanone **25** to the corresponding alcohol **26**.



Scheme 6. Hydrosilylation of 1-(4-methyl)ethanone **25**.

The transformation was carried out as described for acetophenone **15** above. The percentage conversion was calculated by comparing the amount of unreacted 1-(4-methyl)ethanone **25** [signal at 2.54 ppm (3H)] to that of the product **26** [signal at 1.48 ppm (3H)] in the ¹H-NMR spectrum. Following purification, the ee was measured by chiral GC. The results of the reactions using each ligand are shown in Table 4 below.

Table 4. Hydrosilylation of 1-(4-methyl)ethanone

Entry	Ligand	Conversion (%)	ee (%) ^a
1	19	>99	38
2	20	3	38
3	21	92	10.5
4	23	91	3
5	24	7	35

^a determined by chiral GC (Cyclodex β column, 120 °C, 20 min)

The same general trend can be seen to be as was seen for acetophenone (Table 3). Low selectivities were again observed for ligands **21** and **23** and again the remaining ligands showed almost consistent selectivities. Although an overall decrease in enantioselectivity was seen for ligands **19**, **20** and **24** (Entries 1, 2 and 5) compared to their performance with acetophenone. We can conclude that the addition of the methyl group has an overall negative effect on the selectivity of the reaction.

To that point we had seen that ligands with an ester side-arm showed good selectivity in the reduction of acetophenone. Our initial benzoate gave the highest ee with variations in the aromatic group leading to decreased selectivity and no clear correlation in performance with electronic effects in the aromatic group. The larger naphthyl ester showed no selectivity. We noted that the adamantyl ester performed quite well and we prepared a number of other non-aromatic esters **27-30** to assess whether these would give better performance and a clearer correlation between performance and the ester group. Moving away from aromatic esters also allowed us to try chiral esters to see if there was an additive effect from multiple chiral centres in the ligand. We also prepared the pyridin-2-ylmethyl ether **30** to see what effect a third coordinating nitrogen might have on the selectivity. Thus, using similar chemistry to before, we prepared ligands **27-29** (Figure 4)









Figure 4. Third group of pyridin-2-yl ligands prepared for this study.

Table 5. Catalytic results in the hydrosilylation of acetophenone using ligands **27-30**





Ligand	Conversion (%)	ee (%)
 27	99	57 (R)
 28	98	52 (R)
 29	99	55 (R)
 30	85	40 (R)

These proved to be equally active to those studied previously, giving conversions to the chiral product ranging from 85-99%. The ee obtained with the pyridin-2-ylmethyl ether **30** (40%) was similar to that obtained with the alcohol ligand **1** (42%). This result added to our suspicion that the carbonyl group of the ligand ester was a key feature required to achieve high enantioselectivity using this type of ligand. The similar ees achieved in the reaction by ligands **1** and **30** also indicated that the coordination of the side-arm pyridin-2-yl nitrogen to the silicon centre was less likely.

Interestingly, varying the steric bulk of the group attached to the carbonyl group did not appear to have substantial influence on enantioselectivity. In fact, ligands bearing different ester side-arms achieved similar enantioselectivities in the reactions. The pivalate ester **27** delivered the highest ee (57%) among the four modified ligands. The benzoate ester **29** exhibited similar enantioselectivity (52%) to the ligand bearing a chiral group on the ester **28** (55%). It had been hoped that this chiral ester group would show a substantial additive (or reductive) effect on the stereoselectivity.

As they gave very promising selectivities, ligands **4** and **27**, were selected to study the effect of reaction temperature on the reactivity and selectivity.

Table 6. Catalytic results of **4** and **27** with variation in reaction temperature

Ligands	Temp (°C)	Time (h)	Conversion(%)	ee (%)
 4	-20	24	99	62 (R)
	-40	24	99	53 (R)
 27	-20	24	99	57 (R)
	-40	24	96	65 (R)
	-78	6	14	70 (R)

As shown in Table 6, ligand **4** was highly active even at -40 °C, although its enantioselectivity was slightly reduced at that temperature. Similar precipitation was observed during the addition of trichlorosilane, as was described above for the reaction catalysed by ligand **3**. It was once again speculated that the hypervalent silicon complex was precipitated due to reduced solubility, which contributed to the decline of selectivity in the reaction because this reduced the effective concentration of the catalyst, which could lead to this selectivity decrease.

Ligand **27** also demonstrated impressive activities in this reaction. When the reaction was running at -40 °C a slight drop in conversion (3%) was observed compared to the reaction at -20 °C. At -78 °C this ligand was still reasonably active as 14% conversion was achieved in 6 hours. During the tests with ligand **27**, improvements in selectivity were seen as the temperature was reduced. The ees ranged from from 57% at -20 °C to 65% at -40 °C and finally 70% at -78 °C.

Conclusions

We have reported the synthesis of four novel ligands and their application in asymmetric reduction reactions. The ligands acted as organocatalysts demonstrating consistently high activities in the hydrosilylation of a

ketone and an imine. The exception was ligand **3**, which produced only a 20% conversion in the hydrosilylation of a ketone, possibly due to the solubility issue as was discussed previously.

The ees obtained in the hydrosilylation of acetophenone were improved by the modifications. The ees obtained in this reaction increased from 41% using the original alcohol **1** to 62% using ligand **4** and in general, ligands with ester side-arms were more selective. With the *t*-Bu substituted ligand which showed good selectivity and activity in the reaction at -20 °C, reduction at -40 °C still gave good conversion and an ee of 65%. At -78 °C we achieved an ee of 70% albeit with a decrease in catalyst activity.

The ees obtained in the reduction of *N*-(1-phenylethylidene)aniline were adversely affected by the modifications to the side-arm as the highest ee (28%) was produced by the alcohol **1**.

We are continuing to develop these ligands with modifications based on these results and the application these ligands to other reactions as organocatalysts and as part of catalytic complexes.

Experimental Section

General. All procedures were carried out under an atmosphere of nitrogen unless otherwise stated. All solvents were distilled prior to use as follows: dichloromethane and acetonitrile were distilled from calcium hydride, toluene and tetrahydrofuran were distilled from sodium benzophenone ketyl or dried through an Innovative Technology Pure Solv PS-MD-5 Purification System. Organic phases were dried using anhydrous magnesium sulphate or sodium sulphate. All chemicals were purchased from Aldrich Chemical Company, Fisher Scientific or Acros Organics and were used without further purification. All asymmetric reactions were carried out using standard Schlenk line techniques and all Schlenk tubes were heat dried under vacuum to remove moisture prior to use.

Melting points were measured on a Stuart Scientific SMP3 or SMP11 apparatus and are uncorrected. IR spectra were measured on a Perkin Elmer Spectrum One FT-IR, in which liquid samples were measured as thin films and solids were measured directly. Optical rotations were measured on a Uniplol L1000 polarimeter at 589 nm (Na) in a 1 dm cell; concentrations (*c*) are expressed in g/100 mL. $[\alpha]_D$ is the specific optical rotation of a compound and is measured in units of $\text{deg}\cdot\text{mL}\cdot\text{g}^{-1}\cdot\text{dm}^{-1}$. Thin layer chromatography (TLC) was carried out on pre-coated silica gel plates (Merck 60 F254); column chromatography was carried out using Apollo Scientific silica gel 40-63 micron. Visualisation was achieved by UV (254 nm) light detection, vanillin stain or ninhydrin stain. High resolution mass spectra were carried out using electrospray ionisation (ESI) on a Walters LCT Premier XE spectrometer by manual peak matching. ^1H NMR (400 MHz) and ^{13}C NMR (100 MHz) spectra were recorded on a JEOL ECX-400 NMR spectrometer. All spectra were recorded at probe temperatures (~ 20 °C) in deuterated chloroform (CDCl_3) unless otherwise stated, using tetramethylsilane (TMS) as an internal standard. Chemical shifts are expressed in parts per million (ppm) and coupling constants in Hertz (Hz), unless indicated all coupling constants represent 3 bond couplings.

All chiral HPLC analysis was carried out on a Varian Prostar instrument, with a UV/Vis detector at the specified wavelength, with a CHIRACEL OD 0.46 cm Φ x 25 cm column under conditions described for each experiment. All chiral GC analysis was carried out on a Varian 3900 instrument, using helium as the mobile phase and a FID (Flame Ionisation Detector), with a CYCLODEX- β 0.25 mm Φ x 30 m column under conditions described for each experiment.

Methyl L-serinate hydrochloride (7). Thionyl chloride (4.4 mL, 60 mmol) was added dropwise to a suspension of L-serine **5** (6.34 g, 60 mmol) in anhydrous MeOH (60 mL) at 0 °C. The mixture was allowed to warm to rt and

stirred for a further 2 h at rt. The reaction was then heated at reflux overnight. The solvent was removed under reduced pressure and the resulting solid residue was stirred in Et₂O until a fine powder precipitated. The powder was removed by filtration and the supernatant dried under vacuum to yield compound **7** (9.33 g, 59.9 mmol, 99%) as a white powder.

¹H NMR spectroscopic data was in accordance with the literature.^{24,29}

¹H NMR (D₂O, 400 MHz): δ (ppm) 4.29 (1H, t, *J* 3.8), 4.08 (2H, m, *J* 12.5, 3.8), 3.87 (3H, s); ¹³C NMR (D₂O, 100 MHz): δ (ppm) 169.0, 59.3, 54.7, 53.7.

Methyl (4S)-2-(pyridin-2-yl)-4,5-dihydrooxazole-4-carboxylate (6). Sodium metal (560 mg, 24 mmol) was added to anhydrous MeOH (60 mL) and the mixture was stirred until bubbling ceased at which stage 2-cyanopyridine **8** (6.24 g, 60 mmol) was added. After stirring for 24 h, the solvent was removed under reduced pressure and the crude product **9** was transferred to a suspension of compound **7** (9.33 g, 59.9 mmol) in anhydrous CH₂Cl₂ (60 mL). The reaction mixture was then heated at reflux for 2.5 h, cooled and the inorganic precipitate filtered off. The solvent was removed under reduced pressure and the resulting crude product was purified by column chromatography (silica gel, EtOAc-PE, 80:20) to yield the oxazoline ester **6** (9.66 g, 46.8 mmol, 78%) as a yellow oil. [α]_D²⁰ +96.8 (c 0.50, CHCl₃); ¹H NMR (CDCl₃, 400 MHz): δ (ppm) 8.46 (1H, d, *J* 4.7 Hz), 7.85 (1H, d, *J* 7.9 Hz), 7.56 (1H, td, *J* 7.8, 1.2 Hz), 7.23 – 7.15 (1H, m), 4.80 (1H, dd, *J* 10.8, 8.2 Hz), 4.55 – 4.42 (2H, m), 3.57 (3H, s); ¹³C NMR (CDCl₃, 101 MHz): δ (ppm) 171.1, 165.2, 149.7, 145.8, 136.7, 126.0, 124.3, 70.1, 68.6, 52.6; IR (cm⁻¹): 2954, 1737, 1636, 1581, 1570, 1436, 1364, 1204.

[(4R)-2-(Pyridin-2-yl)-4,5-dihydrooxazol-4-yl]methanol (1). Sodium borohydride pellets (1.52g, 40 mmol) were added to an ice-cold solution of compound **6** (4.12 g, 20 mmol) in MeOH (200 mL). After gas evolution had ceased, the reaction was allowed to warm up to rt and was further stirred for 4 h. The solvent was removed and the solid residue was partitioned between H₂O (50 mL) and CH₂Cl₂ (50 mL), the layers were separated and the aqueous layer was extracted with CH₂Cl₂ (50 mL x 3). The combined organic layers were dried (Na₂SO₄) and filtered, and the solvent removed under reduced pressure. The crude product was then purified by crystallization (CH₂Cl₂/EtOAc) to yield the oxazoline alcohol **1** (2.55 g, 14.3 mmol, 72%) as a white powder.

All spectroscopic data was in accordance with the literature.²⁷

¹H NMR (CDCl₃, 400 MHz): δ (ppm) 8.69 (1H, ddd, *J* 4.8, 1.6, 0.9 Hz), 7.93 (1H, dt, *J* 7.9, 1.2 Hz), 7.76 (1H, td, *J* 7.8, 1.6 Hz), 7.35 (1H, ddd, *J* 7.6, 4.8, 0.9 Hz), 4.58 (1H, dd, *J* 9.7, 7.3 Hz), 4.57 – 4.45 (1H, m), 4.41 (1H, t, *J* 7.4 Hz), 3.98 (1H, dd, *J* 11.4, 3.1 Hz), 3.70 (1H, dd, *J* 11.0, 3.4 Hz), 2.45 (1H, s); ¹³C NMR (CDCl₃, 101 MHz): δ (ppm) 164.5, 150.0, 146.3, 136.8, 125.8, 124.0, 70.0, 68.5, 64.2.

(4R)-2-(Pyridin-2-yl)-4,5-dihydrooxazol-4-ylmethyl pyridine-2-carboxylate (3). Dicyclohexylcarbodiimide (680 mg, 3.3 mmol) was added portionwise to an ice-cold solution of oxazoline alcohol **1** (534 mg, 3 mmol), picolinic acid (406 mg, 3.3 mmol) and 4-dimethylaminopyridine (36.5 mg, 0.3 mmol) in anhydrous CH₂Cl₂ (20 mL). After stirring for 10 min, the reaction was allowed to warm up to rt and stirred for a further 4 h. The mixture was filtered and concentrated under reduced pressure. The resulting crude product was then purified by column chromatography (silica gel, CH₂Cl₂-MeCN, 95:5) to yield picolinate ester **3** (453 mg, 1.6 mmol, 53%) as a colorless oil. [α]_D²⁰ +70.5 (c 0.49, CHCl₃); ¹H NMR (CDCl₃): δ 8.78-8.72 (2H, m), 8.14 – 8.05 (2H, m), 7.87 – 7.76 (2H, m), 7.52 – 7.39 (2H, m), 4.79 – 4.90 (1H, m), 4.74 – 4.64 (2H, m), 4.57 (1H, dd, *J* 11.2, 6.5 Hz), 4.50 (1H, dd, *J* 8.8, 7.6 Hz); ¹³C NMR (CDCl₃): δ 164.8, 164.5, 149.9, 149.8, 147.5, 146.2, 137.0, 136.7, 127.0, 125.8, 125.3, 124.2, 70.8, 66.9, 65.6; IR (cm⁻¹): 3422, 2964, 1732, 1640, 1583, 1133; ESI-HRMS calcd for C₁₅H₁₃N₃O₃ 284.1035 [M+H⁺], found *m/z* 284.1034.

(4R)-2-(Pyridin-2-yl)-4,5-dihydrooxazol-4-ylmethyl benzoate (4). Benzoyl chloride (510 mg, 3.6 mmol) was added slowly to an ice-cold solution of oxazoline alcohol **1** (534 mg, 3 mmol) and Et₃N (450 mg, 4.5 mmol) in

anhydrous CH_2Cl_2 (20 mL). The reaction was allowed to warm up to rt and stirred for a further 3 h before it was quenched by H_2O (2 mL). The resulting mixture was washed with H_2O (2 x 15 mL), dried (Na_2SO_4) and concentrated under reduced pressure to give the crude product, which was then purified by column chromatography (silica gel, EtOAc-PE, 90:10) to yield benzoate ester **4** (744 mg, 2.64 mmol, 88%) as a light yellow oil. $[\alpha]_{\text{D}}^{20} +16.7$ (c 0.28, CHCl_3); ^1H NMR (CDCl_3): δ 8.72 (1H, dd, J 4.8, 1.7 Hz), 8.06 (1H, d, J 8.0 Hz), 7.99 (2H, dd, J 8.1, 1.1 Hz), 7.79 (1H, td, J 7.8, 1.8 Hz), 7.57–7.48 (1H, m), 7.45–7.34 (3H, m), 4.83–4.71 (1H, m), 4.66 (1H, dd, J 9.8, 8.6 Hz), 4.59 (1H, dd, J 11.3, 4.3 Hz), 4.53–4.44 (2H, m); ^{13}C NMR (CDCl_3): δ 166.4, 164.5, 149.8, 146.3, 136.7, 133.1, 129.7, 128.5, 126.0, 124.2, 70.6, 66.1, 65.8; IR (cm^{-1}): 2969, 1717, 1640, 1582, 1110; ESI-HRMS calcd for $\text{C}_{16}\text{H}_{14}\text{N}_2\text{O}_3$ Na 305.0902 [$\text{M}+\text{Na}^+$], found m/z 305.0904.

2-[(4R)-4-Methoxymethyl-4,5-dihydrooxazol-2-yl]pyridine (2). Sodium hydride (60% dispersion in mineral oil, 192 mg, 4.8 mmol) was added to an ice-cold solution of oxazoline alcohol **1** (534 mg, 3 mmol) in anhydrous THF (20 mL). After the bubbling caused by gas evolution ceased, MeI (511 mg, 3.6 mmol) was added slowly and the reaction was allowed to warm up to rt. After stirring the mixture for a further 3 h, the precipitate was filtered off and the filtrate was concentrated under reduced pressure. The resulting residue was partitioned between H_2O (15 mL) and CH_2Cl_2 (15 mL), and the separated aqueous layer was further extracted with CH_2Cl_2 (2 x 15 mL). The combined organic layers were dried (Na_2SO_4), filtered and concentrated under reduced pressure to give the crude product, which was then purified by column chromatography (silica gel, EtOAc-MeOH, 95:5) to yield the methyl ether **2** (437 mg, 2.28 mmol, 76%) as a light-yellow oil. $[\alpha]_{\text{D}}^{20} +40.0$ (c 1.00, CHCl_3); ^1H NMR (CDCl_3): δ 8.69 (1H, ddd, J 4.6, 1.6, 0.9 Hz), 8.04 (1H, d, J 7.8 Hz), 7.76 (1H, td, J 7.8, 1.8 Hz), 7.43–7.34 (1H, m), 4.62–4.48 (2H, m), 4.41–4.33 (1H, m), 3.73–3.64 (1H, m), 3.54–3.45 (1H, m), 3.39 (3H, s); ^{13}C NMR (CDCl_3): δ 164.0, 149.8, 146.6, 136.7, 125.7, 124.1, 74.6, 71.2, 66.6, 59.4; IR (cm^{-1}): 3412, 2901, 1641, 1583, 1113; ESI-HRMS calcd for $\text{C}_{10}\text{H}_{12}\text{N}_2\text{O}_2$ 193.0977 [$\text{M}+\text{H}^+$], found m/z 193.0975.

Asymmetric trichlorosilane-assisted hydrosilylation of aromatic ketone and imine. General procedure B

All Schlenk tubes and round-bottom flasks were flame dried under vacuum and re-filled with nitrogen gas before being used in these reactions. Aromatic ketone or imine (1.0 equiv) was added to a solution of ligand (20 mol %) in anhydrous CH_2Cl_2 (2 mL) at the appropriate temperature. Trichlorosilane (2.1 equiv) was then added and the reaction mixture was stirred at that temperature for a specific period of time.

The reaction was quenched by aqueous NaHCO_3 (2 mL) and the resulting mixture was extracted with CH_2Cl_2 (2 x 5 mL). The combined organic layers were dried (Na_2SO_4) and concentrated under reduced pressure to give the crude product. A ^1H NMR spectrum of the crude product was recorded to determine the percentage conversion of this reaction. [For the reduction of acetophenone, the CH_3 signal of the unreacted ketone appeared at 2.58 ppm (3H) versus the CH_3 signal of the product, appearing at 1.52 ppm (3H); for the reduction of *N*-(1-phenylethylidene)aniline, the CH_3 signal of the unreacted imine appeared at 2.26 ppm (3H) versus the CH_3 signal of the product, at 1.43 ppm (3H)]. The crude product was then purified by column chromatography (silica gel). The enantiomeric excess (ee) of the purified product was measured using chiral GC (Cyclodex- β 30 m x 0.252 mm 0.25 μm) or chiral HPLC (DAICEL Chiracel OD 25 cm x 4.6 mm 5 μm).

Methyl-(4S)-2-(pyridin-4-yl)-4,5-dihydrooxazole-4-carboxylate (13). To a stirring solution of anhydrous MeOH (80 mL) at rt was added Na metal (760 mg, 32 mmol) under N_2 . Upon cessation of gas evolution, 4-cyanopyridine (8.48 g, 0.08 mol) was added and the resulting mixture was stirred at rt for 40 h. Concentration at reduced pressure yielded an off-white solid, which was dissolved in CH_2Cl_2 (60 mL). Following this, ester **7** (7.05 g, 0.06 mmol) was added and the resulting mixture was heated to reflux and stirred for 2.5 h. The solution was allowed to cool to rt and excess solvent was removed under vacuum. The resulting oil was dissolved in EtOAc/ H_2O (30 mL/30 mL) and the separated aqueous layer was extracted with EtOAc (3 x 30 mL). The combined organic extracts were dried over Na_2SO_4 , filtered and concentrated to yield **13** as brown oil.

Purification of crude **13** was achieved by column chromatography in EtOAc:MeOH (95:5) to yield **13** as a clear oil (7.61 g, 61%). $[\alpha]_D +103.2$ (c 5.05 $\times 10^{-3}$, CHCl₃, 20 °C), ¹H NMR (400 MHz, CDCl₃) δ 8.69 (2H, d, J 5.9 Hz), 7.78 (2H, d, J 5.9 Hz), 4.96 (1H, dd, J 10.7, 8.2 Hz), 4.71 (1H, m), 4.61 (1H, m), 3.80 (3H, s) ppm; ¹³C NMR (100 MHz, CDCl₃) δ 171.1, 164.7, 150.3, 134.2, 122.1, 69.9, 68.7, 53.0 ppm; IR 2983, 2887, 1741, 1530 cm⁻¹; ESI-HRMS calcd. for C₁₀O₃N₂H₁₀ 207.0769 (M+H)⁺ found m/z 207.0748.

[(4R)-2-(Pyridin-4-yl)-4,5-dihydrooxazol-4-yl]methanol (12). To a stirring solution of **13** (4.86 g, 23.34 mmol) at 0 °C in MeOH (360 mL) was added NaBH₄ (3.20 g, 83.6 mmol). Upon cessation of gas evolution the resulting solution was heated to rt and stirred under N₂ for 4 h. Excess MeOH was removed under vacuum, and the resulting yellow solid was dissolved in CH₂Cl₂/H₂O (15 mL/15 mL). The aqueous layer was extracted with CH₂Cl₂ (5 x 15 mL). The combined organic extracts were dried over NaSO₄, filtered and concentrated under reduced pressure to yield crude **12** as an off-white solid. Recrystallisation of the crude product from CH₂Cl₂/EtOAc at 0 °C yielded **12** as a white solid (3.23g, 77%). $[\alpha]_D +63.1$ (c 5.10 $\times 10^{-3}$, CHCl₃, 20 °C); ¹H NMR (400 MHz, CDCl₃) δ 8.62 (2H, d, J 4.5 Hz), 7.64 (2H, d, J 4.5 Hz), 4.54-4.38 (3H, m), 4.00 (1H, d, J 11.6 Hz), 3.68 (1H, d, J 11.7 Hz) ppm; ¹³C NMR (400 MHz, CDCl₃) δ 163.8, 150.2 (2 x CH), 134.5, 122.0 (2 xCH), 69.6, 68.4, 63.5; IR 3293, 2903, 1604, 845 cm⁻¹; MP 49-51 °C; ESI-HRMS calcd. for C₉O₂N₂H₁₀ 179.0820 (M+H)⁺ found m/z 179.0812.

[(4S)-2-(Pyridin-4-yl)-4,5-dihydrooxazol-4-yl]methyl-4-methylbenzoate (14). To a stirring solution of **12** (216 mg, 1.2 mmol) in CH₂Cl₂ (5 mL) at 0 °C was added NEt₃ (0.33 mL, 2.4 mmol) and *p*-toluoyl chloride (0.24 mL, 1.8 mmol). The reaction was stirred at 0 °C after which it was heated to rt and stirred for 18 h. Addition of H₂O (5 mL) was followed by extraction of the aqueous layer with CH₂Cl₂ (3 x 5 mL). The combined organic extracts were washed with brine (10 mL), dried over NaSO₄, filtered and concentrated *in vacuo*. The resulting oil was purified by column chromatography (PE:EtOAc, 9:1) to yield the mono(oxazoline) ligand **14** as a white solid (233 mg, 68%). $[\alpha]_D +18.0$ (c 5.00 $\times 10^{-3}$, CHCl₃, 20 °C); ¹H NMR (400 MHz, CDCl₃) δ 8.72 (2H, d, J 6.0 Hz), 7.85 (2H, d, J 8.2 Hz), 7.79 (2H, d, J 6.0), 7.18 (2H, d, J 8.2 Hz), 4.74-4.70 (1H, m), 4.60-4.52 (2H, m), 4.46-4.40 (2H, m), 2.38 (3H, s) ppm; ¹³C NMR (100 MHz, CDCl₃) δ 166.5, 163.6, 150.3 (2 x CH), 143.9, 134.7, 129.7 (2 x ArCH), 129.2 (2 x ArCH), 126.9, 122.0 (2 x CH), 70.4, 65.9, 65.9, 21.8 ppm; IR 2959, 2898, 1715, 1648, 837 cm⁻¹; mp 105-108 °C; ESI-HRMS calcd. for C₁₇O₃N₂H₁₆ 297.1239 found (M+H)⁺ m/z 297.1225.

[(4S)-2-(Pyridin-2-yl)-4,5-dihydrooxazol-4-yl]methylnaphthalene-1-carboxylate (21). This was prepared according to the procedure reported above for benzoate **3** using oxazoline **1** (288 mg, 1.6 mmol) in CH₂Cl₂ (5 mL), NEt₃ (0.45 mL, 3.2 mmol) and 1-naphthoyl chloride (0.37 mL, 2.4 mmol). The resulting crude product was purified by column chromatography (PE:EtOAc, 9:1) to yield the mono(oxazoline) ligand **21** as a viscous straw-coloured oil (489 mg, 75%). $[\alpha]_D +5.1$ (c 5.10 $\times 10^{-3}$, CHCl₃, 20 °C); ¹H-NMR (400 MHz, CDCl₃) δ 8.90-8.85 (1H, m), 8.73 (1H, d, J 4.2 Hz), 8.15 (1H, d, J 7.3 Hz), 8.07 (1H, d, J 7.8 Hz), 7.99 (1H, d, J 8.2 Hz), 7.87-7.82 (1H, m), 7.81-7.86 (1H, m), 7.51-7.47 (2H, m), 7.43-7.39 (2H, m), 4.87-4.79 (1H, m), 4.72-4.59 (3H, m), 4.55-4.51 (1H, m) ppm; ¹³C-NMR (100 MHz, CDCl₃) δ 167.4, 164.6, 150.0, 146.4, 136.8, 133.9, 133.8, 131.4, 130.6, 128.6, 127.9, 127.0, 126.3, 126.0, 125.8, 124.6, 124.2, 70.7, 66.2, 65.9 ppm; IR 3371, 3056, 2953, 1710, 1509, 1238 cm⁻¹; ESI-HRMS calcd. for C₂₀O₃N₂H₁₆ 333.1293 (M+H)⁺ found m/z 333.1196.

(4S)-2-(Pyridin-2-yl)-4,5-dihydrooxazol-4-ylmethyl adamantane-1-carboxylate (24). This was prepared according to the procedure reported above for benzoate **3** using oxazoline **1** (288 mg, 1.6 mmol) in CH₂Cl₂ (5 mL), NEt₃ (0.45 mL, 3.2 mmol) and 1-adamantanecarbonyl chloride (483 mg, 2.4 mmol). After purification the reaction yielded the mono(oxazoline) ligand **24** as a viscous yellow oil (476 mg, 87%). $[\alpha]_D +20.0$ (c 4.90 $\times 10^{-3}$, CHCl₃, 20 °C); ¹H-NMR (400 MHz, CDCl₃) δ 8.70 (1H, d, J 4.8 Hz), 8.03 (1H, d, J 7.9 Hz), 7.78 (1H, td, J 7.8, 1.9 Hz), 7.40 (1H, dd, J 6.4, 4.8 Hz), 4.64-4.53 (2H, m), 4.37-4.30 (2H, m), 4.22 (1H, dd, J 11.3, 5.2), 1.93-1.58 (15H, m) ppm; ¹³C-NMR (100 MHz, CDCl₃) δ 177.5, 164.5, 149.9, 146.5, 136.8, 125.9, 124.1, 70.6, 65.8, 65.1, 40.9,

38.8, 36.5, 27.9 ppm; IR 3391, 2904, 2851, 1724, 1641, 1071 cm^{-1} ; ESI-HRMS calcd. for $\text{C}_{20}\text{O}_3\text{N}_2\text{H}_{24}$ 341.1865 found $(\text{M}+\text{H})^+$ m/z 341.1831.

(4S)-2-(Pyridin-2-yl)-4,5-dihydrooxazol-4-ylmethyl 4-chlorobenzoate (20). This was prepared according to the procedure reported above for benzoate **3** using oxazoline **1** (288 mg, 1.6 mmol) in CH_2Cl_2 (5 mL), NEt_3 (0.45 mL, 3.2 mmol) and 4-chlorobenzoyl chloride (0.3 mL, 2.4 mmol). After purification the reaction yielded the mono(oxazoline) **20** ligand as a pale yellow solid (362 mg, 76%). $[\alpha]_D^{20} +17.1$ (c 5.50 $\times 10^{-3}$, CHCl_3 , 20 $^\circ\text{C}$); $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 8.72 (1H, d, J 4.8 Hz), 8.05 (1H, d, J 7.9 Hz), 7.92 (2H, d, J 8.4 Hz), 7.79 (1H, td, J 7.8, 1.7 Hz), 7.42 (1H, dd, J 6.5, 4.8 Hz), 7.36 (2H, d, J 8.4 Hz), 4.79-4.72 (1H, m), 4.66 (1H, dd, J 9.8, 8.7 Hz), 4.56 (1H, dd, J 11.3, 4.3 Hz), 4.51-4.45 (2H, m) ppm; $^{13}\text{C-NMR}$ (400 MHz, CDCl_3) δ 165.6, 164.6, 150.0, 146.4, 139.7, 136.9, 131.2, 128.8, 128.3, 126.0, 124.2, 70.6, 66.3, 65.8 ppm; IR 2968, 1716, 1639, 1591, 1276 cm^{-1} ; mp 92-97 $^\circ\text{C}$; ESI-HRMS calcd. for $\text{C}_{16}\text{O}_3\text{N}_2\text{H}_{13}\text{Cl}$ 317.0693 $(\text{M}+\text{H})^+$ found m/z 317.0646.

(4S)-2-(Pyridin-2-yl)-4,5-dihydrooxazol-4-ylmethyl 4-nitrobenzoate (22). This was prepared according to the procedure reported above for benzoate **3** using oxazoline **1** (288 mg, 1.6 mmol) in $\text{CH}_2\text{m/zCl}_2$ (5 mL), NEt_3 (0.45 mL, 3.2 mmol) and 4-nitrobenzoyl chloride (410 mg, 2.4 mmol). After purification the reaction yielded the mono(oxazoline) ligand **22** as a yellow solid (394 mg, 75%). $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 8.72 (1H, d, J 4.3 Hz), 8.22 (2H, d, J 9.0 Hz), 8.15 (2H, d, J 8.9 Hz), 8.04 (1H, d, J 7.9 Hz), 7.80 (1H, td, J 7.8, 1.8 Hz), 7.43 (1H, dd, J 6.6, 4.9 Hz), 4.82-4.76 (1H, m), 4.69-4.65 (1H, m), 4.58 (2H, d, J 4.6 Hz), 4.47 (1H, dd, J 8.6, 7.3 Hz) ppm; $^{13}\text{C-NMR}$ (100 MHz, CDCl_3) δ 164.7, 164.5, 150.6, 149.9, 146.1, 136.8, 135.0, 130.8, 126.0, 124.1, 123.5, 70.3, 66.8, 65.5 ppm; IR 2902, 1726, 1640, 1516, 1270 cm^{-1} ; MP 117-120 $^\circ\text{C}$; ESI-HRMS calcd. for $\text{C}_{16}\text{O}_5\text{N}_3\text{H}_{13}$ 328.0933 $(\text{M}+\text{H})^+$ found m/z 328.0901;

(4S)-2-(Pyridin-2-yl)-4,5-dihydrooxazol-4-ylmethyl 4-methylbenzoate (19). This was prepared according to the procedure reported above for benzoate **3** using oxazoline **1** (288 mg, 1.6 mmol) in CH_2Cl_2 (5 mL), NEt_3 (0.45 mL, 3.2 mmol) and *p*-toluoyl chloride (0.32 mL, 2.4 mmol). After purification the reaction yielded the mono(oxazoline) ligand **19** as a white solid (320 mg, 75%). $[\alpha]_D^{20} +13.0$ (c 2.15 $\times 10^{-3}$, CHCl_3 , 20 $^\circ\text{C}$); $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 8.72 (1H, d, J 4.8 Hz), 8.06 (1H, d, J 7.9 Hz), 7.88 (2H, d, J 8.2 Hz), 7.79 (1H, td, J 7.4, 1.3 Hz), 7.41 (1H, dd, J 6.6, 4.8 Hz), 7.19 (2H, d, J 8.0 Hz), 4.80-4.73 (1H, m), 4.66-4.63 (1H, m), 4.57 (2H, d, J 4.4 Hz), 4.50-4.44 (1H, m), 2.38 (3H, s) ppm; $^{13}\text{C-NMR}$ (100 MHz, CDCl_3) δ 167.7, 165.4, 149.9, 146.2, 143.7, 136.8, 129.8, 129.2, 126.7, 125.9, 124.2, 70.7, 66.0, 65.9, 21.7 ppm; IR 2957, 2901, 1713, 1653, 1255 cm^{-1} ; MP 93-96 $^\circ\text{C}$; ESI-HRMS calcd. for $\text{C}_{17}\text{O}_3\text{N}_2\text{H}_{16}$ 297.1239 $(\text{M}+\text{H})^+$ found m/z 297.1188.

(4S)-2-(Pyridin-2-yl)-4,5-dihydrooxazol-4-ylmethyl 4-methoxybenzoate (23). This was prepared according to the procedure reported above for benzoate **3** using oxazoline **1** (288 mg, 1.6 mmol) in CH_2Cl_2 (5 mL), NEt_3 (0.45 mL, 3.2 mmol) and 4-methoxybenzoyl chloride (0.33 mL, 2.4 mmol). After purification the reaction yielded the mono(oxazoline) ligand **23** as a white solid (372 mg, 78%). $[\alpha]_D^{20} +10.4$ (c 5.20 $\times 10^{-3}$, CHCl_3 , 20 $^\circ\text{C}$); $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 8.56 (1H, d, J 4.7 Hz), 8.19 (2H, d, J 7.8 Hz), 8.01 (2H, d, J 8.6 Hz), 7.85 (1H, td, J 7.7, 1.7 Hz), 7.43 (1H, dd, J 6.6, 4.9 Hz), 6.92 (2H, d, J 8.9 Hz), 4.61-4.45 (5H, m) 3.86 (3H, s, OCH_3) ppm; $^{13}\text{C-NMR}$ (100 MHz, CDCl_3) δ 166.6, 164.6, 149.9, 146.4, 144.0, 136.8, 130.8, 129.2, 127.0, 125.9, 124.2, 70.7, 66.0, 65.9, 55.5 ppm; IR 2959, 2930, 1706, 1603, 863 cm^{-1} ; MP 102-105 $^\circ\text{C}$; ESI-HRMS calcd. for $\text{C}_{17}\text{O}_4\text{N}_2\text{H}_{16}$ 313.1188 $(\text{M}+\text{H})^+$ found m/z 313.1147.

Acknowledgements

We acknowledge the College of Science NUI Galway for studentships provided to N. McC. and Z.C.

References

1. Malacea, R.; Poli, R.; Manoury, E. *Coord. Chem. Rev.* **2010**, *254*, 729.
<https://doi.org/10.1016/j.ccr.2009.09.033>
2. Ohkuma, T.; Ishii, D.; Takeno, H.; Noyori, R. *J. Am. Chem. Soc.* **2000**, *122*, 6510.
<https://doi.org/10.1021/ja001098k>
3. Schnider, P.; Koch, G.; Prétôt, R.; Wang, G.; Bohnen, F. M.; Krüger, C.; Pfaltz, A. *Chem. Eur. J.* **1997**, *3*, 887.
<https://doi.org/10.1002/chem.19970030609>
4. Echeverria, P.-G.; Féraud, C.; Phansavath, P.; Ratovelomanana-Vidal, V. *Catal. Commun.* **2015**, *62*, 95.
<https://doi.org/10.1016/j.catcom.2015.01.012>
5. Uematsu, N.; Fujii, A.; Hashiguchi, S.; Ikariya, T.; Noyori, R. *J. Am. Chem. Soc.* **1996**, *118*, 4916.
<https://doi.org/10.1021/ja960364k>
6. Ou, Z.-M.; Shi, H.-B.; Sun, X.-Y.; Shen, W.-H. *J. Mol. Catal. B: Enzym.* **2011**, *72*, 294.
<https://doi.org/10.1016/j.molcatb.2011.07.004>
7. Brunner, H.; Henrichs, C. *Tetrahedron: Asymmetry* **1995**, *6*, 653.
[https://doi.org/10.1016/0957-4166\(95\)90125-6](https://doi.org/10.1016/0957-4166(95)90125-6)
8. Nishibayashi, Y.; Takei, I.; Uemura, S.; Hidai, M. *Organometallics* **1998**, *17*, 3420.
<https://doi.org/10.1021/om980399o>
9. Malkov, A. V.; Stewart-Liddon, A. J. P.; McGeoch, G. D.; Ramirez-Lopez, P.; Kocovsky, P. *Org. Biomol. Chem.* **2012**, *10*, 4864.
<https://doi.org/10.1039/c2ob25472g>
10. Iwasaki, F.; Onomura, O.; Mishima, K.; Maki, T.; Matsumura, Y. *Tetrahedron Lett.* **1999**, *40*, 7507.
[https://doi.org/10.1016/S0040-4039\(99\)01554-3](https://doi.org/10.1016/S0040-4039(99)01554-3)
11. Iwasaki, F.; Onomura, O.; Mishima, K.; Kanematsu, T.; Maki, T.; Matsumura, Y. *Tetrahedron Lett.* **2001**, *42*, 2525.
[https://doi.org/10.1016/S0040-4039\(01\)00219-2](https://doi.org/10.1016/S0040-4039(01)00219-2)
12. Malkov, A. V.; Mariani, A.; MacDougall, K. N.; Kočovský, P. *Org. Lett.* **2004**, *6*, 2253.
<https://doi.org/10.1021/ol049213+>
13. Malkov, A. V.; Figlus, M.; Stončius, S.; Kočovský, P. *J. Org. Chem.* **2007**, *72*, 1315.
<https://doi.org/10.1021/jo062215i>
14. Zhou, L.; Wang, Z.; Wei, S.; Sun, J. *Chem. Commun.* **2007**, 2977.
<https://doi.org/10.1039/b703307a>
15. Zheng, H.; Deng, J.; Lin, W.; Zhang, X. *Tetrahedron Lett.* **2007**, *48*, 7934.
<https://doi.org/10.1016/j.tetlet.2007.09.064>
16. Guizzetti, S.; Benaglia, M.; Cozzi, F.; Annunziata, R. *Tetrahedron* **2009**, *65*, 6354.
<https://doi.org/10.1016/j.tet.2009.06.015>
17. Gautier, F.-M.; Jones, S.; Li, X.; Martin, S. J. *Org. Biomol. Chem.* **2011**, *9*, 7860.
<https://doi.org/10.1039/c1ob05965c>
18. Pei, D.; Wang, Z.; Wei, S.; Zhang, Y.; Sun, J. *Org. Lett.* **2006**, *8*, 5913.
<https://doi.org/10.1021/ol062633+>
19. Malkov, A. V.; Liddon, A.; Ramirez-Lopez, P.; Bendova, L.; Haigh, D.; Kocovsky, P. *Angew. Chem. Int. Ed.* **2006**, *45*, 1432.
<https://doi.org/10.1002/anie.200503941>
20. Frain, D.; Kirby, F.; McArdle, P.; O'Leary, P. *Synlett* **2009**, 1261.

<https://doi.org/10.1055/s-0029-1216731>

21. Frain, D.; Kirby, F.; McArdle, P.; O'Leary, P. *Tetrahedron Lett.* **2010**, *51*, 4103.
<https://doi.org/10.1016/j.tetlet.2010.05.135>
22. Kirby, F.; Frain, D.; McArdle, P.; O'Leary, P. *Catal. Commun.* **2010**, *11*, 1012.
<https://doi.org/10.1016/j.catcom.2010.05.003>
23. Balavoine, G.; Clinet, J. C.; Lellouche, I. *Tetrahedron Lett.* **1989**, *30*, 5141.
[https://doi.org/10.1016/S0040-4039\(01\)93469-0](https://doi.org/10.1016/S0040-4039(01)93469-0)
24. 24 Çelebi-Ölçüm, N.; Lam, Y.-h.; Richmond, E.; Ling, K. B.; Smith, A. D.; Houk, K. N. *Angew. Chem. Int. Ed.* **2011**, *50*, 11478.
<https://doi.org/10.1002/anie.201105412>
25. Kenworthy, M. N.; Kilburn, J. P.; Taylor, R. J. K. *Org. Lett.* **2004**, *6*, 19.
<https://doi.org/10.1021/ol0360039>
26. Castellano, S.; Kuck, D.; Sala, M.; Novellino, E.; Lyko, F.; Sbardella, G. *J. Med. Chem.* **2008**, *51*, 2321.
<https://doi.org/10.1021/jm7015705>
27. Schulz, M.; Christoffers, J. *Synthesis* **2014**, *46*, 81.
<https://doi.org/10.1055/s-0033-1340044>
28. Motloch, P.; Valterová, I.; Kotora, M. *Adv. Synth. Catal.* **2014**, *356*, 199.
<https://doi.org/10.1002/adsc.201300849>
29. 29 Kim, Y.-A.; Chung, H.-M.; Park, J.-S.; Choi, W.; Min, J.; Park, N.-H.; Kim, K.-H.; Jhon, G.-J.; Han, S.-Y. *J. Org. Chem.* **2003**, *68*, 10162.
<https://doi.org/10.1021/jo034969s>

This paper is an open access article distributed under the terms of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>)