

Supplementary Material

Computational insights into substituent effects on the stability and reactivity of flavylium cation analogs of anthocyanins

Adilson A. Freitas,^{a,*} Karina Shimizu,^a Luis G. Dias,^b and Frank H. Quina^c

^a*Centro de Química Estrutural, Instituto Superior Técnico, Universidade de Lisboa, Lisbon, Portugal*

^b*Departamento de Química, Faculdade de Filosofia, Ciências e Letras de Ribeirão Preto, Universidade de São Paulo, Ribeirão Preto, Brazil*

^c*Departamento de Química Fundamental, Instituto de Química, Universidade de São Paulo, São Paulo, Brazil
Email: adilsondefreitas@tecnico.ulisboa.pt*

Table of Contents

Table S1. Gas phase energies, solvation free energies (in kcal.mol ⁻¹) and pK _h equilibrium constants calculated at T = 25 °C.....	S2
Table S2. Hammett σ parameters for the substituents of the flavylium salts studied.....	S3
Table S3. Electronic energies (in Hartrees) of the species in its normal oxidation state [E(N)], reduced state [E(N+1)] and oxidized state [E(N-1)], Vertical Ionization Potential (VIP) and Vertical Electron Affinity (VEA) in eV, calculated at the mPW1PW91/6-31G(d) level.....	S4
Table S4. Core-Electron Binding Energies (in eV) of some selected compounds compiled from NIST X-ray Photoelectron Spectroscopy Database ^a and values estimated based on the Koopmans' theorem, calculated at the mPW1PW91/6-31G(d) level.....	S8
Table S5. Core-Electron Binding Energies (in eV) of the most relevant centers in flavylium salts, taken from Natural Population Analysis calculated at the mPW1PW91/6-31G(d) level.....	S9
Table S6. Core-Electron Binding Energies (in eV) of the most relevant centers in natural anthocyanins, taken from Natural Population Analysis calculated at the mPW1PW91/6-31G(d) level.....	S10
Table S7. Hammett σ parameters for the substituents of the natural anthocyanins studied.....	S11
Figure S1. Experimental binding energies (orbital 1s) and the corresponding calculated values at the mPW1PW91/6-31G(d) level with solvent effects for (a) carbon, (b) oxygen and (c) nitrogen data on Table S4.....	S12
Figure S2. Representations of (a) pK _h and (b) -log(K _t) as a function of the difference between the core-electron binding energy of C2(1s) of AH ⁺ and the corresponding value for the unsubstituted compound 1 , ΔCEBE; (c) -log(K _i) as a function of the average ΔCEBE of C2'(1s) and C6'(1s) in the C _z form; and (d) pK _a as a function of ΔCEBE of the O(1s) of the oxygen of the hydroxyl groups of the AH ⁺ species. The ΔCEBE values of the flavylium salt species (full symbols) and anthocyanins (open symbols) were determined from Natural Population analysis at the mPW1PW91/6-31G(d) level of theory.....	S13

Table S1. Gas phase energies, solvation free energies (in kcal.mol⁻¹) and pK_h equilibrium constants calculated at T = 25 °C

cpd	ΔG_g^* (AH ⁺) ^a	ΔG_{solv}^* (AH ⁺) ^b	ΔG_{solv}^* (AH ⁺) ^c	ΔG_{solv}^* (AH ⁺) ^d	ΔG_{solv}^* (AH ⁺) ^e	ΔG_{solv}^* (B) ^a	ΔG_{solv}^* (B) ^b	ΔG_{solv}^* (B) ^c	ΔG_{solv}^* (B) ^d	pK _h ^b	pK _h ^c	pK _h ^d	pK _h ^e
1	-144.5	-46.9	-40.3	-45.9	-39.2	-6.82	-6.43	-9.19	-5.98	3.12	2.91	2.67	3.39
2	-141.5	-50.2	-46.3	-50.5	-44.3	-11.0	-12.3	-14.7	-11.1	4.70	5.20	4.20	5.63
3	-141.0	-50.1	-46.2	-50.1	-43.5	-11.0	-12.6	-14.7	-10.9	4.90	5.21	4.24	5.44
4	-137.7	-48.7	-44.7	-48.9	-42.6	-11.1	-11.5	-14.2	-10.9	6.21	7.30	6.14	7.23
5	-139.1	-46.5	-39.7	-44.7	-38.2	-8.64	-8.05	-10.6	-7.16	5.45	5.25	4.74	5.81
6	-142.6	-45.7	-38.9	-44.2	-38.0	-6.51	-6.16	-8.79	-5.72	3.85	3.47	3.11	4.12
7	-140.3	-53.4	-44.2	-51.9	-44.4	-16.2	-12.8	-18.4	-14.3	4.12	4.14	3.38	4.21
8	-140.8	-51.6	-47.5	-52.2	-45.2	-14.1	-15.0	-18.5	-13.1	3.94	4.62	3.12	5.30
9	-138.7	-49.1	-44.9	-48.8	-43.1	-10.9	-12.1	-14.2	-10.8	6.05	6.38	5.40	7.01
10	-152.0	-60.7	-54.2	-60.1	-51.4	-13.8	-14.9	-17.5	-12.7	2.57	1.33	1.47	1.89
11	-129.5	-47.0	-43.5	-46.6	-39.0	-13.4	-14.0	-16.1	-11.5	9.34	10.7	9.14	10.2
12	-134.9	-47.3	-41.0	-45.9	-42.0	-12.4	-10.2	-13.8	-10.9	6.38	7.70	6.30	8.87
13	-142.4	-50.9	-46.9	-51.2	-44.0	-11.1	-12.6	-14.8	-10.9	4.45	4.75	3.99	4.81
14	-137.1	-50.4	-46.1	-49.7	-43.4	-12.7	-13.8	-15.8	-11.8	6.75	7.17	6.03	7.67
MAD^f										0.64	0.68	0.55	0.99
15	-138.2	-53.9	-52.4	-55.0	-48.4	-15.2	-18.4	-19.9	-15.5	6.67	7.59	6.10	7.75
16	-139.3	-57.9	-61.5	-60.7	-55.4	-22.9	-28.1	-29.8	-23.4	3.22	6.34	2.18	6.34
17	-134.0	-51.4	-49.3	-51.7	-46.1	-16.5	-18.5	-20.4	-16.1	6.99	8.30	6.34	8.70
18	-131.4	-50.4	-48.7	-50.3	-44.8	-16.2	-17.9	-19.6	-15.0	8.37	10.4	7.86	10.8
19	-136.1	-50.6	-49.0	-51.2	-45.4	-14.2	-16.2	-18.2	-13.8	6.59	8.27	6.15	8.44
20	-139.8	-46.8	-39.6	-45.0	-38.2	-8.42	-7.57	-10.4	-6.85	5.29	5.00	4.59	5.52
21	-136.5	-55.4	-56.4	-57.5	-51.8	-18.9	-23.3	-24.3	-19.9	6.32	8.19	5.87	8.22

^a $\Delta G_g^* = \Delta G_g^o(AH^+) - \Delta G_g^{o\rightarrow\bullet}$ ^b ΔG_{solv}^* calculated by the PCM2 model and $\Delta\Delta G_{solv}^*$ evaluated employing the experimental⁴⁶ $\Delta G_{solv}^*(OH^-) = -104.6$ kcal.mol⁻¹;^c ΔG_{solv}^* calculated by the IEFPCM HF(UAHF) model and $\Delta\Delta G_{solv}^*$ evaluated with $\Delta G_{solv}^*(OH^-) = -110.52$ kcal.mol⁻¹, estimated at the same level;^d ΔG_{solv}^* calculated by the SMD model with $\Delta G_{solv}^*(OH^-) = -107.37$ kcal.mol⁻¹, estimated at the same level;^e ΔG_{solv}^* calculated by the SM5.4 model and $\Delta\Delta G_{solv}^*$ evaluated with $\Delta G_{solv}^*(OH^-) = -111.87$ kcal.mol⁻¹, estimated at the same level;^f Mean Absolute Deviation.

Table S2. Hammett σ parameters for the substituents of the flavylium salts studied. The σ -values were compiled from reference 16

compound	C3 (σ_m)	C4 (σ_R)	C5 (σ_R)	C6 (σ_m)	C7 (σ_R)	C8 (σ_m)	C4' (σ_R)	$\Sigma\sigma_X$
1	0	0	0	0	0	0	0	0
2	0	0	0	0	-0.7	0	0	-0.7
3	0	0	0	0	0	0	-0.7	-0.7
4	0	-0.18	0	0	-0.7	0	0	-0.88
5	0	0	0	0	0	0	-0.56	-0.56
6	0	0	0	0	0	0	-0.18	-0.18
7	0	0	0	0	0	0	-0.31	-0.31
8	0	0	0	0	-0.7	0.12	0	-0.58
9	0	0	0	0	-0.7	0	-0.18	-0.88
10	0	0	0	0.71	0	0	-0.7	0.01
11	0	0	0	0.12	0	0	-1.85 ^a	-1.73
12	0	0	0	0	0	0	-1.38 ^a	-1.38
13	0	0	0	0.12	0	0	0	0.12
14	0	0	0	0	-0.7	0	-0.56	-1.26
15	0	0	0	0	-0.7	0	-0.7	-1.40
16	0.12	0	-0.7	0	-0.7	0	-0.7	-1.98
17	0	0	-0.56	0	-0.7	0	-0.7	-1.96
18	0	-0.18	-0.7	0	-0.7	0	-0.56	-2.14
19	0	-0.18	-0.7	0	-0.7	0	0	-1.58
20	0	0	0	0	-0.56	0	0	-0.56
21	0	0	-0.7	0	-0.7	0	-0.7	-2.10

^a through-resonance parameter.

Table S3. Electronic energies (in Hartrees) of the species in its normal oxidation state [E(N)], reduced state [E(N+1)] and oxidized state [E(N-1)], Vertical Ionization Potential (VIP) and Vertical Electron Affinity (VEA) in eV, calculated at the *mPW1PW91/6-31G(d)* level. Note that 1 Hartree = 27.2114 eV

compound	E(N)	E(N+1)	E(N-1)	VEA	VIP
AH ⁺ form					
1	-653.1404	-653.2778	-652.8785	3.74	7.13
2	-728.3548	-728.4860	-728.1082	3.57	6.71
3	-728.3554	-728.4860	-728.1158	3.55	6.52
4	-767.6664	-767.7957	-767.4192	3.52	6.73
5	-767.6413	-767.7732	-767.3998	3.59	6.57
6	-692.4511	-692.5869	-692.1954	3.70	6.96
7	-861.1234	-861.2572	-860.8853	3.64	6.48
8	-803.5541	-803.6852	-803.3214	3.57	6.33
9	-767.6653	-767.7951	-767.4222	3.53	6.62
10	-932.8043	-932.9453	-932.5579	3.84	6.70
11	-862.2998	-862.4224	-862.0897	3.34	5.72
12	-708.5036	-708.6256	-708.2874	3.32	5.88
13	-728.3495	-728.4847	-728.1053	3.68	6.65
14	-842.8553	-842.9812	-842.6217	3.43	6.35
15	-803.5694	-803.6942	-803.3377	3.39	6.30
16	-953.9798	-954.1028	-953.7615	3.35	5.94
17	-918.0664	-918.1889	-917.8366	3.33	6.25
18	-957.3749	-957.4940	-957.1438	3.24	6.29
19	-842.8753	-842.9986	-842.6351	3.36	6.53
20	-767.6411	-767.7728	-767.3923	3.58	6.77
21	-878.7800	-878.9018	-878.5504	3.31	6.25

Table S3. (continued)

compound	E(N)	E(N+1)	E(N-1)	VEA	VIP
B form					
1	-729.1045	-729.1456	-728.8868	1.12	5.92
2	-804.3137	-804.3469	-804.1101	0.90	5.54
3	-804.3136	-804.3532	-804.0990	1.08	5.84
4	-843.6233	-843.6544	-843.4195	0.85	5.55
5	-843.6006	-843.6407	-843.3851	1.09	5.86
6	-768.4135	-768.4540	-768.1966	1.10	5.90
7	-937.0838	-937.1251	-936.8675	1.12	5.89
8	-879.5128	-879.5451	-879.3111	0.88	5.49
9	-843.6227	-843.6551	-843.4194	0.88	5.53
10	-1008.774	-1008.865	-1008.551	2.48	6.06
11	-938.2485	-938.2876	-938.0578	1.06	5.19
12	-784.4544	-784.4928	-784.2548	1.05	5.43
13	-804.3105	-804.3521	-804.1071	1.13	5.53
14	-918.8096	-918.8414	-918.6078	0.87	5.49
15	-879.5226	-879.5536	-879.3212	0.84	5.48
16	-1029.937	-1029.951	-1029.753	0.36	5.01
17	-994.0181	-994.0477	-993.8189	0.81	5.42
18	-1033.324	-1033.348	-1033.125	0.67	5.41
19	-918.8280	-918.8534	-918.6285	0.69	5.43
20	-843.6008	-843.6354	-843.3961	0.94	5.57
21	-954.7312	-954.7591	-954.5327	0.76	5.40

Table S3. (continued)

compound	E(N)	E(N+1)	E(N-1)	VEA	VIP
	C _Z form				
1	-729.0942	-729.1700	-728.8777	2.06	5.89
2	-804.3046	-804.3791	-804.0990	2.03	5.59
3	-804.3055	-804.3751	-804.0902	1.89	5.86
4	-843.6123	-843.6845	-843.4059	1.96	5.62
5	-843.5914	-843.6635	-843.3747	1.96	5.90
6	-768.4038	-768.4771	-768.1878	2.00	5.88
7	-937.0749	-937.1484	-936.8589	2.00	5.88
8	-879.5042	-879.5784	-879.2972	2.02	5.63
9	-843.6139	-843.6873	-843.4088	2.00	5.58
10	-1008.767	-1008.856	-1008.537	2.43	6.27
11	-938.2414	-938.3069	-938.0447	1.78	5.35
12	-784.4482	-784.5133	-784.2436	1.77	5.57
13	-804.2994	-804.3746	-804.0999	2.05	5.43
14	-918.8022	-918.8727	-918.5984	1.92	5.55
15	-879.5155	-879.5849	-879.3119	1.89	5.54
16	-1029.927	-1029.987	-1029.739	1.63	5.13
17	-994.0112	-994.0797	-993.8086	1.86	5.51
18	-1033.317	-1033.382	-1033.112	1.75	5.57
19	-918.8195	-918.8896	-918.6141	1.91	5.59
20	-843.5914	-843.6666	-843.3853	2.05	5.61
21	-954.7246	-954.7926	-954.5223	1.85	5.51

Table S3. (continued)

compound	E(N)	E(N+1)	E(N-1)	VEA	VIP
	C _E form				
1	-729.1064	-729.1924	-728.8869	2.34	5.97
2	-804.3178	-804.3990	-804.1101	2.21	5.65
3	-804.3178	-804.3993	-804.1002	2.22	5.92
4	-843.6148	-843.6915	-843.4069	2.09	5.66
5	-843.5765	-843.6854	-843.3842	2.96	5.23
6	-768.4160	-768.5008	-768.1970	2.31	5.96
7	-937.0871	-937.1723	-936.8688	2.32	5.94
8	-879.5171	-879.5991	-879.3083	2.23	5.68
9	-843.6273	-843.7075	-843.4201	2.18	5.64
10	-1008.777	-1008.870	-1008.548	2.53	6.23
11	-938.2540	-938.3326	-938.0581	2.14	5.33
12	-784.4604	-784.5384	-784.2567	2.12	5.54
13	-804.3115	-804.3979	-804.1078	2.35	5.54
14	-918.8155	-918.8934	-918.6099	2.12	5.59
15	-879.5289	-879.6059	-879.3230	2.10	5.60
16	-1029.936	-1030.007	-1029.742	1.94	5.28
17	-994.0235	-994.0987	-993.8204	2.05	5.53
18	-1033.319	-1033.389	-1033.112	1.91	5.62
19	-918.8214	-918.8952	-918.6138	2.01	5.65
20	-843.6043	-843.6864	-843.3960	2.23	5.67
21	-954.7372	-954.8117	-954.5342	2.03	5.52

Table S4. Core-Electron Binding Energies (in eV) of some selected compounds compiled from NIST X-ray Photoelectron Spectroscopy Database^a and values estimated based on the Koopmans' theorem, calculated at the *mPW1PW91/6-31G(d)* level. The asterisk (*) in XPS formula indicates the specific atom which is being observed

Compound	XPS Formula	Experimental C(1s)	Calculated C(1s)
benzene	(C*H)	284.8	274.8
toluene	C ₆ H ₅ C*H ₃	284.7	274.9
toluene	C ₆ H ₅ CH ₃ (C*-CH ₃)	285.1	274.9
toluene	C ₆ H ₅ CH ₃ (C*H)	285.0	274.8
naphthalene	(C*H)	284.3	274.8
trimethoxymethane	HC*(OCH ₃) ₃	289.7	279.8
benzonitrile	(C*H)	285.4	275.5
acetonitrile	C*H ₃ CN	286.3	276.4
acetonitrile	CH ₃ C*N	287.2	276.8
methanol	C*H ₃ OH	286.8	276.7
phenol	C ₆ H ₅ OH (C*H)	284.9	274.7
phenol	C ₆ H ₅ OH (C*-OH)	286.7	276.7
aniline	C ₆ H ₅ NH ₂ (C*H)	284.8	274.5
aniline	C ₆ H ₅ NH ₂ (C*-NH ₂)	285.8	275.8
hexafluorobenzene	(C*F)	288.7	278.9
anisole	C ₆ H ₅ OCH ₃ (C*H)	284.9	274.9
acetophenone	C ₆ H ₅ C(=O)CH ₃ (C*H)	285.1	275.1
nitrobenzene	C ₆ H ₅ NO ₂ (C*H)	285.3	275.5
Compound	XPS Formula	Experimental O(1s)	Calculated O(1s)
methanol	CH ₃ O*H	539.1	518.4
2-propanol	(O*H)	538.5	517.9
cyclohexanol	(O*H)	538.4	517.8
formic acid	HC(=O)O*H	540.7	519.8
trifluoroacetic acid	CF ₃ C(=O)O*H	541.3	520.3
phenol	C ₆ H ₅ O*H	539.2	518.6
methyl formate	HC(=O)O*-CH ₃	539.9	519.8
anisole	(C-O*-C)	538.7	518.7
acetophenone	(C=O*)	532.3	516.1
nitrobenzene	C ₆ H ₅ NO* ₂	533.6	518.9
Compound	XPS Formula	Experimental N(1s)	Calculated N(1s)
benzonitrile	(CN*)	399.2	385.4
acetonitrile	(CN*)	399.4	385.3
aniline	C ₆ H ₅ N*H ₂	400.3	387.6
nitrobenzene	C ₆ H ₅ N*O ₂	405.9	394.8
trimethylamine N-oxide	(CH ₃) ₃ N*O	403.0	391.5

^a NIST X-ray Photoelectron Spectroscopy Database, NIST Standard Reference Database Number 20, National Institute of Standards and Technology, Gaithersburg MD, 20899 (2000).

Table S5. Core-Electron Binding Energies (in eV) of the most relevant centers in flavylium salts (see the structure in Scheme 1), taken from Natural Population Analysis calculated at the *mPW1PW91/6-31G(d)* level

compound	OH(1s)	AH ⁺ form		C _Z form	
		O1(1s)	C2(1s)	C2'(1s)	C6'(1s)
1	—	520.99	279.09	274.91	275.01
2	519.50	520.82	278.87	274.88	274.98
3	519.35	520.79	278.92	274.95	275.08
4	519.46	520.75	278.81	274.87	274.97
5	—	520.83	278.96	274.95	275.09
6	—	520.94	279.05	274.86	274.97
7	—	520.84	278.96	274.98	275.06
8	519.55	520.86	278.89	274.88	274.98
9	519.48	520.78	278.84	274.83	274.94
10	519.49	521.02	279.16	275.01	275.13
11	519.08	520.54	278.62	274.77	274.87
12	—	520.51	278.64	274.79	274.90
13	519.23	520.97	278.99	274.91	275.01
14	519.44	520.73	278.78	274.95	275.09
15	519.42	520.69	278.74	274.92	275.05
16	519.33	520.69	278.70	274.97	275.04
17	519.44	520.65	278.69	274.91	275.04
18	519.37	520.54	278.57	274.94	275.08
19	519.43	520.63	278.65	274.87	274.98
20	—	520.89	278.90	274.88	274.98
21	519.40	520.63	278.67	274.91	275.04

Table S6. Core-Electron Binding Energies (in eV) of the most relevant centers in natural anthocyanins (see the structure in Scheme 1), taken from Natural Population Analysis calculated at the *mPW1PW91/6-31G(d)* level

compound	AH ⁺ form			C _Z form	
	OH(1s)	O1(1s)	C2(1s)	C2'(1s)	C6'(1s)
malvin	519.38	520.84	278.82	274.98	274.98
cianin	519.33	520.79	278.75	274.88	274.91
oenin	519.39	520.81	278.80	274.98	274.91
peonin	519.38	520.82	278.77	274.97	274.97
kuromarin	519.37	520.76	278.73	274.89	274.91
callistephin	519.38	520.77	278.73	274.99	275.03
hibiscin	519.32	520.79	278.82	274.91	274.91
myrthillin	519.38	520.77	278.73	274.78	274.78
petunidin 3- <i>O</i> -glucoside	519.38	520.79	278.76	274.86	274.84
oxycocccyanin	519.38	520.79	278.75	274.97	274.97

Table S7. Hammett σ parameters for the substituents of the natural anthocyanins studied. The σ -values were compiled from reference 16

compound ^a	C3 (σ_m)	C5 (σ_R)	C7 (σ_R)	C3' (σ_m)	C4' (σ_R)	C5' (σ_m)	$\Sigma\sigma_x$
malvin	-0.1	-0.18	-0.7	0.12	-0.7	0.12	-1.44
cianin	-0.1	-0.18	-0.7	0.12	-0.7	0	-1.56
oenin	-0.1	-0.7	-0.7	0.12	-0.7	0.12	-1.96
peonin	-0.1	-0.18	-0.7	0.12	-0.7	0	-1.56
kuromarin	-0.1	-0.7	-0.7	0.12	-0.7	0	-2.08
callistephin	-0.1	-0.7	-0.7	0	-0.7	0	-2.20
hibiscin	-0.1	-0.7	-0.7	0.12	-0.7	0.12	-1.96
myrthillin	-0.1	-0.7	-0.7	0.12	-0.7	0.12	-1.96
petunidin 3-O-glucoside	-0.1	-0.7	-0.7	0.12	-0.7	0.12	-1.96
oxycocccyanin	-0.1	-0.7	-0.7	0.12	-0.7	0	-2.08

^a O-glucoside and O-sambubioside units were represented with electronic and steric parameters (E_S) of the *t*-butyl group ($\sigma_R = -0.18$; $\sigma_m = -0.1$; $E_S = -1.78$)




Figure S1. Experimental binding energies (orbital 1s) and the corresponding calculated values at the mPW1PW91/6-31G(d) level with solvent effects for (a) carbon, (b) oxygen and (c) nitrogen data on Table S4.




Figure S2. Representations of (a) pK_h and (b) $-\log(K_t)$ as a function of the difference between the core-electron binding energy of C2(1s) of AH^+ and the corresponding value for the unsubstituted compound **1**, ΔCEBE ; (c) $-\log(K_i)$ as a function of the average ΔCEBE of C2'(1s) and C6'(1s) in the C_z form; and (d) pK_a as a function of ΔCEBE of the O(1s) of the oxygen of the hydroxyl groups of the AH^+ species. The ΔCEBE values of the flavylium salt species (full symbols) and anthocyanins (open symbols) were determined from Natural Population analysis at the *mPW1PW91/6-31G(d)* level of theory.