

## The synthesis of 7,9-dimethoxy-3-propyl-3,4-dihydro-1*H*-benzo[*g*]isochromene-1,5,10-trione: A potential monomer for the synthesis of the natural product xylindein

Md. Firoj Hossain, Andreas Lemmerer,<sup>II</sup> and Charles B. de Koning\*

Molecular Sciences Institute, School of Chemistry, University of the Witwatersrand, PO Wits 2050, Johannesburg, South Africa

Email: [Charles.deKoning@wits.ac.za](mailto:Charles.deKoning@wits.ac.za)


Received 07-24-2020

Accepted 08-22-2020

Published on line 09-09-2020

### Abstract


A novel synthesis of a 3-propyl-substituted-benzo[*g*]isochromene quinone, a potential monomer of the natural product xylindein, was accomplished in 9 steps (overall yield of 8.2%) from 2,4-dimethoxybenzaldehyde. Key steps included the use of a cross-metathesis reaction in which ethyl-3-allyl-4-(benzyloxy)-1,6,8-trimethoxy-2-naphthoate was converted into ethyl-4-(benzyloxy)-1,6,8-trimethoxy-3-(4-oxopent-2-enyl)-2-naphthoate as a mixture of (*E*)- and (*Z*)-isomers. Following an oxa-Michael addition reaction, a racemic mixture of the desired product, 5-(benzyloxy)-7,9,10-trimethoxy-3-(2-oxopropyl)-3,4-dihydro-1*H*-benzo[*g*]isochromen-1-one, the basic xylindein lactone skeleton, was obtained.



**Keywords:** Xylindein, benzo[*g*]isochromene quinone, cross-metathesis, oxa-Michael addition

## Introduction

Anhydrofusarubin lactone (**1**) and 5-methoxy-3,4-dehydrosemixanthomegnin (**2**) are examples of lactone-containing aromatic compounds. Both compounds belong to an expanding class of quinone-containing compounds (Figure 1).<sup>1</sup> Anhydrofusarubin lactone (**1**)<sup>2</sup> is found in several fungi such as *Nectria haematococca* and *Fusarium solani*, while 5-methoxy-3,4-dehydrosemixanthomegnin (**2**)<sup>3</sup> has been isolated from *Paepalanthus latipes* Silveira (family Eriocaulaceae). Related aromatic compounds, but not quinones, are paepalantine (**3**) and 9-*O*-methyl paepalantine (**4**), as well as the dimer of paepalantine (**5**), have been isolated from a related Eriocaulaceae species *Paepalanthus bromelioides*<sup>4</sup> and *Paepalanthus vellozioides*.<sup>5</sup> A more complex, related dimeric-fused lactone-containing aromatic compound possessing a stereogenic center at C-3 is xylindein (**6**), produced by fungi from genus *Chlorociboria aeruginosa* (Figure 2). This pigment is known for the green staining of wood. The artificial coloring of wood with *C. aeruginosa* is a patented process, and oak staining using this procedure is used in decorative woodworking such as *Tunbridge ware*.<sup>6</sup>



**Figure 1.** Examples of naturally occurring lactone-containing aromatic compounds.

The structure of xylindein was elucidated, independently, by Todd et al.<sup>7</sup> and by Edwards and Kale.<sup>8</sup> More recently, the absolute configuration and tautomeric structure of xylindein have been described.<sup>9</sup> An interesting potential application of xylindein is that it has been investigated for use as a sustainable organic agent for optical- and electronic-property applications.<sup>10</sup>


Two research groups have attempted the synthesis of xylindein. Giles et al.<sup>11</sup> and Green et al.<sup>12</sup> have postulated that the possible biosynthetic precursor to xylindein was the lactone-containing quinone **7** (Figure 2). While the racemic quinone **7** could be synthesized in very poor yields, the authors concluded that “new synthetic strategies to the elusive  $\delta$ -lactone **7** will have to be sought.”<sup>12</sup>

Gill and Donner<sup>13</sup> postulated that the naphthol **8**, or even the lactone deficient aromatic compound **9**, might be better building blocks for the formation of xylindein.<sup>14</sup> The use of these building blocks was postulated on the basis of previous literature. In this research, the extended quinone, xylaphin, was prepared from a *pyran*-containing aromatic compound rather than a *lactone* precursor. Naphthol **8** was synthesized by Gill using Staunton-Weinreb annulation methodology, while **9** was synthesized by Donner using the Diels-Alder

reaction as a key step. None of these approaches, however, allowed for the synthesis of the target compound, xylindein.

In our laboratories, we have explored the synthesis of lactone-containing aromatic compounds. This has resulted in the synthesis of 5-methoxy-3,4-dehydrosemixanthomegnin (**2**) and 9-*O*-methylpaepalantine (**5**), a methoxy derivative of paepalantine (**3**).<sup>15</sup> Hence, we believed that we would be able to synthesize the quinone-containing lactone **7**. In principle, dehydration of **7** should result in the formation of xylindein.

In this paper, we disclose novel methodology we have developed for the racemic synthesis of 7,9-dimethoxy-3-propyl-3,4-dihydro-1*H*-benzo[*g*]isochromenene-1,5,10-trione (**10**), a potential monomer *en route* to the synthesis of xylindein (**6**).




**Figure 2.** Postulated monomeric aromatic building blocks for the assembly of xylindein.

## Results and Discussion

We have previously described the synthesis of the trioxygenated naphthalene **11**<sup>16a</sup> as well as related PIFA-mediated methodology<sup>16b</sup> for the synthesis of the correctly substituted 1,4,6,8-tetraoxygenated naphthalene nucleus **12** of the xylindein monomer. We envisaged that the ester at C-2 and the allyl substituent at C-3 of the naphthalene **12** would allow for possible synthetic routes for the synthesis of the desired isochromenone moiety of **10**.

2,4-Dimethoxybenzaldehyde was treated under Stobbe reaction conditions with diethyl succinate in the presence of *t*-BuOH and *t*-BuOK to afford the intermediate half-ester, which was subjected to Ac<sub>2</sub>O and NaOAc to afford the required naphthalene, ethyl 4-acetoxy-6,8-dimethoxynaphthalene-2-carboxylate, along with a minor product, *t*-butyl-4-acetoxy-6,8-dimethoxy-2-naphthoate (see experimental). Following a further synthetic step described in the literature,<sup>16a</sup> the desired trioxygenated naphthalene **11** containing a C-allyl substituent was furnished in good yield. Using the PIFA-mediated Kozłowski conditions<sup>17</sup> allowed for the introduction of the required methoxy substituent at C-1 of **11** to afford **12** in good yield of 75% (Scheme 1). It

now remained to protect the naphthol **12** with an appropriate protecting group that would allow for selective removal at a later stage of the synthesis. *O*-Benzylation of **12** afforded **13** in excellent yield.





**Scheme 1.** Reagents and Conditions: (i) (a) PIFA, MeOH, rt, 15 min, (b) <sup>t</sup>BuOK, EtOH, 20 min, 75%; (ii) PhCH<sub>2</sub>Br, K<sub>2</sub>CO<sub>3</sub>, acetone, 95%; (iii) Grubbs II catalyst, methyl vinyl ketone, CH<sub>2</sub>Cl<sub>2</sub>, reflux, 15 min, 70%; (iv) BF<sub>3</sub>·OEt<sub>2</sub>, 0 °C, 64%.

The next step in the synthesis was the construction of the appropriate 3-propyl-substituted chromanone ring *via* a cross metathesis reaction of **13** with methyl vinyl ketone in the presence of the Grubbs II catalyst. This resulted in the formation of the  $\alpha,\beta$ -unsaturated carbonyl compound **14** as a mixture of (*E*)- and (*Z*)-isomers. The Michael acceptor **14** was exposed to BF<sub>3</sub>·OEt<sub>2</sub> at 0 °C to give the desired isochromene-1-one **15** in good yield. Treatment of **14** with a 10% potassium hydroxide solution in a mixture of water and ethanol resulted in the formation of anthracene **16** in a poor yield of 20%.<sup>18</sup>

The next step required the removal of the ketone carbonyl of **15**. The thioacetal **17** was readily formed by treatment of **15** with ethanedithiol under acidic conditions as shown in Scheme 2 (structure confirmed by X-ray crystallography<sup>19</sup>), along with a minor product, which indicated that one of the methyl ether groups was no longer present, and had resulted in the formation of naphthol **17a**. Our basis for suggesting that the 10-hydroxynaphthalene **17a** had formed was that a signal for the hydroxyl H was observed at  $\delta$  12.99 in the <sup>1</sup>H NMR spectrum, indicating hydrogen bonding to the adjacent lactone carbonyl.

The conversion of the thioacetal **17** to liberate the propyl substituent was the next step. As the thioacetal **17** also contained an *O*-benzyl ether, we believed that reaction conditions could be manipulated to efficiently afford the trimethoxynaphthol, 5-hydroxy-7,9,10-trimethoxy-3-propyl-3,4-dihydro-1*H*-benzo[*g*]isochromen-1-one that would facilitate the oxidation step to the required naphthoquinone **10**. Thioacetal **17** was exposed to Raney nickel in refluxing ethanol for 72 h. Much to our surprise, not only was the thioacetal no longer present, but the methoxy substituent at C-10 that we had previously introduced was no longer evident. As expected, however, under the reaction conditions, the *O*-benzyl substituent was no longer present, and the C-5 naphthol, 5-hydroxy-7,9-dimethoxy-3-propyl-3,4-dihydro-1*H*-benzo[*g*]isochromen-1-one (**18**) had formed. Evidence for this was that no hydrogen-bonded phenol was observed in the <sup>1</sup>H NMR spectrum of **18**, and the proton at C-10 of the <sup>1</sup>H NMR spectrum which appeared at  $\delta$  8.65 seemed to be incorrect for the C-10 naphthol. In fact, if the regioisomer, 10-hydroxy-7,9-dimethoxy-3-propyl-3,4-dihydro-1*H*-benzo[*g*]isochromen-

1-one, had been formed, the  $^1\text{H}$  NMR spectrum would show a more shielded singlet at about  $\delta$  7. This was corroborated by examining the  $^1\text{H}$  NMR of a related product, 10-hydroxy-7,9-dimethoxy-3-methyl-1H-benzo[*g*]isochromen-1-one, where the proton attached to C-5 was observed at 6.92 in the  $^1\text{H}$  NMR spectrum.<sup>20</sup> Therefore, the isochromenone **18** containing a 5,7,9-trihydroxynaphthalene had been formed. The  $^1\text{H}$  NMR spectrum showed one aromatic singlet at  $\delta$  8.65, as well as two *meta*-coupled doublets at  $\delta$  6.93 (1H, d, *J* 1.9 Hz) and 6.47 (1H, d, *J* 1.9 Hz) while, in the 135 DEPT spectrum, there were three aromatic C-H signals at  $\delta$  119.78, 98.13 and 91.5. Further evidence for the formation of the naphthol **18** was obtained from HRMS (calcd. for  $\text{C}_{18}\text{H}_{21}\text{O}_5$  ( $\text{M}+\text{H}$ )<sup>+</sup> 317.1390, found 317.1390). Undeterred by this result, we exposed the naphthol **18** to CAN which furnished the desired quinone **10** in good yield.



**Scheme 2. Reagents and Conditions:** (i) ethanedithiol, p-TsOH, toluene, reflux, 18 h, 70%; (ii) Raney Ni, EtOH, reflux, 72 h, 79%; (iii) CAN, MeCN, 0 °C, 30 min, 80%.

## Conclusions

Using 2,4-dimethoxybenzaldehyde as the starting material, we have been able to develop a novel synthesis of a 3-propyl-substituted benzo[*g*]isochromenone quinone, a potential monomer of the natural product xylindein, in 9 steps with an overall yield of 8.2%. Currently, we are exploring the utilization of this quinone as the monomer unit for the assembly of xylindein. As the methoxy substituent at C-10 was unexpectedly lost in the conversion of **17** into **18**, attempts are being made to conduct the same synthetic steps commencing with compound **13**, lacking the methoxy substituent, however, at C-1.

## Experimental Section

**General.**  $^1\text{H}$  NMR and  $^{13}\text{C}$  NMR spectra were recorded on a Bruker AVANCE 300 spectrometer. All chemical shift values are reported in parts per million (ppm) referenced against TMS which is given an assignment of zero ppm. Coupling constants (*J*-values) are given in Hertz (Hz). All mass spectroscopy data were collected on a Waters Acquity UPLC system coupled to a Waters HDMS G1 QTOF mass spectrometer. UPLC settings: Analytical column: BEH C18 150 × 2.1 mm; Column temperature: 60 °C; Mobile phase: 90% water (0.1% formic acid): 5% acetonitrile; Flow rate: 0.4 mL/min. MS settings: Mode: VTOF; Ionisation: ESIPos and ESINeg; Scan range: 100-1000 Da; Scan speed: 0.1 second; Run time: 10 mins. Infrared spectra were recorded on a Bruker Tensor 27 standard system spectrometer with diamond ATR attachment. Macherey-Nagel Kieselgel 60

(particle size 0.063-0.200mm) was used for conventional silica gel column chromatography with various EtOAc and hexane mixtures as the mobile phase. TLC was performed on aluminum-backed Macherey-Nagel Alugram Sil G/UV254 plates pre-coated with 0.25mm silica gel 60. The Bruker D8 VENTURE PHOTON CMOS area detector diffractometer, equipped with a graphite monochromated Mo K $\alpha_1$  radiation (50 kV, 30 mA), was used to collect all the intensity data. The program *SAINT+*, v. 6.02<sup>21</sup> was used to reduce the data, and the program *SADABS* was used to make corrections to the empirical absorptions. Space group assignments were made using *XPREP*<sup>21</sup> on all compounds. In all cases, the structures were solved in the *WinGX*<sup>22</sup> Suite of programs by direct methods using *SHELXS-97*<sup>23</sup> and refined using full-matrix least-squares/difference Fourier techniques on  $F^2$  using *SHELXL-97*.<sup>23</sup> All non-hydrogen atoms were refined anisotropically. All C-H hydrogen atoms were placed at idealized positions and refined as riding atoms with isotropic parameters 1.2 or 1.5 times those of the 'heavy' atoms to which they are attached. Diagrams and publication material were generated using *ORTEP-3*,<sup>24</sup> and *PLATON*.<sup>25</sup> Experimental details of the X-ray analyses are given for *t*-butyl-4-acetoxy-6,8-dimethoxy-2-naphthoate in the experimental section and for 5-(benzyloxy)-7,9,10-trimethoxy-3-(2-methyl-1,3-dithiolan-2-yl)methyl-3,4-dihydro-1*H*-benzo[*g*]isochromen-1-one (**17**) in the experimental section.

**Ethyl-4-acetoxy-6,8-dimethoxynaphthalene-2-carboxylate and *t*-butyl-4-acetoxy-6,8-dimethoxy-2-naphthoate.** To a solution of 2,4-dimethoxybenzaldehyde (5.45 g, 32.8 mmol) and diethyl succinate (8.18 mL, 49.2 mmol) in dry *t*-BuOH (80 mL), stirred under N<sub>2</sub>, was added *t*-BuOK (5.52 g, 49.2 mmol). The reaction mixture was heated under reflux for 2 h and then allowed to cool to rt, poured onto ice and made acidic to pH 3.0 with conc. aqueous HCl. The product precipitated before it was dissolved and extracted with ethyl acetate (2 × 100 mL). The combined extracts were dried with MgSO<sub>4</sub>, filtered through Celite, and the filtrate was concentrated *in vacuo*. The resultant oily residue and anhydrous NaOAc (6.72 g, 81.9 mmol) were dissolved in Ac<sub>2</sub>O (100 mL) and refluxed for 3 h before being allowed to cool. The remaining Ac<sub>2</sub>O was removed *in vacuo*, H<sub>2</sub>O (100 mL) was added, and the product was extracted with EtOAc (3 × 100 mL). Drying of the combined EtOAc extracts (MgSO<sub>4</sub>), followed by filtration through Celite, and removal of the solvent *in vacuo*, gave a brown semi-solid which was purified by silica gel column chromatography (28% ethyl acetate-hexane). Initially, a minor product, *t*-butyl-4-acetoxy-6,8-dimethoxy-2-naphthoate was eluted and subsequently obtained as a yellow crystalline solid (1.14 g, 10%), mp 139-141 °C. IR (solid,  $\nu_{\max}$  cm<sup>-1</sup>): 1760, 1702. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>):  $\delta$  (ppm) 8.74 (1H, d, *J* 1.6), 7.74 (1H, d, *J* 1.6), 6.65 (1H, d, *J* 2.0), 6.52 (1H, d, *J* 2.0), 3.97 (3H, s), 3.91 (3H, s), 2.46 (3H, s), 1.62 (9H, s). <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>):  $\delta$  (ppm) 169.3, 165.4, 160.8, 157.9, 145.4, 131.0, 126.1, 123.0, 122.3, 119.2, 98.5, 91.6, 81.1, 55.8, 55.4, 28.3 (× 3), 21.0. Crystal data for *t*-butyl-4-acetoxy-6,8-dimethoxy-2-naphthoate, assigned CCDC no. 1030315, C<sub>19</sub>H<sub>22</sub>O<sub>6</sub>:  $M_r$  346.37 g.mol<sup>-1</sup>; crystal dimensions (mm) 0.40 × 0.34 × 0.23; crystal system, monoclinic; space group, *C2/c*; unit cell dimensions and volume, *a* = 17.2795(11) Å, *b* = 9.8328(4) Å, *c* = 22.8241(9) Å,  $\alpha$  = 90°,  $\beta$  = 104.799(2)°,  $\gamma$  = 90°, *V* = 3749.3(3) Å<sup>3</sup>, *Z* = 8; calculated density  $\rho_{\text{calcd}}$ , 1.227 Mg/m<sup>3</sup>; linear absorption coefficient,  $\mu$  = 0.091 mm<sup>-1</sup>; radiation and wavelength, MoK $\alpha_1$  = 0.71073 Å; temperature of measurement, 173(2) K,  $2\theta_{\max}$  28.00°; 24655 reflections measured, 4512 unique reflections, 3293 observed reflections [*I* > 2 $\sigma$ (*I*)]; *R*<sub>int</sub> = 0.0402; *R*<sub>1</sub> [*I* > 2.0 $\sigma$ (*I*)] = 0.0539, *wR2* [all] = 0.1536, GoF = 1.016, refined on  $F^2$ ; residual electron density, 0.456 and -0.369 eÅ<sup>-3</sup>.

The major desired product, ethyl 4-acetoxy-6,8-dimethoxynaphthalene-2-carboxylate (8.10 g, 77%) was obtained as a white solid. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>):  $\delta_H$  (ppm) 8.79 (1H, d, *J* 1.1), 7.81 (1H, d, *J* 1.1), 6.66 (1H, d, *J* 1.9), 6.51 (1H, d, *J* 1.9), 4.42 (2H, q, *J* 7.1), 3.98 (3H, s), 3.91 (3H, s), 2.46 (3H, s), 1.42 (3H, t, *J* 7.1). <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>):  $\delta_C$  (ppm) 169.3, 166.3, 161.0, 157.9, 145.4, 133.2, 124.4, 123.4, 122.2, 119.1, 98.6, 91.7, 61.0, 55.8, 55.4, 20.9, 14.4.<sup>16a</sup>

Ethyl-4-acetoxy-6,8-dimethoxynaphthalene-2-carboxylate was converted into ethyl-3-allyl-4-hydroxy-6,8-dimethoxynaphthalene-2-carboxylate **11** in a reproducible yield of 75% using a known literature procedure.<sup>15,16a</sup>

**Ethyl-3-allyl-4-hydroxy-1,6,8-trimethoxynaphthalene-2-naphthoate (12).** Ethyl-3-allyl-4-hydroxy-6,8-dimethoxynaphthalene-2-carboxylate **11** (6.02 g, 16.85 mmol) was dissolved in MeOH (100 mL) in a round-bottomed flask. PIFA (9.4 g, 21.90 mmol) was added and the reaction was stirred for 15 min at rt. A saturated aqueous NaHCO<sub>3</sub> solution was added until effervescence ceased and then the MeOH was removed *in vacuo* to leave only an aqueous medium. The aqueous layer was extracted with ethyl acetate (3 × 70 mL). The organic layers were combined, dried over anhydrous MgSO<sub>4</sub>, filtered, and concentrated under reduced pressure. The residue was then treated with an ethanolic solution of <sup>t</sup>BuOK (10.4 g, 92.69 mmol) with vigorous stirring for 20 min at rt. The reaction was then quenched with the addition of a saturated aqueous NH<sub>4</sub>Cl solution (30 mL). Ethyl acetate (100 mL) was added and the resulting upper organic layer was removed, dried over anhydrous MgSO<sub>4</sub>, filtered, and concentrated *in vacuo*. Column chromatography (10% ethyl acetate-hexane) of the residue eventually afforded **12** as a yellow solid (4.37 g, 75%). mp 128-130 °C. IR (solid,  $\nu_{\max}$  cm<sup>-1</sup>): 3243, 1685. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>):  $\delta$  (ppm) 7.05 (1H, br s), 6.50 (1H, br s), 5.95 (1H, br s), 5.75 (1H, s), 5.21 (1H, s), 5.15 (1H, s), 4.39 (2H, q, *J* 7.1), 3.89 (3H, s), 3.81 (6H, s), 3.41 (2H, br s), 1.36 (3H, t, *J* 7.1). <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>):  $\delta$  (ppm) 168.6, 158.7, 157.4, 147.5, 145.4, 135.4, 129.5, 124.7, 116.9, 115.1, 99.4, 92.9, 63.9, 61.3, 56.0, 55.3, 52.2, 32.7, 14.3. HRMS (ESI) *m/z* calcd for C<sub>19</sub>H<sub>23</sub>O<sub>6</sub> (M+H)<sup>+</sup> 347.1493, found 347.1493.<sup>15</sup>

**Ethyl-3-allyl-4-(benzyloxy)-1,6,8-trimethoxy-2-naphthoate (13).** To a solution of the phenol **12** (1.70 g, 4.40 mmol) in acetone (60 mL) was added benzyl bromide (0.78 mL, 6.60 mmol) and K<sub>2</sub>CO<sub>3</sub> (911 mg, 6.60 mmol). The mixture was heated at reflux under N<sub>2</sub> for 18 h. After cooling to rt, the mixture was filtered through Celite and the filtrate concentrated *in vacuo*. The resultant residue was purified by silica gel column chromatography (7% ethyl acetate-hexane) to afford the product **13** as a pale liquid (1.82 g, 95%). IR (liquid,  $\nu_{\max}$  cm<sup>-1</sup>): 2937, 1721, 1618, 1580, 1342. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>):  $\delta$  (ppm) 7.56-7.37 (5H, m), 7.00 (1H, d, *J* 2.2), 6.53 (1H, d, *J* 2.2), 6.07-5.94 (1H, m), 5.12-5.04 (2H, m), 4.97 (2H, s), 4.41 (2H, q, *J* 7.1), 3.96 (3H, s), 3.88 (3H, s), 3.78 (3H, s), 3.65 (1H, t, *J* 1.5), 3.63 (1H, t, *J* 1.5), 1.41 (3H, t, *J* 7.1). <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>):  $\delta$  (ppm) 168.1, 159.3, 157.9, 150.6, 148.3, 137.6, 136.3, 132.9, 128.7 (× 2), 128.1, 127.7 (× 2), 126.9, 125.1, 116.0, 115.4, 99.3, 93.6, 75.9, 63.9, 61.2, 56.1, 55.2, 32.0, 14.3. HRMS (ESI) *m/z* calcd for C<sub>26</sub>H<sub>29</sub>O<sub>6</sub> (M+H)<sup>+</sup> 437.1968, found 437.1968.

**Ethyl-4-(benzyloxy)-1,6,8-trimethoxy-3-(4-oxopent-2-enyl)-2-naphthoate (14).** To a solution of **13** (100 mg, 0.23 mmol) in anhydrous, degassed CH<sub>2</sub>Cl<sub>2</sub> (10 mL), methyl vinyl ketone (0.03 mL, 0.34 mmol) was added as well as the Grubbs II catalyst (3.9 mg, 0.005 mmol). The reaction mixture was then refluxed for 15 min. Following removal of the solvent, the crude mass obtained was purified by silica gel column chromatography (40% ethyl acetate-hexane) to afford the  $\alpha,\beta$ -unsaturated carbonyl compound **14** (77 mg, 70%) as a white solid. mp 91-93 °C. IR (solid,  $\nu_{\max}$  cm<sup>-1</sup>): 1728, 1670, 1658. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>):  $\delta$  (ppm) 7.50-7.36 (5H, m), 6.96 (1H, d, *J* 2.3), 6.92-6.80 (1H, m), 6.54 (1H, d, *J* 2.3), 6.05-6.00 (1H, m), 4.94 (2H, s), 4.36 (2H, q, *J* 7.1), 3.96, (3H, s), 3.86 (3H, s), 3.77 (3H, s), 3.71 (1H, d, *J* 1.6), 3.69 (1H, d, *J* 1.7), 2.17 (3H, s), 1.36 (3H, t, *J* 7.1). <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>):  $\delta$  (ppm) 198.4, 167.9, 159.6, 158.0, 150.95, 148.5, 145.4, 137.2, 132.8, 132.2, 128.8, 128.3, 127.6, 124.8, 124.7, 115.8, 99.6, 93.6, 77.3, 76.1, 63.9, 61.3, 56.2, 55.2, 52.2, 30.8, 26.8, 14.3. HRMS (ESI) *m/z* calcd for C<sub>28</sub>H<sub>31</sub>O<sub>7</sub> (M+H)<sup>+</sup> 479.2072, found 479.2072.

**5-(Benzyloxy)-7,9,10-trimethoxy-3-(2-oxopropyl)-3,4-dihydro-1H-benzo[g]isochromen-1-one (15).** To a solution of the  $\alpha,\beta$ -unsaturated carbonyl compound **14** (480 mg, 1.00 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (20 mL) was added BF<sub>3</sub>·OEt<sub>2</sub> (1.85 mL, 15.00 mmol) at 0 °C. The reaction mixture was stirred for 1 h and then quenched with saturated aqueous NaHCO<sub>3</sub> solution (6 mL). The organic layer was separated and the aqueous part was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 × 15 mL). The combined organic layers were dried over anhydrous MgSO<sub>4</sub>,

concentrated *in vacuo*, and the residue was purified by silica gel column chromatography (50% ethyl acetate-hexane) to afford **15** (290 mg, 64%) as a viscous liquid. IR (liquid,  $\nu_{\max}$   $\text{cm}^{-1}$ ): 1715.  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  (ppm) 7.44-7.37 (5H, m), 6.93 (1H, d,  $J$  2.0), 6.52 (1H, d,  $J$  2.0), 4.96 (1H, d,  $J$  11.3), 4.89 (1H, d,  $J$  11.3), 4.80-4.72 (1H, m), 3.96 (3H, s), 3.95 (3H, s), 3.80 (3H, s), 3.27 (1H, dd,  $J$  15.90, 2.58), 3.02 (1H, dd,  $J$  17.01, 6.38), 2.73-2.63 (2H, m), 2.21 (3H, s).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ ):  $\delta$  (ppm) 204.9, 162.2, 161.1, 159.8, 168.6, 145.3, 136.9, 135.4, 128.7 ( $\times 2$ ), 128.5, 128.2 ( $\times 2$ ), 127.2, 116.7, 112.6, 99.7, 93.3, 75.8, 73.1, 63.4, 56.4, 55.3, 48.0, 30.9, 28.9. HRMS (ESI)  $m/z$  calcd for  $\text{C}_{26}\text{H}_{27}\text{O}_7$  ( $\text{M}+\text{H}$ ) $^+$  451.1761, found 451.1761.

**1-[10-(Benzyloxy)-1-hydroxy-6,8,9-trimethoxyanthracen-2-yl]ethanone (16)**. The  $\alpha,\beta$ -unsaturated carbonyl compound **15** (200 mg, 0.46 mmol) was treated with 10% aqueous ethanolic KOH solution (3 mL) for 1 h under refluxing conditions. After complete disappearance of the starting material as monitored by TLC, the reaction mixture was cooled down to rt and the solution was extracted with ethyl acetate ( $3 \times 10$  mL). The organic layers were then combined, dried over anhydrous  $\text{MgSO}_4$  and concentrated under reduced pressure. The crude material was purified by silica gel column chromatography (5% ethyl acetate-hexane) to obtain a yellow solid following evaporation of the eluent, **16** (40 mg, 20%). mp 140-142  $^\circ\text{C}$ . IR (liquid,  $\nu_{\max}$   $\text{cm}^{-1}$ ): 3403, 1715, 1668.  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  (ppm) 15.31 (1H, br s), 7.61-7.58 (4H, m), 7.50-7.36 (3H, m), 7.04 (1H, d,  $J$  2.2), 6.49 (1H, d,  $J$  2.2), 5.08 (2H, s), 4.02 (3H, s), 4.00 (3H, s), 3.84 (3H, s), 2.70 (3H, s).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ ):  $\delta$  (ppm) 202.8, 167.2, 159.8, 159.3, 155.8, 144.8, 137.6, 131.9, 129.1, 128.8 ( $\times 2$ ), 128.3, 127.9 ( $\times 2$ ), 125.0, 116.2, 115.9, 111.96, 111.8, 99.4, 91.6, 76.3, 63.9, 56.4, 55.3, 27.2. HRMS (ESI)  $m/z$  calcd for  $\text{C}_{26}\text{H}_{25}\text{O}_6$  ( $\text{M}+\text{H}$ ) $^+$  433.1648, found 433.1648.

**5-(Benzyloxy)-7,9,10-trimethoxy-3-(2-methyl-1,3-dithiolan-2-yl)methyl-3,4-dihydro-1H-benzo[g]-isochromen-1-one (17) and 5-(benzyloxy)-10-hydroxy-7,9-dimethoxy-3-(2-methyl-1,3-dithiolan-2-yl)methyl-3,4-dihydro-1H-benzo[g]isochromen-1-one (17a)**. To a solution of the lactone **15** (280 mg, 0.62 mmol) in toluene (20 mL) was added ethanedithiol (0.10 mL, 1.24 mmol) followed by the addition of *p*-toluenesulfonic acid monohydrate (5 mg). The solution was heated to reflux for 18 h, cooled down to rt, and extracted with ethyl acetate ( $3 \times 10$  mL). The organic layers were combined, dried over anhydrous  $\text{MgSO}_4$ , concentrated *in vacuo*, and the crude product was purified by column chromatography to afford **17** as a solid white product (220 mg, 70%) along with a minor viscous liquid **17a** (50 mg, 16%) following evaporation of the eluent. **17**: mp 168-170  $^\circ\text{C}$ . IR (solid,  $\nu_{\max}$   $\text{cm}^{-1}$ ): 1711.  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  (ppm) 7.47-7.37 (5H, m), 6.96 (1H, d,  $J$  2.3), 6.52 (1H, d,  $J$  2.3), 4.97 (1H, d,  $J$  11.3), 4.91 (1H, d,  $J$  11.3), 4.63-4.57 (1H, m), 3.97 (3H, s), 3.96 (3H, s), 3.82 (3H, s), 3.37-3.26 (5H, m), 2.78 (1H, dd,  $J$  15.9, 10.4), 2.52 (1H, dd,  $J$  15.0, 6.9), 2.24 (1H, dd,  $J$  15.0, 3.7), 1.90 (3H, s).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ ):  $\delta$  (ppm) 162.3, 161.0, 159.7, 158.4, 145.3, 136.9, 135.3, 128.8 ( $\times 2$ ), 128.5, 128.2 ( $\times 2$ ), 127.4, 116.7, 112.95, 99.7, 93.2, 76.3, 75.9, 64.7, 63.4, 56.4, 55.3, 49.8, 39.9, 39.8, 32.9, 30.1. HRMS (ESI)  $m/z$  calcd for  $\text{C}_{28}\text{H}_{31}\text{O}_6\text{S}_2$  ( $\text{M}+\text{H}$ ) $^+$  527.1564, found 527.1564. **17a**: IR (liquid,  $\nu_{\max}$   $\text{cm}^{-1}$ ): 1719, 1649, 1607, 1558.  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  (ppm) 12.99 (1H, s), 7.45-7.37 (5H, m), 6.93 (1H, d,  $J$  2.3), 6.50 (1H, d,  $J$  2.3), 4.90 (2H, s), 4.78-4.71 (1H, m), 3.99 (3H, s), 3.83 (3H, s), 3.41-3.27 (5H, m), 2.73 (1H, dd,  $J$  16.3, 10.6), 2.53 (1H, dd,  $J$  15.1, 7.1), 2.26 (1H, dd,  $J$  15.1, 3.6), 1.89 (3H, s).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ ):  $\delta$  (ppm) 170.6, 162.2, 161.2, 161.0, 141.0, 137.0, 136.6, 128.7 ( $\times 2$ ), 128.5, 128.3 ( $\times 2$ ), 124.7, 111.4, 99.6, 98.7, 93.7, 77.8, 75.6, 64.6, 56.3, 55.4, 49.9, 39.92 ( $\times 2$ ), 32.94, 28.9. HRMS (ESI)  $m/z$  calcd for  $\text{C}_{27}\text{H}_{29}\text{O}_6\text{S}_2$  ( $\text{M}+\text{H}$ ) $^+$  513.1407, found 513.1407. Crystal data for **17**, assigned CCDC no. 1030317,  $\text{C}_{28}\text{H}_{30}\text{O}_6\text{S}_2$ :  $M_r$  526.64  $\text{g}\cdot\text{mol}^{-1}$ ; crystal dimensions (mm)  $0.31 \times 0.16 \times 0.07$ ; crystal system, monoclinic; space group,  $P2_1/c$ ; unit cell dimensions and volume,  $a = 9.3246(2)$   $\text{\AA}$ ,  $b = 21.4592(5)$   $\text{\AA}$ ,  $c = 12.9652(3)$   $\text{\AA}$ ,  $\alpha = 90^\circ$ ,  $\beta = 101.039(1)^\circ$ ,  $\gamma = 90^\circ$ ,  $V = 2546.3(1)$   $\text{\AA}^3$ ,  $Z = 4$ ; calculated density  $\rho_{\text{calcd}}$ , 1.374  $\text{Mg}/\text{m}^3$ ; linear absorption coefficient,  $\mu = 0.129$   $\text{mm}^{-1}$ ; radiation and wavelength,  $\text{MoK}\alpha_1 = 0.71073$   $\text{\AA}$ ; temperature of measurement, 173(2) K,  $2\theta_{\max}$   $28.00^\circ$ ; 31024 reflections measured, 6106 unique reflections,



4962 observed reflections [ $I > 2\sigma(I)$ ];  $R_{\text{int}} = 0.0256$ ;  $R_1 [I > 2.0\sigma(I)] = 0.0463$ ,  $wR_2 [\text{all}] = 0.1287$ ,  $\text{GoF} = 1.039$ , refined on  $F^2$ ; residual electron density, 1.055 and  $-0.484 \text{ e}\text{\AA}^{-3}$ .

**5-Hydroxy-7,9-dimethoxy-3-propyl-3,4-dihydro-1H-benzo[g]isochromen-1-one (18).** To a solution of **17** (200 mg, 0.38 mmol) in EtOH (50 mL), was added Raney nickel (5.0 g, 50% slurry in  $\text{H}_2\text{O}$ ). The heterogeneous mixture was heated to reflux for 72 h and then gravity filtered through filter paper to remove the solid. The solvent was then removed under reduced pressure to give **18** (95 mg, 79%) as a white solid. mp 161–163 °C. IR (solid,  $\nu_{\text{max}} \text{ cm}^{-1}$ ): 3216, 1691.  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  (ppm) 8.65 (1H, s), 6.93 (1H, d,  $J$  1.9), 6.47 (1H, d,  $J$  1.9), 4.55–4.46 (1H, m), 3.94 (3H, s), 3.94 (3H, s), 3.18 (1H, dd,  $J$  16.1, 3.1), 2.82 (1H, dd,  $J$  16.1, 11.2), 1.89–1.82 (1H, m), 1.76–1.62 (2H, m), 1.61–1.46 (2H, m), 0.95 (3H, t,  $J$  7.3).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ ):  $\delta$  (ppm) 166.8, 160.9, 158.2, 145.8, 129.2, 121.1, 119.8, 119.5, 118.5, 98.2, 91.5, 78.1, 55.7, 55.6, 37.3, 27.5, 18.3, 13.9. HRMS (ESI)  $m/z$  calcd for  $\text{C}_{18}\text{H}_{21}\text{O}_5 (\text{M}+\text{H})^+$  317.1390, found 317.1390.

**7,9-dimethoxy-3-propyl-3,4-dihydro-1H-benzo[g]isochromen-1,5,10-trione (10).** Naphthalene **18** (40 mg, 0.12 mmol) was dissolved in  $\text{CH}_3\text{CN}$  (2 mL) and ceric ammonium nitrate (CAN) (131 mg, 0.24 mmol) in  $\text{H}_2\text{O}$  (1 mL) was added dropwise at 0 °C. The reaction mixture was stirred for 30 min, followed by the addition of  $\text{H}_2\text{O}$  (2 mL) and ethyl acetate (10 mL). The organic layer was separated, dried over anhydrous  $\text{MgSO}_4$ , filtered and the filtrate was then concentrated *in vacuo*. The residue was purified by silica gel column chromatography (10% ethyl acetate-hexane) to afford the quinone **10** (30.5 mg, 80%) as an orange solid. mp 107–108 °C. IR (solid,  $\nu_{\text{max}} \text{ cm}^{-1}$ ): 1734  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  (ppm) 7.08 (1H, d,  $J$  2.1), 6.69 (1H, d,  $J$  1.1), 4.49–4.39 (1H, m), 3.89 (3H, s), 3.88 (3H, s), 3.04 (1H, dd,  $J$  18.7, 2.9), 2.44 (1H, dd,  $J$  18.7, 11.9), 1.86–1.77 (1H, m), 1.72–1.63 (1H, m), 1.59–1.43 (2H, m), 0.95 (3H, t,  $J$  7.3).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ ):  $\delta$  (ppm) 184.1, 177.9, 164.6, 162.0, 160.3, 147.7, 134.9, 130.7, 114.8, 105.1, 103.3, 77.3, 56.5, 56.0, 36.6, 26.9, 18.0, 13.8. HRMS (ESI)  $m/z$  calcd for  $\text{C}_{18}\text{H}_{19}\text{O}_6 (\text{M}+\text{H})^+$  331.1187, found 331.1187.

## Acknowledgements

This research was supported by the National Research Foundation (NRF, GUN 2053652 and IRDP of the NRF (South Africa) and by the University of the Witwatersrand (Science Faculty Research Council). We also gratefully acknowledge the University of the Witwatersrand for a postdoctoral Fellowship to MFH. Mr B Moolman and Dr M Stander from the MS service at the University of Stellenbosch are thanked for recording the HRMS.

## Supplementary Material

Copies of  $^1\text{H}$  NMR and  $^{13}\text{C}$  NMR spectra presented in the Supplementary Material are available in the online version.

## References

<sup>II</sup>For queries regarding X-ray crystallography, contact [Andreas.Lemmerer@wits.ac.za](mailto:Andreas.Lemmerer@wits.ac.za)

<https://doi.org/10.1039/C4NP00127C>.

2. (a) Parisot, D.; Devys, M.; Férézou, J.-P.; Barbier, M. *Phytochem.* **1983**, 22, 1301–1303.  
[https://doi.org/10.1016/0031-9422\(83\)80253-2](https://doi.org/10.1016/0031-9422(83)80253-2).  
 (b) Tatum, J. H.; Baker, R. A.; Berry, R. E. *Phytochem.* **1989**, 28, 283–284.  
[https://doi.org/10.1016/0031-9422\(89\)85062-9](https://doi.org/10.1016/0031-9422(89)85062-9).
3. Kitagawa, R. R.; Raddi, M. S. G.; dos Santos, L. C.; Vilegas, W. *Chem. Pharm. Bull.* **2004**, 52, 1487–1488.
4. Devienne, K. F.; Raddi, M. S. G.; Varanda, E. A.; Vilegas, W. Z. *Naturforsch. C* **2002**, 57, 85–88.
5. Varanda, E. A.; Raddi, M. S. G.; de Luz Dias, F.; Araújo, M. C. P.; Gibran, S. C. A.; Takahashi, C. S.; Vilegas, W. *Teratogenesis, Carcinogenesis, and Mutagenesis* **1997**, 17, 85–95.  
[https://doi.org/10.1002/\(SICI\)1520-6866\(1997\)17:2<85::AID-TCM4>3.0.CO;2-A](https://doi.org/10.1002/(SICI)1520-6866(1997)17:2<85::AID-TCM4>3.0.CO;2-A).
6. *Wood and Tree Fungi: Biology, Damage, Protection, and Use*, Schmidt, O. Springer-Verlag Berlin Heidelberg 2006, Chapter 4, p. 120, ISBN-10 3-540-32138-1.
7. Blackburn, G. M.; Ekong, D. E. U.; Neilson, A. H.; Todd, L. *Chimia* **1965**, 19, 208–212.
8. Edwards, R. L.; Kale, N. *Tetrahedron* **1965**, 21, 2095–2107.  
[https://doi.org/10.1016/S0040-4020\(01\)98346-2](https://doi.org/10.1016/S0040-4020(01)98346-2).
9. Saikawa, Y.; Watanabe, T.; Hashimoto, K.; Nakata, M. *Phytochem.* **2000**, 55, 237–240.  
[https://doi.org/10.1016/S0031-9422\(00\)00282-X](https://doi.org/10.1016/S0031-9422(00)00282-X).
10. Giesbers, G., Van Schenck, J.; Gutierrez, S. M. V.; Robinson, S. C.; Ostroverkhova, O. *ACS Omega*, **2019**, 4, 13309–13318.d  
<https://doi.org/10.1021/acsomega.9b01490>.
11. Giles, R. G. F.; Reuben, M. K.; Roos, G. P. S. *Afr. J. Chem.* **1979**, 32, 127–129.
12. Giles, R. G. F.; Green, I. R.; Hugo, V. I., S. *Afr. J. Chem.* **1990**, 43, 28–33.
13. Donner, C. D.; Gill, M.; Tewierik, L. M. *Molecules* **2004**, 9, 498–512.
14. Donner, C.; Cuzzupe, A.; Falzon, C.; Gill, M. *Tetrahedron*, **2012**, 68, 2799–2805.  
<https://doi.org/10.1016/j.tet.2012.02.009>.
15. Sumani, J. E. Y.; Ngwira, K. J.; Lemmerer A.; de Koning, C. B. *Eur. J. Org. Chem.* **2019**, 1145–1153.  
<https://doi.org/10.1002/ejoc.201801595>.
16. (a) Mmutlane, E. M. Michael, J. P.; Green, I. R.; de Koning, C. B. *Org. & Biomol. Chem.* **2004**, 2, 2461–2470.  
<https://doi.org/10.1039/B407208A>.  
 (b) Pillay, A.; Rousseau, A. L.; Fernandes, M. A.; de Koning, C. B. *Org. & Biomol. Chem.* **2012**, 10, 7809–7819.  
<https://doi.org/10.1039/C2OB26126J>.
17. Lowell, A. N.; Fennie, M. W.; Kozlowski, M. C. *J. Org. Chem.*, **2008**, 73, 1911–1918.  
<https://doi.org/10.1021/jo7024114>.
18. Anthracene **16** is presumably formed by abstraction of a benzylic hydrogen of the allyl side chain of **15** which results in nucleophilic attack at the ester resulting in the formation of the resultant naphthol **16**. We were unable to improve the yield of this aromatic ring forming reaction by attempting the transformation under a variety of reaction conditions and different bases.
19. X-ray crystallographic data for compound **17** is given in the experimental section.
20. Wadsworth, A. D.; Jonathan Sperry, J.; Brimble, M. A. *Synthesis*, **2010**, 2604–2608.  
<https://doi.org/10.1055/s-0029-1218832>.
21. Bruker, SAINT+, version 6.02 (including XPREP), Bruker AXS Inc., Madison, WI, USA, 1999.
22. Farrugia, L. J. *App. Cryst.* **1999**, 32, 837–838.
23. Sheldrick, G. M. SHELX, release 97-2 (includes SHELXS and SHELXL), University of Göttingen, 1997.
24. Farrugia, L. J. *App. Cryst.* **1997**, 30, 565.

25. Spek, A. J. *App. Cryst.* **2003**, 36, 7-13.

This paper is an open access article distributed under the terms of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>)