

Copper-catalyzed tandem reaction of cyclic esterification/selenoxidation for 11oxo-11H-5-oxa-11-selena-benzo[*a*] fluoren-6-ones synthesis

Xinyan Zhou^a, Yueqin Xue^b, Yiying Cheng,^a and Runsheng Xu^{a*}

^a Department of Biology and Environment, Jiyang College of Zhejiang A &F University, Shaoxing 311800, Zhejiang, China ^b LinYi Central Hospital, Linyi 276400, ShanDong, China Email: <u>20140041@zafu.edu.cn</u>

Received	mm-dd-yyyy	Accepted	mm-dd-yyyy		Published on line	mm-dd-yyyy	
Dates to be inserted by editorial office							

Abstract

Organoselenium compounds have attracted much attention for their wide use in a variety of fields. A transition-metal-catalyzed cross-coupling reaction is the mostly commonly used methodology for incorporation of a Se atom into aromatic structures. A copper-catalyzed tandem reaction of cyclic esterification/selenoxidation has been developed. Starting from sample 3-(2-hydroxy-phenyl)-acrylic acids with 2-halide-benzeneselenols, 11-oxo-11H-5-oxa-11-selena-benzo[*a*]fluoren-6-ones, with reported biological-activity importance, were efficiently synthesized in good-to-high yields. This new methodology provides an approach toward C(sp²)-Se bond formation.



Keywords: Copper-catalyzed, tandem reaction, cyclic esterification, selenoxidation, C(sp²)-Se bond formation, organoselenium, organometallic

Introduction

Due to their important applications in the synthesis of materials,¹⁻³ pharmaceutical agents,^{4,5} fluorescent probes,⁶ and functional organic materials,⁷ organoselenium compounds syntheses have attracted extensive attention of synthetic chemists. It is known that a transition-metal-catalyzed cross-coupling reaction is the methodology used mostly for the incorporation of a Se atom into aromatic frameworks,^{8,9} however, pre-functionalization of the substrate is generally required. Simple methods of C-Se bonds formation have been scarcely described.¹⁰⁻¹²

In recent decades, the activation of C-H bonds is considered to be one of the most useful C-Se bondformation strategies (Scheme1, **a**). In comparison with the C-X (X = I, Br, Cl) cross-coupling reactions, however, the C(sp²)-Se bond selenoxidation direct-cross-coupling reactions need harsher conditions, and activated reaction systems (Scheme1, **b**). ^{13,14} Given the present challenges, the development of more efficient, and environmentally friendly, chemical processes for drug discovery is required.^{15,16}



Scheme 1. C(sp²)-Se bond formation synthesis approaches

11-Oxo-11H-5-oxa-11-selena-benzo[*a*]fluoren-6-one constitutes the central core unit of a variety of natural polycyclic lactones with important biological activities including anticancer, antibacterial, antimyotoxic, and phytoalexine effects.^{17,18} A wide range of biological properties make 11-oxo-11H-5-oxa-11-selena-benzo[*a*]fluoren-6-ones an interesting synthetic target for chemists. A number of synthetic methods have been developed for the construction of this privileged structural unit. Most of the reported procedures involve multiple steps with moderate overall yields. The starting materials are often not very readily available, and harsh reaction conditions are usually required. In view of these limitations, the development of an efficient strategy for the 11-oxo-11H-5-oxa-11-selena-benzo[*a*]fluoren-6-ones synthesis is highly desirable.

Herein, we report on a novel copper-catalyzed tandem reaction of cyclic esterification/selenoxidation (Scheme 2). Versatile 11-oxo-11H-5-oxa-11-selena-benzo[a]fluoren-6-one compounds were efficiently synthesized in good-to-high yields under mild conditions. This new methodology provides an approach towards C(sp²)-Se bond formation.



Scheme 2. Copper-catalyzed tandem reaction for cyclic esterification/selenoxidation

Results and Discussion

First, reaction conditions were screened based on the model reaction of 3-(2-hydroxy-phenyl)-acrylic acid **1a** with 2-iodine-benzeneselenol **2a** (Table 1). The structure of **3a** was confirmed by NMR and MS analysis (Please see Supplementary Material). The copper catalysts displayed good catalytic activity in the presence of Cs_2CO_3 (entries 1-6). In addition, CuI exhibited superior catalytic efficiency over all of the other examined copper catalysts (entry 7). These results indicated that Cs_2CO_3 was the optimal base (entry 12), which produced the product **3a** in 84% overall yield. It was also noted that the product yield was decreased when the reaction temperature was less or greater than 100 °C (entries 13 and 14). Furthermore, the results also show that the reaction yield with DMSO as solvent is higher than that of other solvents (entries 15 and 16). Thus, the optimum reaction condition was determined to be a **1a:2a** ratio of 1:1.5 in the presence of CuI (10 mol%) and Cs_2CO_3 (2 equiv) in DMSO (5 mL) at 100 °C for 12 hours (Table 1, entry 12).





12	Cul	Cs ₂ CO ₃	1:1.5	84
13	Cul	Cs ₂ CO ₃	1:1.5	71 ^c
14	Cul	Cs ₂ CO ₃	1:1.5	78 ^d
15	Cul	Cs ₂ CO ₃	1:1.5	76 ^e
16	Cul	Cs ₂ CO ₃	1:1.5	57 ^f

^a Unless otherwise noted, reaction conditions were **1a** (0.5 mmol), **2a** (0.5 mmol), copper catalyst (10 mol%), base (2 equiv), DMSO (5 mL), 100 °C for 12 h. b Isolated yield. c At 90 °C. ^d At 110 °C. ^e In CHCl₃. *f* In DMF.

Next, a wide array of 3-(2-hydroxy-phenyl)-acrylic acids **1** and 2-iodine-benzeneselenols **2** were subjected to this reaction conditions, providing the products **3** in good-to-excellent yields (71-90%, Table 2). 3-(2-Hydroxy-phenyl)-acrylic acids **1** bearing an electron-donating group (Me and MeO) demonstrated better activity than those bearing an electron-withdrawing group (F, Cl, and Br). It was notable that very strong electron-withdrawing groups, e.g., the trifluoromethyl and nitro groups, failed to obtain the corresponding products.





4	4-CH₃O	Н	MeO 3d	90
5	4-CH₃O	4,5-diCH₃O	MeO 3e	78
6	4-CH₃O	Acene	MeO O O O O O O O O O O O O O O O O O O	83
7	5-F	Н	F 3g	73
8	5-Cl	Н	Cl Cl O O O O Se O O O O O O O O O O O O O O	77
9	5-Br	Н	Br Br 3i	71
10	4,5-diCH₃	Н	Me Me 3j	76



^{*a*} Unless otherwise noted, reaction conditions were **1** (0.5 mmol), **2** (0.75 mmol), Cul (10 mol%), Cs₂CO₃ (2 equiv), DMSO (5 mL), 100 °C for 12 h. ^{*b*} Isolated yield.

Other 3-(2-hydroxy-phenyl)-acrylic acids **1** and 2-bromo-benzeneselenols **4** also successfully provided the corresponding products in 54-77% yields, respectively (Table 3). The 3-(2-hydroxy-4,5-dimethoxy-phenyl)-acrylic acid displayed a moderate reactivity with 2-bromo-benzeneselenol with 66% yield (entry 4). Furthermore, to our delight, reactants with more substituents also performed smoothly which displayed a moderate reactivity 6).

Table 3 Copper-catalyzed tandem reaction of cyclic esterification/selenoxidation^{*a*}





^{*a*} Unless otherwise noted, reaction conditions were **1** (0.5 mmol), **3** (0.75 mmol), Cul (10 mol%), NaOEt (2 equiv), DMSO (5 mL), 100 °C for 12 h. ^{*b*} Isolated yield.

We propose a possible reaction mechanism in Scheme 3. At the beginning of the reaction, the Cul generated a Cu^I intermediate **6**. Next, intermediate **7** was provided from **2** via an oxidation addition step. Intermediate **8** was then provided from **1** via intramolecular esterification. Finally, through the oxidation reaction by DMSO and CS_2CO_3 , intermediate **9** generated the desired products **3**, and, concomitantly, formed intermediate **6**, which then re-entered the catalytic cycle.



Scheme 3. Proposed mechanism

Conclusions

In summary, we report a copper-catalyzed tandem reaction of cyclic esterification/selenoxidation. Starting from sample 3-(2-hydroxy-phenyl)-acrylic acids with 2-halide-benzeneselenols, a number of 11-oxo-11H-5-oxa-11-selena-benzo[*a*]fluoren-6-ones were efficiently synthesized in good-to-high yields. This new methodology provides an economical approach toward C(sp²)-Se bond formation for biologically-active 11-oxo-11H-5-oxa-11-selena-benzo[a]fluoren-6-one compounds.

Experimental Section

General. All reagents used in the experiment were obtained from commercial sources and used without further purification. Solvents for chromatography were of technical grade and distilled prior to use. Solvent mixtures are understood to be volume/volume. Chemical yields refer to pure isolated substances. Catalysts were purchased from Alfa Aesar (analytical reagent). Thin layer chromatography (TLC) employed glass 0.25 mm silica gel plates with an F-254 indicator, visualized by irradiation with UV light. The NMR spectra were recorded on a Bruker AVANCE III-400 spectrometer at 400 MHz and 100 MHz for ¹H and ¹³C NMR, respectively, in CDCl₃. The NMR chemical shifts were reported in ppm relative to 7.26 and 77 ppm of CDCl₃ as internal standards for ¹H and ¹³C NMR, respectively. The mass spectra were performed on a Bruker Esquire 3000 plus mass spectrometer equipped with an ESI interface and ion trap analyzer. The ESI-HRMS was tested on a Bruker 7-tesla FT-ICR MS equipped with an electrospray source.

General synthesis methods of 3a-3l

A solution of 3-(2-hydroxy-phenyl)-acrylic acids **1** (0.5 mmol), 2-halide-benzeneselenols **2** or **4** (0.75 mmol), Cul (10 mol%) and Cs_2CO_3 (2 equiv) in DMSO (5 mL) was stirred under air. After stirring at 100 °C for 12 h, it was cooled to room temperature. The reaction mixture was then quenched with a saturated salt-water solution (10 mL). After that, the solution was extracted with ethyl acetate (3 × 10 mL), and then washed with saturated Na₂CO₃ solution. The organic layers were combined and dried by Na₂SO₄ and concentrated in vacuo. The pure product **3** or **5** was afforded by flash column chromatography on silica gel (cyclohexane/ethyl acetate).

11-Oxo-11*H***-5-oxa-11-selena-benzo**[*a*]**fluoren-6-one (3a)** white solid, mp 171-172 °C, was afforded by flash column chromatography on silica gel (cyclohexane/ethyl acetate = 3:1); ¹H NMR (500 MHz, CDCl₃): δ 8.15 (dd, J_1 2.4 Hz, J_2 6.6 Hz, 1H), 8.05 (dd, J_1 1.5 Hz, J_2 7.8 Hz, 1H), 7.68 (dd, J_1 1.7 Hz, J_2 6.9 Hz, 1H), 7.60-7.64 (m, 1H), 7.41-7.52 (m, 4H); ¹³C NMR (125 MHz, CDCl₃): δ 160.0, 158.1, 155.6, 153.7, 131.9, 126.8, 125.2, 124.7, 123.5, 121.9, 117.5, 112.7, 111.8, 105.9 (one peak is missing due to overlap); HRMS (+ESI) Calcd for C₁₅H₉O₃Se [M+H]⁺: 316.9717, found 316.9719.

2-Methyl-11-oxo-11*H***-5-oxa-11-selena-benzo**[*a*]**fluoren-6-one (3b)** yellow solid, mp 153-154 °C, was afforded by flash column chromatography on silica gel (cyclohexane/ethyl acetate = 5:1); ¹H NMR (500 MHz, CDCl₃): δ 8.07-8.18 (m, 1H), 7.80 (br s, 1H), 7.59-7.70 (m, 1H), 7.32-7.51 (m, 4H), 2.48 (s, 3H); ¹³C NMR (125 MHz, CDCl₃): δ 160.0, 158.2, 155.5, 151.9, 134.5, 133.0, 126.6, 125.1, 123.5, 121.8, 121.5, 117.2, 112.3, 111.7, 105.8, 20.9; HRMS (+ESI) Calcd for C₁₆H₁₁O₃Se [M+H]⁺: 330.9873, found 330.9875.

2-Methyl-13-oxo-13*H***-5-oxa-13-selena-dibenzo**[*a*,*h*]**fluoren-6-one (3c)** yellow solid, mp 141-143 °C, was afforded by flash column chromatography on silica gel (cyclohexane/ethyl acetate = 5:1); ¹H NMR (500 MHz, CDCl₃): δ 8.59 s, 1H), 7.94-8.06 (m, 3H), 7.86 (s, 1H), 7.42-7.56 (m, 4H), 2.51 (s, H); ¹³C NMR (125 MHz, CDCl₃):

δ 162.4, 158.3, 153.9, 152.4, 35.7, 134.7, 133.7, 132.1, 128.5, 127.9, 126.2, 125.3, 123.4, 121.8, 20.4, 117.3, 112.1, 107.7, 105.2, 20.9 ; HRMS (+ESI) Calcd for C₂₀H₁₃O₃Se [M+H]⁺: 381.0030, Found 381.0032.

3-Methoxy-11-oxo-11*H***-5-oxa-11***I***4-selena-benzo**[*a*]**fluoren-6-one (3d)** yellow solid, mp 178-179 °C, was afforded by flash column chromatography on silica gel (cyclohexane/ethyl acetate = 5:1); ¹H NMR (500 MHz, CDCl₃): δ 8.04–8.14 (m, 1H), 7.91 (dd, *J*₁ 1.9 Hz, *J*₂ 7.3 Hz, 1H), 7.57-7.67 (m, H), 7.39-7.47 (m, 2H), 6.94-7.02 (m, 2H), 3.91 (s, 3H); ¹³C NMR 125 MHz, CDCl₃): δ 163.0, 160.6, 158.3, 155.5, 155.2, 126.0, 125.0, 23.5, 122.8, 121.4, 113.0, 111.4, 105.8, 103.3, 101.3, 55.7; HRMS (+ESI) calcd for C₁₆H₁₁O₄Se [M+H]⁺: 346.9823, found 346.9825.

3,8,9-Trimethoxy-11-oxo-11H-5-oxa-11-selena-benzo[*a*]**fluoren-6-one (3e)** yellow solid, mp 130-133 °C , was afforded by flash column chromatography on silica gel (cyclohexane/ethyl acetate = 5:1); ¹H NMR (500 MHz, CDCl₃): δ 7.85 (d, *J* 8.4 Hz, 1H), 7.51 (s, 1H), 7.13 (s, 1H), 6.94–7.04 (m 2H), 4.00 (s, 3H), 3.96 (s, 3H), 3.90 (s, 3H); ¹³C NMR (125 MHz, CDCl₃): δ 162.4, 159.7, 158.8, 154.9, 149.9, 149.0, 147.9, 122.3, 115.5, 113.0, 106.3, 103.8, 102.2, 101.4, 95.5, 56.5, 56.4, 55.8; HRMS (+ESI) Calcd for C₁₈H₁₅O₆Se [M+H]⁺: 407.0034, found 407.0036.

3-Methoxy-13-oxo-13*H***-5-oxa-13-selena-dibenzo**[*a*,*h*]**fluoren-6-one (3f)** yellow solid, mp: 157-159 °C m, was afforded by flash column chromatography on silica gel (cyclohexane/ethyl acetate = 5:1); ¹H NMR (500 MHz, CDCl₃): δ 8.55 (s, 1H), 7.87-8.12 (m, 4H), 7.43-7.60 (m, 2H), 6.92-7.10 (m, 2H), 3.93 (s, 3H); ¹³C NMR (125 MHz, CDCl₃): δ 163.5, 162.9, 158.2, 156.1, 153.8, 131.8, 131.4, 128.3, 127.9, 125.9, 125.2, 123.4, 123.2, 119.8, 113.2, 107.5, 105.6, 102.6, 101.4, 55.8 ; HRMS (+ESI) Calcd for C₂₀H₁₃O₄Se [M+H]⁺: 396.9979, found 396.9979.

2-Fluoro-11-oxo-11*H***-5-oxa-11-selena-benzo**[*a*]**fluoren-6-one (3g)** yellow solid, mp 178-180 °C, was afforded by flash column chromatography on silica gel (cyclohexane/ethyl acetate = 5:1); ¹H NMR (500 MHz, CDCl₃): δ 8.15 (d, *J* 7.4 Hz, 1H), 7.62-7.74 (m, 2H), 7.45-7.55 (m, 3H), 7.29-7.34 (m, 1H); ¹³C NMR (125 MHz, CDCl₃): δ 159.0 (d, *J* 2.8 Hz), 158.9 (d, *J* 245.7 Hz), 157.6, 155.7, 149.8 (d, *J* 1.6 Hz), 127.2, 126.4, 123.2, 122.0, 119.4 (d, *J* 34.7 Hz), 119.3, 113.4 (d, *J* 9.8 Hz), 111.8, 107.4 (d, *J* 25.8 Hz), 106.6; HRMS (+ESI) Calcd for C₁₅H₈FO₃Se [M+H]⁺: 334.9623, found 334.9625.

2-Chloro-11-oxo-11H-5-oxa-11l4-selena-benzo[a]fluoren-6-one (3h) yellow solid, mp 189-191 °C, was afforded by flash column chromatography on silica gel (cyclohexane/ethyl acetate = 5:1); ¹H NMR (500 MHz, CDCl₃): δ 8.14 (dd, J_1 1.2 Hz, J_2 7.9 Hz, 1H), 8.01 (d, J 2.5 Hz, 1H), 7.68 (d, J 7.6 Hz, 1H), 7.55 (dd, J_1 2.5 Hz, J_2 8.9 Hz, 1H), 7.47-7.52 (m, 2H), 7.45 (d, J 8.9 Hz, 1H); ¹³C NMR (125 MHz, CDCl₃): δ 158.5, 157.4, 155.6, 151.8, 131.8, 130.2, 127.1, 125.4, 123.1, 121.9, 121.2, 118.8, 113.6, 111.8, 106.5; HRMS (+ESI) Calcd for C₁₅H₈ClO₃Se [M+H]⁺: 350.9327, found 350.9329.

2-Bromo-11-oxo-11*H***-5-oxa-11-selena-benzo**[*a*]**fluoren-6-one (3i)** yellow solid, mp 176-178 °C, was afforded by flash column chromatography on silica gel (cyclohexane/ethyl acetate = 5:1); ¹H NMR (500 MHz, CDCl₃): δ 8.18 (d, *J* 2.3 Hz, 1H), 8.13-8.16 (m, 1H), 7.67-7.71 (m, 2H), 7.46-7.55 (m, 2H), 7.40 (d, *J* 8.8 Hz, 1H); ¹³C NMR (125 MHz, CDCl₃): δ 158.4, 157.3, 155.6, 152.3, 134.6, 127.2, 125.4, 124.3, 123.1, 121.9, 119.1, 117.4, 114.1, 111.8, 106.5; HRMS (+ESI) Calcd for C₁₅H₈BrO3Se [M+H]⁺: 394.8822, found 394.8824.

2,3-Dimethyl-11-oxo-11*H***-5-oxa-11-selena-benzo**[*a*]**fluoren-6-one (3j)** yellow solid, mp 195-197 °C, was afforded by flash column chromatography on silica gel (cyclohexane/ethyl acetate = 5:1); ¹H NMR (500 MHz, CDCl₃): δ 8.03-8.14 (m, 1H), 7.67 (s, 1H), 7.56-7.64 (m, 1H), 7.36-7.50 (m, 2H), 7.21 (s, 1H), 2.34 (s, 6H); ¹³C NMR (125 MHz, CDCl₃): δ 160.2, 158.3, 155.2, 152.1, 142.1, 133.5, 126.2, 124.9, 123.6, 121.6, 121.5, 117.8, 111.5, 109.9, 104.8, 20.4, 19.2; HRMS (+ESI) Calcd for C₁₇H₁₃O₃Se [M+H]⁺: 345.0030, found 345.0032.

2,3-Dimethoxy-13-oxo-13H-5-oxa-13-selena-dibenzo[*a*,*h*]**fluoren-6-one (3k)** yellow solid, was afforded by flash column chromatography on silica gel (cyclohexane/ethyl acetate = 5:1); ¹H NMR (500 MHz, CDCl₃): δ 8.54 (s, 1H), 7.94-8.05 (m, 3H), 7.50-7.55 (m, 2H), 7.39 (s, 1H), 7.00 (s, 1H), 4.04 (s, 3H), 3.99 (s, 3H); ¹³C NMR (125)

MHz, CDCl₃): δ 162.9, 158.5, 153.8, 153.6, 150.4, 146.9, 131.8, 131.5, 128.4, 127.9, 126.0, 125.3, 123.6, 119.9, 114.1, 107.5, 104.4, 102.0, 100.6, 56.6, 56.6; HRMS (+ESI) Calcd for C₂₁H₁₅O₅Se [M+H]⁺: 427.0085, found 427.0087.

2,3-Dimethoxy-11-oxo-11*H***-5-oxa-11***I***4-selena-benzo***[a***]***f***Iuoren-6-one (3I)** yellow solid, mp 207-209 °C, was afforded by flash column chromatography on silica gel (cyclohexane/ethyl acetate = 5:1); ¹H NMR (500 MHz, CDCl₃): δ 8.09 (dd, J_1 3.4 Hz, J_2 5.7 Hz, 1H), 7.62 (dd, J_1 3.0 Hz, J_2 6.2 Hz, 1H), 7.44 (d, J 3.3 Hz, 1H), 7.43 (d, J 3.1 Hz, 1H), 7.35 (s, 1H), 6.99 (s, 1H), 4.02 (s, 3H), 3.97 (s, 3H); ¹³C NMR (125 MHz, CDCl₃): δ 160.5, 158.4, 155.1, 152.9, 149.6, 146.7, 126.1, 125.0, 123.6, 121.5, 111.3, 104.5, 103.6, 101.7, 100.5, 56.3, 56.3; HRMS (+ESI) Calcd for C₁₇H₁₃O₅Se [M+H]⁺: 376.9928, found 376.9930.

Acknowledgements

This research received no external funding by the Natural Science Foundation of China No. 21702186.

Supplementary Material

Supplementary material, consisting of ¹H and ¹³C NMR spectra, can be found in the online version.

References

- 1 Trenner, J.; Depken, C.; Weber, T.; Breder, A. *Angew. Chem., Int. Ed.* **2013**, *52*, 8952-8956. https://doi.org/10.1002/anie.201303662
- 2 Huang, L. W.; Xun, X. D.; Zhao, M.; Xue, J. Z.; Li, G. F.; Hong, L. J. Org. Chem. 2019, 84, 11885-11890. https://doi.org/10.1021/acs.joc.9b01742
- 3 Wei, R. B.; Xiong, H. G.; Ye, C. Q.; Li, Y. J.; Bao, H. L. *Org. Lett.* **2020**, *22*, 3195-3199. https://doi.org/10.1021/acs.orglett.0c00969
- Engman, L.; Stern, D.; Frisell, H.; Vessman, K.; Berglund, M.; Ek, B.; Andersson, C.-M. *Bioorg. Med. Chem.* 1995, *3*, 1255-1262.
 https://doi.org/10.1016/0968-0896(95)00111-S
- 5 Wirth, T. *Angew. Chem., Int. Ed.* **2015**, *54*, 10074-10076. https://doi.org/10.1002/anie.201505056
- 6 Panda, S.; Panda, A.; Zade, S. S. *Coord. Chem. Rev.* **2015**, *300*, 86-100. https://doi.org/10.1016/j.ccr.2015.04.006
- Somasundaram, S.; Chenthamarakshan, C. R.; de Tacconi, N. R.; Ming, Y.; Rajeshwar, K. *Chem. Mater.* 2004, *16*, 3846-3852. https://doi.org/10.1021/cm049293b
- 8 Beletskaya, I. P.; Ananikov, V. P. *Chem. Rev.* **2011**, *111*, 1596-1636. https://doi.org/10.1021/cr100347k
- 9 Wang, Y.; Zhang, W. X.; Wang, Z. T.; Xi, Z. F. *Angew. Chem. Int. Ed.* **2011**, *50*, 8122-8126. https://doi.org/10.1002/anie.201101948
- 10 Qiu, R.; Reddy, V. P.; Iwasaki, T.; Kambe, N. J. Org. Chem. 2015, 80, 367-374.

- 11 Yu, S.; Wan, B.; Li, X. Org. Lett. **2015**, *17*, 58-61. https://doi.org/10.1021/ol503231p
- 12 Xie, W.; Li, B.; Wang, B. J. Org. Chem. **2016**, *81*, 396-403. https://doi.org/10.1021/acs.joc.5b01943
- 13 Duan, F. F.; Song, S. Q.; Xu, R. S. *Chem. Commun.*, **2017**, *53*, 2737-2739. https://doi.org/10.1039/C6CC10303K
- 14 Cai, R. R.; Zhou, Z. D.; Chai, Q. Q.; Zhu, Y. E.; Xu, R. S. *RSC Adv.*, **2018**, 8, 26828-26836. https://doi.10.1039/C8RA05311A
- 15 Noshi, M. N.; El-Awa, A.; Torres, E.; Fuchs, P. L. *J. Am. Chem. Soc.*, **2007**, 129, 11242-11247. https://doi.org/10.1021/ja072890p
- 16 López-Pérez, A.; Robles-Machín, R.; Adrio, J.; Carretero, J. C. Angew. Chem. Int. Edit., **2007**, 46, 9261-9264.

https://doi.org/10.1002/ange.200703258

- 17 Liu, L. L.; Deng, Q. Q.; Weng, S. J.; Yang, X. L.; Zhong, Y. M. *Neuroscience*, **2016**, 332, 53-60. https://doi.org/10.1016/j.neuroscience.2016.06.045
- 18 Zanos, P.; Moaddel, R.; Morris, P. J.; Riggs, L. M.; Highland, J. N.; Georgiou, P.; Pereira, E. F. R.; Albuquerque, E. X.; Thomas, C. J.; Zarate, C. A.; Gould, T. D. *Pharmacol. Rev.*, **2018**, 70, 621-660. <u>https://doi.org/10.1124/pr.117.015198</u>

This paper is an open access article distributed under the terms of the Creative Commons Attribution (CC BY) license (<u>http://creativecommons.org/licenses/by/4.0/</u>)