

Supplementary Material

Synthesis and properties of liquid phenyliodine dicarboxylates



Christina Callovi, Su Wint War, Calvin N. Semczuk, Madison Fisher, Matthew Lucero Rodriguez, Joshua Plaut, Ivana Joshi, I. F. Dempsey Hyatt,* and Daniel L. Silverio*


Department of Chemistry, Adelphi University, One South Avenue, Garden City, NY, 11530



Email: dsilverio@adelphi.edu, ihyatt@adelphi.edu


Table of Contents


¹ H NMR, & ¹³ C NMR, spectra of phenyl- λ^3 -iodanediyl bis(1-adamantoate) (2a)	S2
¹ H NMR, & ¹³ C NMR, spectra of phenyl- λ^3 -iodanediyl bis(2-ethylhexanoate) (2b).....	S4
¹ H NMR, & ¹³ C NMR, spectra of phenyl- λ^3 -iodanediyl bis(2-ethylhexanoate) (2b).....	S5
¹ H NMR, & ¹³ C NMR spectra of phenyl- λ^3 -iodanediyl bis(2-ethylbutanoate) (2c)	S6
Pictures for Miscibility of 2b to 2d	S10
Solubility data for phenyl- λ^3 -iodanediyl bis(1-adamantoate) (2a).....	S13
Partition Coefficient of Compound 2b in Acetonitrile and Hexanes	S13
Partition Coefficient of Compound 2c in Acetonitrile and Hexanes.....	S15
Partition Coefficient of Compound 2d in Acetonitrile and Hexanes	S18
Absorbance Data for Filtered vs. Unfiltered PhI(OAc) ₂ 1a	S21
Absorbance vs. Time Stored for Filtered PhI(OAc) ₂ 1a	S21



^1H NMR, & ^{13}C NMR, spectra of phenyl- λ^3 -iodanediyl bis(1-adamantoate) (2a)



^1H NMR, & ^{13}C NMR, spectra of phenyl- λ^3 -iodanediyl bis(1-adamantoate) (2a)



^1H NMR, & ^{13}C NMR, spectra of phenyl- λ^3 -iodanediyl bis(2-ethylhexanoate) (2b)





^1H NMR, & ^{13}C NMR, spectra of phenyl- λ^3 -iodanediyl bis(2-ethylhexanoate) (2b)


^1H NMR, & ^{13}C NMR spectra of phenyl- λ^3 -iodanediyl bis(2-ethylbutanoate) (**2c**)

^1H NMR, & ^{13}C NMR spectra of phenyl- λ^3 -iodanediyl bis(2-ethylbutanoate) (**2c**)



^1H NMR & ^{13}C NMR of phenyl- λ^3 -iodanediyl bis(2-methylpentanoate) (2d)

^1H NMR & ^{13}C NMR of phenyl- λ^3 -iodanediyl bis(2-methylpentanoate) (2d)


Pictures for Miscibility of **2b** to **2d**compound **2b** with hexanes (2:1,1:1,1:2)compound **2b** with acetonitrile (2:1,1:1,1:2)




compound **2c** with hexanes (2:1,1:1,1:2)





with acetonitrile (2:1,1:1,1:2)





compound **2d** with hexanes (2:1,1:1,1:2)




compound **2d** with acetonitrile (2:1,1:1,1:2)



Solubility data for **phenyl- λ 3-iodanediyl bis(1-adamantoate) (2a)**

Solvent	Solubility (g/ml)
Hexanes	0.005
	0.0051
	0.0044
	0.0053
	0.0042
	0.0046
	0.004
	0.0042
Average	0.0046
Standard Deviation (SD)	0.000481
%RSD	10.5%

Partition Coefficient of Compound **2b** in Acetonitrile and Hexanes**Compound 2b**

Molar mass = 490.42 g/mol


Density = 1.3 g/ml

Acetonitrile layer (0.10 ml of 2b)

	Mass dissolved (g)	Concentration (mol/L)
Trial 1 Vial 1	0.0202	0.0412
Trial 1 Vial 2	0.0184	0.0375
Trial 2 Vial 1	0.0182	0.0371
Trial 3 Vial 1	0.0193	0.0394
Trial 3 Vial 2	0.0198	0.0404
Average	0.0192	0.0391
Standard deviation (SD)	0.00087	0.00177
%RSD	4.52	4.52

Hexanes layer (0.10 ml 2b)

	Mass dissolved (g)	Concentration (mol/L)
Trial 1 Vial 1	0.0095	0.0194
Trial 1 Vial 2	0.0103	0.0210
Trial 2 Vial 1	0.0081	0.0165
Trial 2 Vial 2	0.0109	0.0222
Trial 3 Vial 1	0.0082	0.0167
Trial 3 Vial 2	0.0086	0.0175
Average	0.00927	0.0189
Standard deviation (SD)	0.00116	0.00237
%RSD	12.5	12.5

Partition Coefficient of Compound **2c** in Acetonitrile and Hexanes**Compound 2c**

Molar mass = 434.10 g/mol

Density = 1.3 g/ml

Acetonitrile layer (0.10 ml of 2c)

	Mass dissolved (g)	Concentration (mol/L)
Trial 1 Vial 1	0.0217	0.0500
Trial 1 Vial 2	0.0217	0.0500
Trial 2 Vial 1	0.0202	0.0465
Trial 2 Vial 2	0.0206	0.0475
Average	0.0211	0.0485
Standard deviation (SD)	0.000768	0.00178
%RSD	3.65%	3.67%

Hexanes layer (0.10 ml)

	Mass dissolved (g)	Concentration (mol/L)
Trial 1 Vial 1 (SW-1-67)	0.004	0.00921
Trial 1 Vial 2	0.0037	0.00852
Trial 2 Vial 1 (SW-1-68)	0.0038	0.00875
Trial 2 Vial 2	0.0037	0.00852
Average	0.0038	0.00875
Standard deviation (SD)	0.000141	0.000325
%RSD	3.72%	3.72%

Partition Coefficient = 0.12


Acetonitrile layer (0.20 ml 2c)

	Mass dissolved (g)	Concentration (mol/L)
Trial 1 Vial 1	0.0451	0.104
Trial 1 Vial 2	0.0415	0.0956
Trial 2 Vial 1	0.0423	0.0974
Trial 2 Vial 2	0.0435	0.100
Trial 3 Vial 1	0.0413	0.0951
Trial 3 Vial 2	0.0419	0.0965
Average	0.0426	0.0981
Standard deviation (SD)	0.00145	0.00337
%RSD	3.41%	3.43%

Hexanes layer (0.20 ml of 2c)

	Mass dissolved (g)	Concentration (mol/L)
Trial 1 Vial 1	0.0076	0.0175
Trial 1 Vial 2	0.0076	0.0175
Trial 2 Vial 1	0.0063	0.0145
Trial 2 Vial 2	0.0068	0.0157
Trial 3 Vial 1	0.0073	0.0168
Trial 3 Vial 2	0.0072	0.0166
Average	0.00713	0.0164
Standard deviation (SD)	0.000505	0.00116
%RSD	7.07%	7.05%

Partition Coefficient = 0.11

Partition Coefficient of Compound **2d** in Acetonitrile and Hexanes**Compound 2d (I)**

Molar mass = 434.10 g/mol

Density = 1.3 g/ml

Acetonitrile layer (0.1 ml 2d)

	Mass dissolved (g)	Concentration (mol/L)
Trial 1 Vial 1	0.0296	0.0682
Trial 1 Vial 2	0.0303	0.0698
Trial 2 Vial 1	0.0298	0.0686
Trial 2 Vial 2	0.0285	0.0657
Average	0.0296	0.0681
Standard deviation (SD)	0.000759	0.00172
%RSD	2.57%	2.53%

Hexanes layer (0.1 ml 2d)

	Mass dissolved (g)	Concentration (mol/L)
Trial 1 Vial 1	0.0044	0.0101
Trial 1 Vial 2	0.0054	0.0124
Trial 2 Vial 1	0.0047	0.0108
Trial 2 Vial 2	0.0048	0.0111
Average	0.00483	0.0111
Standard deviation (SD)	0.000419	0.000963
%RSD	8.69%	8.67%

Partition Coefficient = 0.11

Acetonitrile layer (0.2 ml 2d)

	Mass dissolved (g)	Concentration (mol/L)
Trial 1 Vial 1	0.0606	0.14
Trial 1 Vial 2	0.0601	0.138
Trial 2 Vial 1	0.06	0.138
Trial 2 Vial 2	0.0609	0.14
Average	0.0604	0.139
Standard deviation (SD)	0.000424	0.00115
%RSD	0.702%	0.831%

Hexanes layer (0.2 ml 2d)

	Mass dissolved (g)	Concentration (mol/L)
Trial 1 Vial 1	0.011	0.0253
Trial 1 Vial 2	0.0109	0.0251
Trial 2 Vial 1	0.0111	0.0256
Trial 2 Vial 2	0.0111	0.0256
Average	0.011025	0.0254
Standard deviation (SD)	0.0000957	0.000245
%RSD	0.868%	0.964%

Partition Coefficient = 0.12

Absorbance Data for Filtered vs. Unfiltered $\text{PhI}(\text{OAc})_2$ **1a**

Made 20%, 40%, 60%, 80% solution in CH_2Cl_2 of unfiltered PIDA and filtered PIDA. Allowed the solutions to stir for an hour before measurement and the filtration. Absorbances were measured with a Genesys 10S Vis Spectrophotometer equipped with Thermo electron quartz cuvettes with a 10 mm path length immediately following filtration as described in the experimental section of the manuscript.

20% saturation of unfiltered PIDA = 0.884 A

40% saturation of unfiltered PIDA = 1.464 A

60% saturation of unfiltered PIDA = 1.897 A

80% saturation of unfiltered PIDA = 2.216 A

20% saturation of filtered PIDA = 0.000 A

40% saturation of filtered PIDA = 0.000 A

60% saturation of filtered PIDA = 0.000 A

80% saturation of filtered PIDA = 0.001 A

Absorbance vs. Time Stored for Filtered $\text{PhI}(\text{OAc})_2$ **1a**

Filtered **1a** was stored as a solid at 2 °C in a vial. 60% saturated solutions in CH_2Cl_2 were made to measure the absorbance. Following measurement, the solvent was removed in vacuo and the obtained solid **1a** was added back into the vial it came from.

Date	Absorbance (stored under nitrogen atmosphere)	Absorbance (stored under air atmosphere)
07/25/22	0.000 A	0.000 A
07/26/22	0.000 A	0.000 A
07/27/22	0.000 A	0.000 A
07/28/22	0.000 A	0.000 A
08/01/22	0.000 A	0.000 A
08/02/22	0.001 A	0.001 A
08/03/22	0.000 A	0.001 A